arXiv:math/0412065v2 [math.NT] 16 Dec 2004

Hessian polyhedra, invariant theory and Appell hypergeometric partial differential equations

Lei Yang

Contents

1. Introduction
2. The reciprocity law about the Appell hypergeometric functions
3. The algebraic solutions of Appell hypergeometric partial differential equations
4. The Hessian polyhedral equations
5. Invariant theory for the system of algebraic equations
6. Ternary cubic forms associated to Hessian polyhedra
7. Some rational invariants on $\mathbb{CP}^2$

1. Introduction

Geometry is one of the oldest and most basic branches of mathematics, as is algebra. Nowhere is the interplay between the two more pronounced than in group theory, and that interplay with group theory acting as a mediator between geometry and algebra. It was Felix Klein in his "Erlanger Programme" who put group theory at the basis of geometry (see [KlE]). Klein advocated considering geometry as invariance properties under a group of automorphisms.

Polyhedra, particularly the Platonic solids, are the most classical objects in geometry, which have been studied by geometers for thousands of years. Furthermore, finding physical applications of polyhedra is a similarly ancient pursuit. For example, Plato was so captivated by the perfect forms of the five regular solids. Kepler also attributed cosmic significance to the Platonic solids (see [W]).

In his famous lecture: Mathematical problems (see [H]), which was delivered before the International Congress of Mathematicians at Paris in 1900, David Hilbert said: "It often happens that the same special problem finds application in the most unlike branches of mathematical knowledge. For example, how convincingly has Felix Klein, in his work on the icosahedron, pictured the significance which attaches to the problem of the regular polyhedra in elementary geometry, in group theory, in the theory of equations and in that of linear differential equations."

Typeset by $\mathcal{AMS}$-TEX

Klein's lectures on the icosahedron, published in 1884, soon became one of the most famous books on mathematics for a half-century; no other book so forcefully put in the limelight the fundamental unity of mathematics (see J. Dieudonné's review on P. Slodowy's paper [Sl] or see [At]). Klein showed that four apparently disjoint theories: the symmetries of the icosahedron (geometry), the resolution of fifth degree equations (algebra), the differential equation of hypergeometric functions (analysis) and the modular equations of elliptic modular functions (arithmetic) were in fact dominated by the structure of a single object, the simple group A_5 of 60 elements. It is the group of rotations leaving the icosahedron invariant, the monodromy group of a hypergeometric equation, and the commutator subgroup of the Galois group of the general fifth degree equation. Klein also pointed out the connection with invariant theory and algebraic geometry (see [Kl], [Kl9] and [Kl10], especially [Kl1], [Kl2], [Kl3], [Kl4], [Kl5], [Kl6], [Kl7] and [Kl8] for the details).

In a letter to J. D. Gray in response to his questions regarding modern treatments of Klein's fascinating but difficult book, J.-P. Serre said (see [Se]) that there is so much more in Klein's book, and in Fricke's! Invariants, hypergeometric functions, and everywhere, a wealth of beautiful formulae!

Now, let us recall some basic facts in Klein's lecture.

There is a long history for solving algebraic equations by means of transcendental functions (see [Kl9] or [G], [Hu]). Put

$$K=\int_0^{\frac{\pi}{2}}\frac{d\theta}{\sqrt{1-k^2\sin^2\theta}},\quad K'=\int_0^{\frac{\pi}{2}}\frac{d\theta}{\sqrt{1-k'^2\sin^2\theta}},\quad k^2+k'^2=1.$$

Let λ be a modulus related to k by a 5th order transformation. In 1829, Jacobi showed that $\sqrt[4]{k}=u$ and $\sqrt[4]{\lambda}=v$ are related by the modular equation (Jacobi's equation):

$$u^6-v^6+5u^2v^2(u^2-v^2)+4uv(1-u^4v^4)=0.$$

In 1858, Hermite (see [He], pp.5-12) found the resolvent of fifth degree which derives from Jacobi's equation of sixth degree. He did this as follows: set

$$y=(v_\infty-v_0)(v_1-v_4)(v_2-v_3),$$

where v_i are the roots of Jacobi's equation. The corresponding resolvent is

$$y^5-2^4\cdot 5^3u^4(1-u^8)^2y-2^6\cdot\sqrt{5^5}u^3(1-u^8)^2(1+u^8)=0.$$

This takes the form of a Bring equation

$$t^5-t-A=0$$

by setting

$$y = 2 \cdot \sqrt[4]{5^3} \cdot u\sqrt{1-u^8} \cdot t, \quad A = \frac{2}{\sqrt[4]{5^5}} \cdot \frac{1+u^8}{u^2\sqrt{1-u^8}}.$$

Hence, by a Tschirnhaus transformation any quintic equation can be brought into the Bring form, which can be solved by the elliptic functions for the transformations of fifth order, this gives a solution of the quintic equation.

The story of the general equation of fifth degree is ultimately culminating in Klein's book [Kl9]. Here geometry, analysis and algebra were brought to bear, showing not only how one can solve an equation of fifth degree, but also the relationships between the different methods which had been used previously. There is a universal Galois resolvent for general quintic equations: the icosahedral equation. Klein built the entire theory of quintic equations on these foundations. He showed that in studying this resolvent one can apply the geometry of the icosahedron, making the algebra very geometric. The solution suggested by Klein had an algebraic and a transcendental part. For the algebraic part, the classical approach is the reduction of equation with solvable Galois group to a pure equation:

$$x^n = X.$$

The icosahedral approach is the reduction of equation of fifth degree to solution of the icosahedral equation:

$$\frac{H^3(x)}{1728 f^5(x)} = X.$$

For the transcendental part, the classical approach is the solution of the pure equation by means of logarithms:

$$x = e^{\frac{1}{n}\log X}.$$

The icosahedral approach is the solution of the icosahedral equation by means of elliptic functions:

$$x = q^{\frac{2}{5}} \frac{\theta_1\left(\frac{2\pi i K'}{K}, q^5\right)}{\theta_1\left(\frac{\pi i K'}{K}, q^5\right)}.$$

In 1888, Klein ([Kl7]) sketched in a letter to Jordan how one could use a similar process to solve the equation of 27th degree determining the 27 lines on a smooth cubic surface. This leads to the development of the invariant theory of several finite collineation groups (unitary reflection groups in projective space). Moreover, this laid the grounds for a theory of arithmetic groups. Following suggestions of Klein and Jordan, Maschke ([Mas1], [Mas2]) and Burkhardt ([Bu1], [Bu2]) solved the equation of 27th degree for the lines of the cubic surface by using hyperelliptic functions. Here, the algebraic part is the reduction to the Maschke form problem:

$$\frac{F_{24}(z)}{F_{12}^2(z)} = \alpha, \quad \frac{F_{30}(z)}{F_{12}(z)F_{18}(z)} = \beta, \quad \frac{F_{18}^2(z)}{F_{12}^3(z)} = \gamma.$$

The transcendental part is the solution of the Maschke equation in terms of the hyperelliptic functions $Z_{\alpha,\beta}$ of

$$\mathbb{P}^3 = \left\{ Z_{\alpha,\beta} = \frac{1}{2}(X_{\alpha,\beta} - X_{-\alpha,-\beta}) \right\}.$$

In fact, thinking of each of the 27 lines as defining a point on the Grassmannian $\mathbf{G}(2,4) \subset \mathbb{P}^5$, one can project to 27 points onto a line, that is, giving the roots of a degree 27 equation. In 1890, Burkhardt showed that the equation could be solved involving the calculation of the periods of an abelian surface. The method, due to Klein and Jordan, carried out by Maschke and Burkhardt, uses the representation of G_{25920} in $\mathbb{P}^3$ and its invariants.

Later on Coble reconsidered the problem, essentially by considering the Galois group $W(E_6)$ not as a reflection group, but as a Cremona group, i.e., as a finite group of birational automorphisms of the plane $\mathbb{P}^2$ (see [Cob1], [Cob2], [Cob3], [Cob4] and [Cob5]). The approach of Coble uses the representation in $\mathbb{P}^4$ and reduces the problem to the form problem for this group.

In the general case, Umemura (see [U]) gave the resolution of algebraic equations by a Siegel modular function which is explicitly expressed by theta constants. For row vectors $n_1, m_2 \in \mathbb{R}^g$, $z \in \mathbb{C}^g$ and a symmetric $g \times g$ matrix τ with positive definite imaginary part, we define the theta function (see [Mu])

$$\theta \begin{bmatrix} m_1 \\ m_2 \end{bmatrix} (z, \tau) = \sum_{\xi \in \mathbb{Z}^g} e\left[\frac{1}{2}(\xi + m_1)\tau^t(\xi + m_1) + (\xi + m_1)^t(z + m_2) \right],$$

where $e(x) = \exp(2\pi i x)$. The theta constant $\theta \begin{bmatrix} m_1 \\ m_2 \end{bmatrix} (0, \tau)$ is denoted by $\theta \begin{bmatrix} m_1 \\ m_2 \end{bmatrix} (\tau)$.

Theorem (see [U]). *Let*

$$a_0 X^n + a_1 X^{n-1} + \cdots + a_n = 0, \quad a_0 \neq 0, a_i \in \mathbb{C}(0 \leq i \leq n)$$

be an algebraic equation irreducible over a certain subfield of $\mathbb{C}$, *then a root of it is given by*

$$\begin{aligned}
&(\theta \begin{bmatrix} \frac{1}{2} & 0 & \cdots & 0 \\ 0 & 0 & \cdots & 0 \end{bmatrix} (\Omega)^4 \theta \begin{bmatrix} \frac{1}{2} & \frac{1}{2} & 0 & \cdots & 0 \\ 0 & 0 & 0 & \cdots & 0 \end{bmatrix} (\Omega)^4 \\
&+ \theta \begin{bmatrix} 0 & \cdots & 0 \\ 0 & \cdots & 0 \end{bmatrix} (\Omega)^4 \theta \begin{bmatrix} 0 & \frac{1}{2} & 0 & \cdots & 0 \\ 0 & 0 & 0 & \cdots & 0 \end{bmatrix} (\Omega)^4 \\
&- \theta \begin{bmatrix} 0 & 0 & \cdots & 0 \\ \frac{1}{2} & 0 & \cdots & 0 \end{bmatrix} (\Omega)^4 \theta \begin{bmatrix} 0 & \frac{1}{2} & 0 & \cdots & 0 \\ \frac{1}{2} & 0 & 0 & \cdots & 0 \end{bmatrix} (\Omega)^4) \\
&\times (2\theta \begin{bmatrix} \frac{1}{2} & 0 & \cdots & 0 \\ 0 & 0 & \cdots & 0 \end{bmatrix} (\Omega)^4 \theta \begin{bmatrix} \frac{1}{2} & \frac{1}{2} & 0 & \cdots & 0 \\ 0 & 0 & 0 & \cdots & 0 \end{bmatrix} (\Omega)^4)^{-1},
\end{aligned}$$

where Ω is the period matrix of a hyperelliptic curve $C : Y^2 = F(X)$ with

$$F(X) = X(X-1)(a_0X^n + a_1X^{n-1} + \cdots + a_n)$$

for n odd and

$$F(X) = X(X-1)(X-2)(a_0X^n + a_1X^{n-1} + \cdots + a_n)$$

for n even.

Let us recall some basic facts about the icosahedral group (see [Kl9] or [F], [We]), which is generated by the following homogeneous substitutions:

$$S : \begin{cases} z_1' = \pm\epsilon^3 z_1, \\ z_2' = \pm\epsilon^2 z_2, \end{cases}$$

$$U : \begin{cases} z_1' = \mp z_2, \\ z_2' = \pm z_1, \end{cases}$$

$$T : \begin{cases} \sqrt{5} \cdot z_1' = \mp(\epsilon - \epsilon^4) z_1 \pm (\epsilon^2 - \epsilon^3) z_2, \\ \sqrt{5} \cdot z_2' = \pm(\epsilon^2 - \epsilon^3) z_1 \pm (\epsilon - \epsilon^4) z_2, \end{cases}$$

where

$$\epsilon = e^{\frac{2\pi i}{5}}.$$

In fact, U can be generated by S and T (see [F]).

The set of forms for the icosahedron is given by

$$f = z_1 z_2 (z_1^{10} + 11 z_1^5 z_2^5 - z_2^{10});$$

$$H = \frac{1}{121} \begin{vmatrix} \frac{\partial^2 f}{\partial z_1^2} & \frac{\partial^2 f}{\partial z_1 \partial z_2} \\ \frac{\partial^2 f}{\partial z_2 \partial z_1} & \frac{\partial^2 f}{\partial z_2^2} \end{vmatrix} = -(z_1^{20} + z_2^{20}) + 228(z_1^{15} z_2^5 - z_1^5 z_2^{15}) - 494 z_1^{10} z_2^{10};$$

$$T = -\frac{1}{20} \begin{vmatrix} \frac{\partial f}{\partial z_1} & \frac{\partial f}{\partial z_2} \\ \frac{\partial H}{\partial z_1} & \frac{\partial H}{\partial z_2} \end{vmatrix} = (z_1^{30} + z_2^{30}) + 522(z_1^{25} z_2^5 - z_1^5 z_2^{25}) - 10005(z_1^{20} z_2^{10} + z_1^{10} z_2^{20}).$$

In fact, $f = 0$ represents the 12 summits of the icosahedron, $H = 0$ represents the 20 summits of the pentagon-dodecahedron and $T = 0$ represents the 30 mid-edge points. Moreover, f, H and T satisfy the identity:

$$T^2 = -H^3 + 1728 f^5.$$

The icosahedral equation is defined by

$$Z = \frac{H^3}{1728 f^5}.$$

In his paper [Kl11], Klein studied the icosahedral irrationality η defined by the equation:

$$\frac{H^3(\eta)}{1728 f^5(\eta)} = J,$$

where J is Klein's J-invariant in the theory of elliptic modular functions.

Let the octahedron be the form:

$$t = z_1^6 + 2z_1^5 z_2 - 5z_1^4 z_2^2 - 5z_1^2 z_2^4 - 2z_1 z_2^5 + z_2^6.$$

The Hessian form of t is

$$W = -z_1^8 + z_1^7 z_2 - 7z_1^6 z_2^2 - 7z_1^5 z_2^3 + 7z_1^3 z_2^5 - 7z_1^2 z_2^6 - z_1 z_2^7 - z_2^8.$$

Put

$$r = \frac{t^2}{f},$$

this gives the function-theoretic resolvent (analytic resolvent):

$$Z : (Z-1) : 1 = (r-3)^3 (r^2 - 11r + 64) : r(r^2 - 10r + 45)^2 : -1728.$$

In fact, set

$$\xi = \frac{125}{z^5 - z^{-5} + 11} = \frac{125 z_1^6 z_2^6}{f(z_1, z_2)},$$

we have (see [F])

$$Z : (Z-1) : 1 = (\xi^2 - 10\xi + 5)^3 : (\xi^2 - 22\xi + 125)(\xi^2 - 4\xi - 1)^2 : -1728\xi,$$

which corresponds to the transformation of fifth order for elliptic functions (see [Kl11]):

$$J : (J-1) : 1 = (\tau^2 - 10\tau + 5)^3 : (\tau^2 - 22\tau + 125)(\tau^2 - 4\tau - 1)^2 : -1728\tau.$$

On the other hand, the form-theoretic resolvent (algebraic resolvent) of t is:

$$t^5 - 10f \cdot t^3 + 45f^2 \cdot t - T = 0.$$

The form-theoretic resolvent (algebraic resolvent) of W is:

$$W^5 + 40f^2 \cdot W^2 - 5fH \cdot W + H^2 = 0.$$

Let

$$u = \frac{12tf^2}{T}, \quad v = \frac{12Wf}{H}.$$

The analytic and algebraic resolvent of u is

$$48u^5(1-Z)^2 - 40u^3(1-Z) + 15u - 4 = 0.$$

The analytic and algebraic resolvent of v is

$$Zv^5 + 40v^2 - 60v + 144 = 0.$$

Furthermore,

$$u = \frac{12}{r^2 - 10r + 45}, \quad v = \frac{12}{r-3}.$$

Put

$$\begin{cases} \phi_\infty = 5z_1^2 z_2^2, \\ \phi_\nu = (\epsilon^\nu z_1^2 + z_1 z_2 - \epsilon^{4\nu} z_2^2)^2, \quad \nu = 0, 1, 2, 3, 4. \end{cases}$$

The form-theoretic resolvent (algebraic resolvent) of ϕ is

$$\phi^6 - 10f \cdot \phi^3 + H \cdot \phi + 5f^2 = 0.$$

Set

$$\zeta = \frac{\phi H}{12f^2}.$$

Then the analytic and algebraic resolvent of ζ is:

$$\zeta^6 - 10Z \cdot \zeta^3 + 12Z^2 \cdot \zeta + 5Z^2 = 0.$$

Klein set

$$\mathbf{A}_0 = z_1 z_2, \quad \mathbf{A}_1 = z_1^2, \quad \mathbf{A}_2 = -z_2^2$$

and

$$A = \mathbf{A}_0^2 + \mathbf{A}_1 \mathbf{A}_2.$$

The invariants f, H and T can be expressed as the functions of $\mathbf{A}_0$, $\mathbf{A}_1$ and $\mathbf{A}_2$:

$$B = 8\mathbf{A}_0^4 \mathbf{A}_1 \mathbf{A}_2 - 2\mathbf{A}_0^2 \mathbf{A}_1^2 \mathbf{A}_2^2 + \mathbf{A}_1^3 \mathbf{A}_2^3 - \mathbf{A}_0(\mathbf{A}_1^5 + \mathbf{A}_2^5),$$

$$C=320\mathbf{A}_0^6\mathbf{A}_1^2\mathbf{A}_2^2-160\mathbf{A}_0^4\mathbf{A}_1^3\mathbf{A}_2^3+20\mathbf{A}_0^2\mathbf{A}_1^4\mathbf{A}_2^4+6\mathbf{A}_1^5\mathbf{A}_2^5+$$
$$-4\mathbf{A}_0(\mathbf{A}_1^5+\mathbf{A}_2^5)(32\mathbf{A}_0^4-20\mathbf{A}_0^2\mathbf{A}_1\mathbf{A}_2+5\mathbf{A}_1^2\mathbf{A}_2^2)+\mathbf{A}_1^{10}+\mathbf{A}_2^{10},$$

$$D=(\mathbf{A}_1^5-\mathbf{A}_2^5)[-1024\mathbf{A}_0^{10}+3840\mathbf{A}_0^8\mathbf{A}_1\mathbf{A}_2-3840\mathbf{A}_0^6\mathbf{A}_1^2\mathbf{A}_2^2+$$
$$+1200\mathbf{A}_0^4\mathbf{A}_1^3\mathbf{A}_2^3-100\mathbf{A}_0^2\mathbf{A}_1^4\mathbf{A}_2^4+\mathbf{A}_1^{10}+\mathbf{A}_2^{10}+2\mathbf{A}_1^5\mathbf{A}_2^5+$$
$$+\mathbf{A}_0(\mathbf{A}_1^5+\mathbf{A}_2^5)(352\mathbf{A}_0^4-160\mathbf{A}_0^2\mathbf{A}_1\mathbf{A}_2+10\mathbf{A}_1^2\mathbf{A}_2^2)].$$

They satisfy the identity

$$D^2=-1728B^5+C^3+720ACB^3-80A^2C^2B+64A^3(5B^2-AC)^2.$$

The actions of the icosahedral group on $\mathbf{A}_0$, $\mathbf{A}_1$ and $\mathbf{A}_2$ are given by

$$S:\quad \mathbf{A}_0'=\mathbf{A}_0,\quad \mathbf{A}_1'=\epsilon\mathbf{A}_1,\quad \mathbf{A}_2'=\epsilon^4\mathbf{A}_2;$$

$$T:\begin{cases}\sqrt{5}\mathbf{A}_0'=\mathbf{A}_0+\mathbf{A}_1+\mathbf{A}_2,\\ \sqrt{5}\mathbf{A}_1'=2\mathbf{A}_0+(\epsilon^2+\epsilon^3)\mathbf{A}_1+(\epsilon+\epsilon^4)\mathbf{A}_2,\\ \sqrt{5}\mathbf{A}_2'=2\mathbf{A}_0+(\epsilon+\epsilon^4)\mathbf{A}_1+(\epsilon^2+\epsilon^3)\mathbf{A}_2;\end{cases}$$

$$U:\quad \mathbf{A}_0'=-\mathbf{A}_0,\quad \mathbf{A}_1'=-\mathbf{A}_2,\quad \mathbf{A}_2'=-\mathbf{A}_1.$$

This leads to the investigation of the binary quadratic forms:

$$\mathbf{A}_1z_1^2+2\mathbf{A}_0z_1z_2-\mathbf{A}_2z_2^2$$

and Hilbert modular surfaces (see [Hi]).

It is well-known that the group A_5 is isomorphic to the finite group I of those elements of $SO(3)$ which carry a given icosahedron centered at the origin of the standard Euclidean space $\mathbb{R}^3$ to itself. The group I operates linearly on $\mathbb{R}^3$ (standard coordinates x_0, x_1, x_2) and thus also on $\mathbb{RP}^2$ and $\mathbb{CP}^2$. We are concerned with the action on $\mathbb{CP}^2$ (see [Hi]). A curve in $\mathbb{CP}^2$ which is mapped to itself by all elements of I is given by a homogeneous polynomial in x_0, x_1, x_2 which is I-invariant up to constant factors and hence I-invariant, because I is a simple group. The graded ring of all I-invariant polynomials in x_0, x_1, x_2 is generated by homogeneous polynomials A, B, C, D of degrees 2, 6, 10, 15 with $A=x_0^2+x_1^2+x_2^2$. The action of I on $\mathbb{CP}^2$ has exactly one minimal orbit where "minimal" means that the number of points in the orbit is minimal. This orbit has six points, they are called poles. These are the points of $\mathbb{RP}^2\subset\mathbb{CP}^2$ which are represented by the six axes through the vertices of the icosahedron. Klein uses coordinates

$$A_0=x_0,\quad A_1=x_1+ix_2,\quad A_2=x_1-ix_2$$

and puts the icosahedron in such a position that the six poles are given by

$$(A_0, A_1, A_2) = \left(\frac{\sqrt{5}}{2}, 0, 0\right), \quad \left(\frac{1}{2}, \varepsilon^{\nu}, \varepsilon^{-\nu}\right)$$

with $\varepsilon = \exp(2\pi i/5)$ and $0 \leq \nu \leq 4$.

The invariant curve $A = 0$ does not pass through the poles. There is exactly one invariant curve $B = 0$ of degree 6 which passes through the poles, exactly one invariant curve $C = 0$ of degree 10 which has higher multiplicity than the curve $B = 0$ in the poles and exactly one invariant curve $D = 0$ of degree 15. In fact, $B = 0$ has an ordinary double point (multiplicity 2) in each pole, $C = 0$ has a double cusp (multiplicity 4) in each pole and $D = 0$ is the union of the 15 lines connecting the six poles. Klein gives formulas for the homogeneous polynomials A, B, C, D (determined up to constant factors). They generate the ring of all I-invariant polynomials. According to Klein the ring of I-invariant polynomials is given as follows:

$$\mathbb{C}[A_0, A_1, A_2]^I = \mathbb{C}[A, B, C, D]/(R(A, B, C, D) = 0).$$

The relation $R(A, B, C, D) = 0$ is of degree 30.

The equations for B and C show that the two tangents of $B = 0$ in the pole $(\sqrt{5}/2, 0, 0)$ are given by $A_1 = 0$, $A_2 = 0$. They coincide with the tangents of $C = 0$ in that pole. Therefore the curves $B = 0$ and $C = 0$ have in each pole the intersection multiplicity 10. Thus they intersect only in the poles.

When we restrict the action of I to the conic $A = 0$, we get the well-known action of I on $\mathbb{CP}^1$. The curves $B = 0$, $C = 0$, $D = 0$ intersect $A = 0$ transversally in 12, 20, 30 points respectively.

It is well-known that Klein's lectures ([Kl9]) on the icosahedron and the solution of equations of fifth degree is one of the most important and influential books of 19th-century mathematics. Since the time (1884) of the publication of Klein's lectures, there has been a standing interest in the mathematics of the icosahedron, i.e. the mathematics where the geometry and the symmetry of the icosahedron play an important role (see [KlS]).

This interest has increased in the last few decades. Let us mention some recent developments: the study of the Klein singularities (Arnold and Brieskorn), the study of the Hilbert modular surfaces (Hirzebruch), the connection between the platonic solids (or the finite subgroups of $SU(2)$) and the complex, simple Lie groups of the type A_r, D_r, E_6, E_7, E_8, discovered by Grothendieck and Brieskorn, the connection between singularities and string duality in the M-theory (Witten, Vafa), the study of the theory of Dessins d'Enfants called by its founder Grothendieck (see [Gro]).

From the point of view of representation theory, by the regular polyhedra, we can obtain dihedral, tetrahedral, octahedral and icosahedral representations. These lead to

Artin's conjecture and Langlands programme (see [L1] and [L2]). Among them, the icosahedral equations remain intractable in general. The equations with icosahedral Galois groups has a greater historical appeal since they are the simplest equations completely inaccessible to the classical theory of equations with abelian Galois groups.

These results have shown how the icosahedron and the platonic solids have thrown new light in a surprising way on previously separate developments.

Recently, polyhedra, which include the Platonic solids, were found to arise naturally in a number of diverse problems in physics, chemistry, biology and a variety of other disciplines. In his book [W], Weyl gave a general discussion of symmetry in the natural world. Atiyah and Sutcliffe ([AtS]) gave some examples which appear in physics, chemistry and mathematics, such as the study of electrons on a sphere, cages of carbon atoms, central configurations of gravitating point particles, rare gas microclusters, soliton models of nuclei (the Skyrme model), magnetic monopole scattering and geometrical problems concerning point particles. Moreover, a particularly interesting application of polyhedra in biology is provided by the structure of spherical virus shells, such as HIV which is built around a trivalent polyhedron with icosahedral symmetry.

In the present paper, we will give the complex counterpart of Klein's book [Kl9], i.e., a story about complex regular polyhedra (which were also called regular pseudo-polyhedra defined over $\mathbb{C}$ by Grothendieck in [Gro]). We will show that the following four apparently disjoint theories: the symmetries of the Hessian polyhedra (geometry), the resolution of some system of algebraic equations (algebra), the system of partial differential equations of Appell hypergeometric functions (analysis) and the modular equation of Picard modular functions (arithmetic) are in fact dominated by the structure of a single object, the Hessian group $\mathfrak{G}'_{216}$. There are two finite unitary groups generated by reflections corresponding to this collineation group. One, of order 648, is the symmetry group of the complex regular polyhedron $3\{3\}3\{3\}3$, and the other, of order 1296, is the symmetry group of the regular complex polyhedron $2\{4\}3\{3\}3$ or its reciprocal $3\{3\}3\{4\}2$.

The significance of the Hessian polyhedron comes from the sequence of polytopes (see [Co2])

$$3\{3\}3, \quad 3\{3\}3\{3\}3, \quad 3\{3\}3\{3\}3\{3\}3$$

(having 8, 27, 240 vertices) which culminates in the four-dimensional honeycomb

$$3\{3\}3\{3\}3\{3\}3\{3\}3.$$

These are very important because of their connection with the celebrated configuration of 27 lines on the general cubic surface in projective 3-space. In fact, the binary tetrahedral group $3[3]3$ is a subgroup of index 9 in the Hessian group of order 216, which is a factor group of the symmetry group $3[3]3[3]3$ of the Hessian polyhedron $3\{3\}3\{3\}3$; this group of order 648 is a subgroup of index 40 in the simple group of order 25920 which is a factor group of the symmetry group $3[3]3[3]3[3]3$ of the Witting polytope $3\{3\}3\{3\}3\{3\}3$.

There are two fundamental reasons for the significance of the triple $W(E_6)$, G_{25920} and G_{648}:

(1) $W(E_6)$ is a reflection group in $\mathbb{P}^5$, G_{25920} is a unitary reflection group in $\mathbb{P}^4$ and $\mathbb{P}^3$, G_{648} is a unitary reflection group in $\mathbb{P}^2$.
(2) $W(E_6)$ is the Galois group of the algebraic equation for the 27 lines on the cubic surface, G_{25920} is the Galois group of the algebraic equation which results after extracting the discriminant of the above, G_{648} is the Galois group of our Hessian polyhedral equations.

In the theory of algebraic curves, if one wants to go beyond the category of hyperelliptic curves, the first non-hyperelliptic curves are plane quartic curves of genus three, such as Klein quartic curve (see [Kl12]), Fermat quartic curve $x^4 + y^4 = 1$ and Picard curves (see [Ho]). Just as the hyperelliptic modular functions (Siegel modular functions) correspond to the general algebraic equation (one variable), we find that the Picard modular functions correspond to some system of algebraic equations (two variables).

In fact, we have the following dictionaries about algebraic equations, groups, integrals of algebraic functions, genus of algebraic curves and transcendental functions (automorphic functions):

(1)

$GL(1)$,
pure equation, $GL(1)$,
rational integrals, rational functions, $g = 0$,
exponential functions,
rational curves.

(2)

$GL(2), SL(2, \mathbb{R})$,
icosahedral equation, quintic equation, $GL(2)$,
elliptic integrals, elliptic functions, $g = 1$,
elliptic modular functions, $SL(2, \mathbb{Z})$,
Dedekind η-function,
Klein J-invariant,
elliptic curves, $SL(2, \mathbb{Z})(2)$,
$y^2 = x(x-1)(x-\lambda)$,
icosahedron, $PSL(2, \mathbb{F}_4) \cong PSL(2, \mathbb{F}_5) \cong A_5$.

(3)

$GL(4), Sp(4, \mathbb{R})$,
27 lines on the cubic surfaces,
hyperelliptic integrals, hyperelliptic functions, $g = 2$,
Siegel modular functions of genus two, $Sp(4, \mathbb{Z})$,
Igusa's J-invariants J_2, J_4, J_6, J_8 and J_{10} (see [I]),
hyperelliptic curves of genus two, $Sp(4, \mathbb{Z})(2)$,
$y^2 = x(x-1)(x-\lambda_1)(x-\lambda_2)(x-\lambda_3)$,
$PSp(4, \mathbb{F}_3)$, order 25920,
Witting polytope, 3[3]3[3]3[3]3.

(4)

$Sp(2g, \mathbb{R}), g > 2$
general algebraic equation,
hyperelliptic integrals of genus $g > 2$,
Siegel modular functions of genus $g > 2$,
hyperelliptic curves of genus $g > 2$.

(5)

$GL(3), U(2,1)$,
Hessian polyhedral equations, system of algebraic equations, $GL(3)$,
Picard integrals, $g = 3$,
Picard modular functions, $U(2, 1; \mathbb{Z}[\omega])$,
our η-function (see [Y]),
our J-invariants J_1 and J_2 (see [Y]),
Picard curves, $U(2, 1; \mathbb{Z}[\omega])(1-\omega)$,
$y^3 = x(x-1)(x-\lambda_1)(x-\lambda_2)$,
Hessian polyhedra, 3[3]3[3]3, 2[4]3[3]3, 3[3]3[4]2.

Main Results.

The Reciprocity Law about the Appell hypergeometric functions
The Appell hypergeometric partial differential equations

$$\left\{\begin{array}{l} x(1-x)\frac{\partial^2 z}{\partial x^2}+y(1-x)\frac{\partial^2 z}{\partial x \partial y}+[c-(a+b+1)x]\frac{\partial z}{\partial x}-by\frac{\partial z}{\partial y}-abz=0, \\ y(1-y)\frac{\partial^2 z}{\partial y^2}+x(1-y)\frac{\partial^2 z}{\partial x \partial y}+[c-(a+b'+1)y]\frac{\partial z}{\partial y}-b'x\frac{\partial z}{\partial x}-ab'z=0, \end{array}\right. \tag{1.1}$$

are equivalent to the following three equations:

(1.2)

$$\begin{cases}\dfrac{\partial^2 z}{\partial x^2}+\left(\dfrac{c-b'}{x}+\dfrac{a+b-c+1}{x-1}+\dfrac{b'}{x-y}\right)\dfrac{\partial z}{\partial x}-\dfrac{by(y-1)}{x(x-1)(x-y)}\dfrac{\partial z}{\partial y}+\dfrac{abz}{x(x-1)}=0,\\ \dfrac{\partial^2 z}{\partial y^2}+\left(\dfrac{c-b}{y}+\dfrac{a+b'-c+1}{y-1}+\dfrac{b}{y-x}\right)\dfrac{\partial z}{\partial y}-\dfrac{b'x(x-1)}{y(y-1)(y-x)}\dfrac{\partial z}{\partial x}+\dfrac{ab'z}{y(y-1)}=0,\\ \dfrac{\partial^2 z}{\partial x\partial y}=\dfrac{b'}{x-y}\dfrac{\partial z}{\partial x}-\dfrac{b}{x-y}\dfrac{\partial z}{\partial y}.\end{cases}$$

There are three linearly independent solutions z_1, z_2 and z_3. Set $u_1=\frac{z_1}{z_3}$ and $u_2=\frac{z_2}{z_3}$.

According to [Y], we define the following four derivatives associated to the group $GL(3)$:

$$\{u_1,u_2;x,y\}_x:=\frac{\begin{vmatrix}\frac{\partial u_1}{\partial x} & \frac{\partial u_2}{\partial x}\\ \frac{\partial^2 u_1}{\partial x^2} & \frac{\partial^2 u_2}{\partial x^2}\end{vmatrix}}{\begin{vmatrix}\frac{\partial u_1}{\partial x} & \frac{\partial u_2}{\partial x}\\ \frac{\partial u_1}{\partial y} & \frac{\partial u_2}{\partial y}\end{vmatrix}},\qquad \{u_1,u_2;x,y\}_y:=\frac{\begin{vmatrix}\frac{\partial u_1}{\partial y} & \frac{\partial u_2}{\partial y}\\ \frac{\partial^2 u_1}{\partial y^2} & \frac{\partial^2 u_2}{\partial y^2}\end{vmatrix}}{\begin{vmatrix}\frac{\partial u_1}{\partial y} & \frac{\partial u_2}{\partial y}\\ \frac{\partial u_1}{\partial x} & \frac{\partial u_2}{\partial x}\end{vmatrix}},$$

$$(1.3)\qquad [u_1,u_2;x,y]_x:=\frac{\begin{vmatrix}\frac{\partial u_1}{\partial y} & \frac{\partial u_2}{\partial y}\\ \frac{\partial^2 u_1}{\partial x^2} & \frac{\partial^2 u_2}{\partial x^2}\end{vmatrix}+2\begin{vmatrix}\frac{\partial u_1}{\partial x} & \frac{\partial u_2}{\partial x}\\ \frac{\partial^2 u_1}{\partial x\partial y} & \frac{\partial^2 u_2}{\partial x\partial y}\end{vmatrix}}{\begin{vmatrix}\frac{\partial u_1}{\partial x} & \frac{\partial u_2}{\partial x}\\ \frac{\partial u_1}{\partial y} & \frac{\partial u_2}{\partial y}\end{vmatrix}},$$

$$[u_1,u_2;x,y]_y:=\frac{\begin{vmatrix}\frac{\partial u_1}{\partial x} & \frac{\partial u_2}{\partial x}\\ \frac{\partial^2 u_1}{\partial y^2} & \frac{\partial^2 u_2}{\partial y^2}\end{vmatrix}+2\begin{vmatrix}\frac{\partial u_1}{\partial y} & \frac{\partial u_2}{\partial y}\\ \frac{\partial^2 u_1}{\partial x\partial y} & \frac{\partial^2 u_2}{\partial x\partial y}\end{vmatrix}}{\begin{vmatrix}\frac{\partial u_1}{\partial y} & \frac{\partial u_2}{\partial y}\\ \frac{\partial u_1}{\partial x} & \frac{\partial u_2}{\partial x}\end{vmatrix}},$$

where x and y are complex numbers, u_1 and u_2 are complex-valued functions of x and y.

We find that (see [Y])

$$(1.4)\qquad \begin{cases}\{u_1,u_2;x,y\}_x=\dfrac{by(y-1)}{x(x-1)(x-y)},\\ \{u_1,u_2;x,y\}_y=\dfrac{b'x(x-1)}{y(y-1)(y-x)},\\ [u_1,u_2;x,y]_x=\dfrac{c-b'}{x}+\dfrac{a+b-c+1}{x-1}+\dfrac{b'-2b}{x-y},\\ [u_1,u_2;x,y]_y=\dfrac{c-b}{y}+\dfrac{a+b'-c+1}{y-1}+\dfrac{b-2b'}{y-x}.\end{cases}$$

THEOREM (see [Y], Main Theorem 1). *For the over-determined system of second order, nonlinear (quasilinear) complex partial differential equations*:

$$\left\{\begin{aligned} \{w_1,w_2;v_1,v_2\}_{v_1} &= F_1(v_1,v_2),\\ \{w_1,w_2;v_1,v_2\}_{v_2} &= F_2(v_1,v_2),\\ [w_1,w_2;v_1,v_2]_{v_1} &= P_1(v_1,v_2),\\ [w_1,w_2;v_1,v_2]_{v_2} &= P_2(v_1,v_2), \end{aligned}\right. \tag{1.5}$$

set the Jacobian

$$\Delta := \frac{\partial(v_1,v_2)}{\partial(w_1,w_2)} = \begin{vmatrix} \frac{\partial v_1}{\partial w_1} & \frac{\partial v_2}{\partial w_1} \\ \frac{\partial v_1}{\partial w_2} & \frac{\partial v_2}{\partial w_2} \end{vmatrix}. \tag{1.6}$$

Then $z=\sqrt[3]{\Delta}$ *satisfies the following linear differential equations*:

$$\left\{\begin{aligned} \frac{\partial^2 z}{\partial v_1^2}+\frac{1}{3}P_1\frac{\partial z}{\partial v_1}-F_1\frac{\partial z}{\partial v_2}+\left(\frac{\partial F_1}{\partial v_2}-\frac{1}{3}\frac{\partial P_1}{\partial v_1}-\frac{2}{9}P_1^2-\frac{2}{3}F_1P_2\right)z=0,\\ \frac{\partial^2 z}{\partial v_1\partial v_2}-\frac{1}{3}P_2\frac{\partial z}{\partial v_1}-\frac{1}{3}P_1\frac{\partial z}{\partial v_2}+\left(\frac{1}{3}\frac{\partial P_2}{\partial v_1}+\frac{1}{3}\frac{\partial P_1}{\partial v_2}+\frac{1}{9}P_1P_2-F_1F_2\right)z=0,\\ \frac{\partial^2 z}{\partial v_2^2}+\frac{1}{3}P_2\frac{\partial z}{\partial v_2}-F_2\frac{\partial z}{\partial v_1}+\left(\frac{\partial F_2}{\partial v_1}-\frac{1}{3}\frac{\partial P_2}{\partial v_2}-\frac{2}{9}P_2^2-\frac{2}{3}F_2P_1\right)z=0. \end{aligned}\right. \tag{1.7}$$

THEOREM 1.1 (MAIN THEOREM 1). *For the system of linear partial differential equations*:

$$\left\{\begin{aligned} \frac{\partial^2 z}{\partial v_1^2}+\frac{1}{3}P_1\frac{\partial z}{\partial v_1}-F_1\frac{\partial z}{\partial v_2}+\left(\frac{\partial F_1}{\partial v_2}-\frac{1}{3}\frac{\partial P_1}{\partial v_1}-\frac{2}{9}P_1^2-\frac{2}{3}F_1P_2\right)z=0,\\ \frac{\partial^2 z}{\partial v_1\partial v_2}-\frac{1}{3}P_2\frac{\partial z}{\partial v_1}-\frac{1}{3}P_1\frac{\partial z}{\partial v_2}+\left(\frac{1}{3}\frac{\partial P_2}{\partial v_1}+\frac{1}{3}\frac{\partial P_1}{\partial v_2}+\frac{1}{9}P_1P_2-F_1F_2\right)z=0,\\ \frac{\partial^2 z}{\partial v_2^2}+\frac{1}{3}P_2\frac{\partial z}{\partial v_2}-F_2\frac{\partial z}{\partial v_1}+\left(\frac{\partial F_2}{\partial v_1}-\frac{1}{3}\frac{\partial P_2}{\partial v_2}-\frac{2}{9}P_2^2-\frac{2}{3}F_2P_1\right)z=0, \end{aligned}\right.$$

put

$$\left\{\begin{aligned} P_1 &= \frac{c-b'}{v_1}+\frac{a+b-c+1}{v_1-1}+\frac{b'-2b}{v_1-v_2},\\ P_2 &= \frac{c-b}{v_2}+\frac{a+b'-c+1}{v_2-1}+\frac{b-2b'}{v_2-v_1},\\ F_1 &= \frac{bv_2(v_2-1)}{v_1(v_1-1)(v_1-v_2)},\\ F_2 &= \frac{b'v_1(v_1-1)}{v_2(v_2-1)(v_2-v_1)}. \end{aligned}\right. \tag{1.8}$$

Set

$$(1.9)\quad w=v_1^{-\frac{1}{3}(c-b')}v_2^{-\frac{1}{3}(c-b)}(v_1-1)^{-\frac{1}{3}(a+b-c+1)}(v_2-1)^{-\frac{1}{3}(a+b'-c+1)}(v_1-v_2)^{-\frac{1}{3}(b+b')}z,$$

then the function $w=w(v_1,v_2)$ *satisfies the following equations*:

$$(1.10)\quad \begin{cases} \dfrac{\partial^2 w}{\partial v_1^2}+\left(\dfrac{c-b'}{v_1}+\dfrac{a+b-c+1}{v_1-1}+\dfrac{b'}{v_1-v_2}\right)\dfrac{\partial w}{\partial v_1}-\dfrac{bv_2(v_2-1)}{v_1(v_1-1)(v_1-v_2)}\dfrac{\partial w}{\partial v_2}+ \\ +\dfrac{ab}{v_1(v_1-1)}w=0, \\ \dfrac{\partial^2 w}{\partial v_2^2}+\left(\dfrac{c-b}{v_2}+\dfrac{a+b'-c+1}{v_2-1}+\dfrac{b}{v_2-v_1}\right)\dfrac{\partial w}{\partial v_2}-\dfrac{b'v_1(v_1-1)}{v_2(v_2-1)(v_2-v_1)}\dfrac{\partial w}{\partial v_1}+ \\ +\dfrac{ab'}{v_2(v_2-1)}w=0, \\ \dfrac{\partial^2 w}{\partial v_1\partial v_2}=\dfrac{b'}{v_1-v_2}\dfrac{\partial w}{\partial v_1}-\dfrac{b}{v_1-v_2}\dfrac{\partial w}{\partial v_2}. \end{cases}$$

A solution is given by

$$(1.11)\qquad w=F_1(a;b,b';c;v_1,v_2),$$

where $F_1=F_1(a;b,b';c;x,y)$ *is the Appell hypergeometric function* (*see* [AK]). *Consequently,*

$$(1.12)\quad \begin{aligned} z=&v_1^{\frac{1}{3}(c-b')}v_2^{\frac{1}{3}(c-b)}(v_1-1)^{\frac{1}{3}(a+b-c+1)}(v_2-1)^{\frac{1}{3}(a+b'-c+1)}(v_1-v_2)^{\frac{1}{3}(b+b')}\times \\ &\times F_1(a;b,b';c;v_1,v_2). \end{aligned}$$

In fact, the above reciprocity law is valid not only for the Appell hypergeometric partial differential equations, but also for the system of nonlinear partial differential equations involving our four derivatives (1.4).

A complete system of invariants for the Hessian groups (the corresponding geometric objects are Hessian polyhedra) has degrees 6, 9, 12, 12 and 18 and can be given explicitly by the following forms:

$$(1.13)\quad \begin{cases} C_6(z_1,z_2,z_3)=z_1^6+z_2^6+z_3^6-10(z_1^3z_2^3+z_2^3z_3^3+z_3^3z_1^3), \\ C_9(z_1,z_2,z_3)=(z_1^3-z_2^3)(z_2^3-z_3^3)(z_3^3-z_1^3), \\ C_{12}(z_1,z_2,z_3)=(z_1^3+z_2^3+z_3^3)[(z_1^3+z_2^3+z_3^3)^3+216z_1^3z_2^3z_3^3], \\ \mathfrak{C}_{12}(z_1,z_2,z_3)=z_1z_2z_3[27z_1^3z_2^3z_3^3-(z_1^3+z_2^3+z_3^3)^3], \\ C_{18}(z_1,z_2,z_3)=(z_1^3+z_2^3+z_3^3)^6-540z_1^3z_2^3z_3^3(z_1^3+z_2^3+z_3^3)^3-5832z_1^6z_2^6z_3^6. \end{cases}$$

They satisfy the following relations:

$$\left\{\begin{aligned} 432C_9^2 &= C_6^3 - 3C_6C_{12} + 2C_{18}, \\ 1728\mathfrak{C}_{12}^3 &= C_{18}^2 - C_{12}^3. \end{aligned}\right. \tag{1.14}$$

For the Hessian group 3[3]3[3]3, of order 6 × 9 × 12, the three forms are C_6, C_9, C_{12}. For the closely related Hessian group 2[4]3[3]3, of order 6 × 12 × 18, they are C_6, C_{12}, C_9^2 (see [Co2]).

In the affine coordinates $\xi = z_1/z_3$ and $\eta = z_2/z_3$, we have

$$\left\{\begin{aligned} C_6 &= C_6(\xi,\eta) = \xi^6 + \eta^6 + 1 - 10(\xi^3\eta^3 + \xi^3 + \eta^3), \\ C_9 &= C_9(\xi,\eta) = (\xi^3 - \eta^3)(\eta^3 - 1)(1 - \xi^3), \\ C_{12} &= C_{12}(\xi,\eta) = (\xi^3 + \eta^3 + 1)[(\xi^3 + \eta^3 + 1)^3 + 216\xi^3\eta^3], \\ \mathfrak{C}_{12} &= \mathfrak{C}_{12}(\xi,\eta) = \xi\eta[27\xi^3\eta^3 - (\xi^3 + \eta^3 + 1)^3], \\ C_{18} &= C_{18}(\xi,\eta) = (\xi^3 + \eta^3 + 1)^6 - 540\xi^3\eta^3(\xi^3 + \eta^3 + 1)^3 - 5832\xi^6\eta^6. \end{aligned}\right. \tag{1.15}$$

In [Y], we gave the modular equation associated to the Picard curves:

Theorem (see [Y], Main Theorem 3). *The following identity of differential forms holds*:

$$\frac{dw_1 \wedge dw_2}{\sqrt[3]{w_1^2 w_2^2 (1 - w_1)(1 - \lambda_1^3 w_1)(1 - \lambda_2^3 w_1)}} = -\frac{5t_1}{t_2^2} \frac{dt_1 \wedge dt_2}{\sqrt[3]{t_1^2 t_2^2 (1 - t_1)(1 - \kappa_1^3 t_1)(1 - \kappa_2^3 t_1)}},$$

where

$$w_1 = \frac{(\beta + \gamma)t_1 + \alpha}{\beta t_1 + (\alpha + \gamma)}, \quad w_2 = \frac{t_1[(\beta + \gamma)t_1 + \alpha]^2[\beta t_1 + (\alpha + \gamma)]}{t_2^5},$$

which is a rational transformation of order five. Here,

$$\left\{\begin{aligned} \alpha &= \frac{(v_1 - 1)(v_2 - 1)(v_1 + v_2 - 2) - (u_1 - 1)(u_2 - 1)(u_1 + u_2 - 2)}{2(u_1 - 1)(u_2 - 1)(v_1 - 1)(v_2 - 1)}, \\ \beta &= \frac{-(v_1 - 1)(v_2 - 1)(2v_1v_2 - v_1 - v_2) + (u_1 - 1)(u_2 - 1)(2u_1u_2 - u_1 - u_2)}{2(u_1 - 1)(u_2 - 1)(v_1 - 1)(v_2 - 1)}, \\ \gamma &= \frac{(v_1 - 1)(v_2 - 1)}{(u_1 - 1)(u_2 - 1)}, \end{aligned}\right.$$

with

$$u_1 = \kappa_1^3, \quad u_2 = \kappa_2^3, \quad v_1 = \lambda_1^3, \quad v_2 = \lambda_2^3.$$

Moreover, the moduli (κ_1, κ_2) *and* (λ_1, λ_2) *satisfy the modular equation*:

$$(\kappa_1^3 - 1)(\kappa_2^3 - 1)(\kappa_1^3 - \kappa_2^3) = (\lambda_1^3 - 1)(\lambda_2^3 - 1)(\lambda_1^3 - \lambda_2^3),$$

which is an algebraic variety of dimension three. The corresponding two algebraic surfaces are:

$$\begin{cases} w_3^3 = w_1^2 w_2^2 (1-w_1)(1-\lambda_1^3 w_1)(1-\lambda_2^3 w_1), \\ t_3^3 = t_1^2 t_2^2 (1-t_1)(1-\kappa_1^3 t_1)(1-\kappa_2^3 t_1), \end{cases}$$

which give the homogeneous algebraic equations of degree seven:

$$\begin{cases} w_3^3 w_4^4 = w_1^2 w_2^2 (w_4-w_1)(w_4-\lambda_1^3 w_1)(w_4-\lambda_2^3 w_1), \\ t_3^3 t_4^4 = t_1^2 t_2^2 (t_4-t_1)(t_4-\kappa_1^3 t_1)(t_4-\kappa_2^3 t_1). \end{cases}$$

In fact, the above modular equation can be written as follows:

$$C_9(\kappa_1, \kappa_2, 1) = C_9(\lambda_1, \lambda_2, 1). \tag{1.16}$$

This shows that our modular equation is intimately connected with the Hessian polyhedra.

Proposition 1.2 (Main Proposition 1). *The following identities hold*:

$$\begin{vmatrix} \frac{\partial C_6}{\partial \xi} & \frac{\partial C_6}{\partial \eta} & 2C_6 \\ \frac{\partial C_9}{\partial \xi} & \frac{\partial C_9}{\partial \eta} & 3C_9 \\ \frac{\partial C_{12}}{\partial \xi} & \frac{\partial C_{12}}{\partial \eta} & 4C_{12} \end{vmatrix} = -864 \mathfrak{C}_{12}^2. \tag{1.17}$$

$$\begin{vmatrix} \frac{\partial C_6}{\partial \xi} & \frac{\partial C_6}{\partial \eta} & 2C_6 \\ \frac{\partial C_9}{\partial \xi} & \frac{\partial C_9}{\partial \eta} & 3C_9 \\ \frac{\partial C_{18}}{\partial \xi} & \frac{\partial C_{18}}{\partial \eta} & 6C_{18} \end{vmatrix} = -1296 C_6 \mathfrak{C}_{12}^2. \tag{1.18}$$

$$\begin{vmatrix} \frac{\partial C_6}{\partial \xi} & \frac{\partial C_6}{\partial \eta} & 2C_6 \\ \frac{\partial C_{12}}{\partial \xi} & \frac{\partial C_{12}}{\partial \eta} & 4C_{12} \\ \frac{\partial C_{18}}{\partial \xi} & \frac{\partial C_{18}}{\partial \eta} & 6C_{18} \end{vmatrix} = 432 \cdot 864 C_9 \mathfrak{C}_{12}^2. \tag{1.19}$$

$$\begin{vmatrix} \frac{\partial C_9}{\partial \xi} & \frac{\partial C_9}{\partial \eta} & 3C_9 \\ \frac{\partial C_{12}}{\partial \xi} & \frac{\partial C_{12}}{\partial \eta} & 4C_{12} \\ \frac{\partial C_{18}}{\partial \xi} & \frac{\partial C_{18}}{\partial \eta} & 6C_{18} \end{vmatrix} = 1296 (C_6^2 - C_{12}) \mathfrak{C}_{12}^2. \tag{1.20}$$

$$\begin{vmatrix} \frac{\partial C_6}{\partial \xi} & \frac{\partial C_6}{\partial \eta} & 2C_6 \\ \frac{\partial C_9}{\partial \xi} & \frac{\partial C_9}{\partial \eta} & 3C_9 \\ \frac{\partial \mathfrak{C}_{12}}{\partial \xi} & \frac{\partial \mathfrak{C}_{12}}{\partial \eta} & 4\mathfrak{C}_{12} \end{vmatrix} = \frac{1}{2} (C_{12}^2 - C_6 C_{18}). \tag{1.21}$$

$$\begin{vmatrix} \frac{\partial C_6}{\partial \xi} & \frac{\partial C_6}{\partial \eta} & 2C_6 \\ \frac{\partial C_{12}}{\partial \xi} & \frac{\partial C_{12}}{\partial \eta} & 4C_{12} \\ \frac{\partial \mathfrak{C}_{12}}{\partial \xi} & \frac{\partial \mathfrak{C}_{12}}{\partial \eta} & 4\mathfrak{C}_{12} \end{vmatrix} = 144 C_9 C_{18}. \tag{1.22}$$

$$\begin{vmatrix} \frac{\partial C_9}{\partial \xi} & \frac{\partial C_9}{\partial \eta} & 3C_9 \\ \frac{\partial C_{12}}{\partial \xi} & \frac{\partial C_{12}}{\partial \eta} & 4C_{12} \\ \frac{\partial \mathfrak{C}_{12}}{\partial \xi} & \frac{\partial \mathfrak{C}_{12}}{\partial \eta} & 4\mathfrak{C}_{12} \end{vmatrix} = \frac{1}{2}(C_6^2 - C_{12}) C_{18}. \tag{1.23}$$

$$\begin{vmatrix} \frac{\partial C_6}{\partial \xi} & \frac{\partial C_6}{\partial \eta} & 2C_6 \\ \frac{\partial C_{18}}{\partial \xi} & \frac{\partial C_{18}}{\partial \eta} & 6C_{18} \\ \frac{\partial \mathfrak{C}_{12}}{\partial \xi} & \frac{\partial \mathfrak{C}_{12}}{\partial \eta} & 4\mathfrak{C}_{12} \end{vmatrix} = 216 C_9 C_{12}^2. \tag{1.24}$$

$$\begin{vmatrix} \frac{\partial C_9}{\partial \xi} & \frac{\partial C_9}{\partial \eta} & 3C_9 \\ \frac{\partial C_{18}}{\partial \xi} & \frac{\partial C_{18}}{\partial \eta} & 6C_{18} \\ \frac{\partial \mathfrak{C}_{12}}{\partial \xi} & \frac{\partial \mathfrak{C}_{12}}{\partial \eta} & 4\mathfrak{C}_{12} \end{vmatrix} = \frac{3}{4}(C_6^2 - C_{12}) C_{12}^2. \tag{1.25}$$

$$\begin{vmatrix} \frac{\partial C_{12}}{\partial \xi} & \frac{\partial C_{12}}{\partial \eta} & 4C_{12} \\ \frac{\partial C_{18}}{\partial \xi} & \frac{\partial C_{18}}{\partial \eta} & 6C_{18} \\ \frac{\partial \mathfrak{C}_{12}}{\partial \xi} & \frac{\partial \mathfrak{C}_{12}}{\partial \eta} & 4\mathfrak{C}_{12} \end{vmatrix} = 0. \tag{1.26}$$

We define the Hessian polyhedral equations as follows:

$$\begin{cases} 432 \dfrac{C_9^2}{C_6^3} = R_1(x, y), \\ 3 \dfrac{C_{12}}{C_6^2} = R_2(x, y), \end{cases} \tag{1.27}$$

where R_1 and R_2 are rational functions of x and y.

Theorem 1.3 (Main Theorem 2). *The relations between Hessian polyhedra and Appell hypergeometric partial differential equations are given by the following identities*:

$$\begin{aligned} & z_1^6 + z_2^6 + z_3^6 - 10(z_1^3 z_2^3 + z_2^3 z_3^3 + z_3^3 z_1^3) \\ =& 2^6 \cdot 3^7 \cdot C^2 \cdot R_1 \left[\frac{1}{4}(R_1 + R_2 - 1)^2 - \frac{1}{27} R_2^3\right]^{\frac{4}{3}} \left[\frac{\partial(R_1, R_2)}{\partial(x, y)}\right]^{-2} H(x, y)^2. \end{aligned} \tag{1.28}$$

$$\begin{aligned}&(z_1^3-z_2^3)(z_2^3-z_3^3)(z_3^3-z_1^3)\\ &=2^7\cdot 3^9\cdot C^3\cdot R_1^2\left[\frac{1}{4}(R_1+R_2-1)^2-\frac{1}{27}R_2^3\right]^2\left[\frac{\partial(R_1,R_2)}{\partial(x,y)}\right]^{-3}H(x,y)^3.\end{aligned}\tag{1.29}$$

$$\begin{aligned}&(z_1^3+z_2^3+z_3^3)[(z_1^3+z_2^3+z_3^3)^3+216z_1^3z_2^3z_3^3]\\ &=2^{12}\cdot 3^{13}\cdot C^4\cdot R_1^2R_2\left[\frac{1}{4}(R_1+R_2-1)^2-\frac{1}{27}R_2^3\right]^{\frac{8}{3}}\left[\frac{\partial(R_1,R_2)}{\partial(x,y)}\right]^{-4}H(x,y)^4.\end{aligned}\tag{1.30}$$

$$\begin{aligned}&z_1z_2z_3[27z_1^3z_2^3z_3^3-(z_1^3+z_2^3+z_3^3)^3]\\ &=2^{10}\cdot 3^{13}\cdot C^4\cdot R_1^2\left[\frac{1}{4}(R_1+R_2-1)^2-\frac{1}{27}R_2^3\right]^3\left[\frac{\partial(R_1,R_2)}{\partial(x,y)}\right]^{-4}H(x,y)^4.\end{aligned}\tag{1.31}$$

$$\begin{aligned}&(z_1^3+z_2^3+z_3^3)^6-540z_1^3z_2^3z_3^3(z_1^3+z_2^3+z_3^3)^3-5832z_1^6z_2^6z_3^6\\ &=2^{17}\cdot 3^{21}\cdot C^6\cdot R_1^3(R_1+R_2-1)\left[\frac{1}{4}(R_1+R_2-1)^2-\frac{1}{27}R_2^3\right]^4\\ &\times\left[\frac{\partial(R_1,R_2)}{\partial(x,y)}\right]^{-6}H(x,y)^6.\end{aligned}\tag{1.32}$$

Here,

$$H(x,y):=x^{b'-c}y^{b-c}(x-1)^{c-a-b-1}(y-1)^{c-a-b'-1}(x-y)^{-b-b'}.\tag{1.33}$$

The above five identities give the relations between the Hessian polyhedra and Appell hypergeometric partial differential equations. More precisely, given the Hessian polyhedra, we can obtain the rational functions R_1, R_2 and the function $H(x,y)$ with parameters a, b, b', c. Then three linearly independent solutions

$$z_1=z_1(x,y),\quad z_2=z_2(x,y),\quad z_3=z_3(x,y),$$

which are the algebraic solutions of the Appell hypergeometric partial differential equations (1.1) with the same parameters a, b, b', c, satisfy the above five algebraic equations (1.27), (1.28), (1.29), (1.30), (1.31). Moreover, the monodromy groups of these Appell hypergeometric partial differential equations are just the Hessian groups.

The Hessian polyhedral irrationality (ξ,η) is defined by the equations:

$$\begin{cases}432\dfrac{C_9^2}{C_6^3}=J_1,\\ 3\dfrac{C_{12}}{C_6^2}=J_2,\end{cases}\tag{1.34}$$

where J_1 and J_2 are J-invariants (see [Y]):

$$\left\{\begin{aligned} J_1 = J_1(\lambda_1, \lambda_2) &= \frac{\lambda_2^2(\lambda_2-1)^2}{\lambda_1^2(\lambda_1-1)^2(\lambda_1-\lambda_2)^2}, \\ J_2 = J_2(\lambda_1, \lambda_2) &= \frac{\lambda_1^2(\lambda_1-1)^2}{\lambda_2^2(\lambda_2-1)^2(\lambda_2-\lambda_1)^2}. \end{aligned}\right. \tag{1.35}$$

The corresponding Picard curve is

$$y^3 = x(x-1)(x-\lambda_1)(x-\lambda_2). \tag{1.36}$$

Here, $\lambda_1 = \lambda_1(\xi, \eta)$ and $\lambda_2 = \lambda_2(\xi, \eta)$ are Picard modular functions.

Denote by

$$(F, G, H) := \frac{\partial(F, G, H)}{\partial(z_1, z_2, z_3)}. \tag{1.37}$$

Proposition 1.4 (Main Proposition 2). *In the projective coordinates, the following duality formulas hold*:

$$\left\{\begin{aligned} (C_6, C_9, C_{12}) &= -2^5 \cdot 3^4 \cdot \mathfrak{C}_{12}^2, \\ (C_6, C_9, C_{18}) &= -2^4 \cdot 3^5 \cdot C_6 \mathfrak{C}_{12}^2, \\ (C_6, C_{12}, C_{18}) &= 2^9 \cdot 3^7 \cdot C_9 \mathfrak{C}_{12}^2, \\ (C_9, C_{12}, C_{18}) &= 2^4 \cdot 3^5 \cdot (C_6^2 - C_{12}) \mathfrak{C}_{12}^2. \end{aligned}\right. \tag{1.38}$$

$$\left\{\begin{aligned} (C_6, C_9, \mathfrak{C}_{12}) &= \frac{3}{2}(C_{12}^2 - C_6 C_{18}), \\ (C_6, C_{12}, \mathfrak{C}_{12}) &= 432 C_9 C_{18}, \\ (C_9, C_{12}, \mathfrak{C}_{12}) &= \frac{3}{2} C_{18}(C_6^2 - C_{12}). \end{aligned}\right. \tag{1.39}$$

$$\left\{\begin{aligned} (C_6, C_{18}, \mathfrak{C}_{12}) &= 648 C_9 C_{12}^2, \\ (C_9, C_{18}, \mathfrak{C}_{12}) &= \frac{9}{4} C_{12}^2 (C_6^2 - C_{12}), \\ (C_{12}, C_{18}, \mathfrak{C}_{12}) &= 0. \end{aligned}\right. \tag{1.40}$$

Put

$$\begin{cases} Y_1 = (C_6, C_{12}, \mathfrak{C}_{12}), \\ Y_2 = (C_9, C_{12}, \mathfrak{C}_{12}), \\ Y_3 = (C_6, C_{12}, C_{18}), \\ Y_4 = (C_9, C_{12}, C_{18}). \end{cases} \tag{1.41}$$

$$\begin{cases} X_1 = (C_6, C_9, C_{12}), \\ X_2 = (C_6, C_9, \mathfrak{C}_{12}), \\ X_3 = (C_6, C_9, C_{18}), \\ X_4 = (C_6, C_{18}, \mathfrak{C}_{12}), \\ X_5 = (C_9, C_{18}, \mathfrak{C}_{12}). \end{cases} \tag{1.42}$$

THEOREM 1.5 (MAIN THEOREM 3). *The functions X_1, X_2, X_3, X_4, X_5 and Y_1, Y_2, Y_3, Y_4 satisfy the system of algebraic equations of the sixth degree*:

$$\begin{cases} 18X_1^2Y_4^3 + 24X_1X_3^2Y_4^2 + 8X_3^4Y_4 + 27X_1^5X_5 = 0, \\ \hfill Y_2Y_3 = Y_1Y_4, \\ \hfill X_4Y_4 = X_5Y_3, \\ \hfill X_3Y_1 - X_2Y_3 = X_1X_4. \end{cases} \tag{1.43}$$

Let

$$\mathcal{A} = \{(C_6, C_9, C_{12}, \mathfrak{C}_{12}, C_{18}) \in \mathbb{C}^5\} \tag{1.44}$$

be the configuration space of the Hessian invariants. Let

$$\mathcal{B} = \{(X_1, X_2, X_3, X_4, X_5, Y_1, Y_2, Y_3, Y_4) \in \mathbb{C}^9\} \tag{1.45}$$

be the dual configuration space of $\mathcal{A}$. Then

$$\dim\mathcal{A} = 3, \quad \dim\mathcal{B} = 5. \tag{1.46}$$

For the Hessian polyhedral equations

$$Z_1 = 432\frac{C_9^2}{C_6^3}, \quad Z_2 = 3\frac{C_{12}}{C_6^2}, \tag{1.47}$$

we find that

$$\begin{aligned}
&Z_1^2 : Z_2^3 : (Z_1+Z_2-1)^2 : [27(Z_1+Z_2-1)^2-4Z_2^3] : 1\\
=&432^2 C_9^4 : 27C_{12}^3 : 4C_{18}^2 : 432^2 \mathfrak{C}_{12}^3 : C_6^6\\
=&432^2(\xi^3-\eta^3)^4(\eta^3-1)^4(1-\xi^3)^4\\
&:27(\xi^3+\eta^3+1)^3[(\xi^3+\eta^3+1)^3+216\xi^3\eta^3]^3\\
&:4[(\xi^3+\eta^3+1)^6-540\xi^3\eta^3(\xi^3+\eta^3+1)^3-5832\xi^6\eta^6]^2\\
&:432^2\xi^3\eta^3[27\xi^3\eta^3-(\xi^3+\eta^3+1)^3]^3\\
&:[\xi^6+\eta^6+1-10(\xi^3\eta^3+\xi^3+\eta^3)]^6.
\end{aligned} \tag{1.48}$$

Put

$$\tau_1 = \xi^3+\eta^3, \quad \tau_2 = \xi\eta. \tag{1.49}$$

Then

$$\begin{cases}
C_6 = \tau_1^2 - 12\tau_2^3 - 10\tau_1 + 1,\\
C_{12} = (\tau_1+1)[(\tau_1+1)^3+216\tau_2^3],\\
\mathfrak{C}_{12} = \tau_2[27\tau_2^3-(\tau_1+1)^3],\\
C_{18} = (\tau_1+1)^6 - 540\tau_2^3(\tau_1+1)^3 - 5832\tau_2^6,\\
C_9^2 = (\tau_1^2-4\tau_2^3)(\tau_2^3-\tau_1+1)^2.
\end{cases} \tag{1.50}$$

Thus,

$$\begin{aligned}
&Z_1^2 : Z_2^3 : (Z_1+Z_2-1)^2 : [27(Z_1+Z_2-1)^2-4Z_2^3] : 1\\
=&432^2(\tau_1^2-4\tau_2^3)^2(\tau_2^3-\tau_1+1)^4\\
&:27(\tau_1+1)^3[(\tau_1+1)^3+216\tau_2^3]^3\\
&:4[(\tau_1+1)^6-540\tau_2^3(\tau_1+1)^3-5832\tau_2^6]^2\\
&:432^2\tau_2^3[27\tau_2^3-(\tau_1+1)^3]^3\\
&:(\tau_1^2-12\tau_2^3-10\tau_1+1)^6.
\end{aligned} \tag{1.51}$$

Put

$$\varphi = z_1z_2z_3, \quad \psi = z_1^3+z_2^3+z_3^3, \quad \chi = z_1^3z_2^3+z_2^3z_3^3+z_3^3z_1^3. \tag{1.52}$$

Let

$$G(z_1,z_2,z_3) := (z_1-z_2)(z_2-z_3)(z_3-z_1). \tag{1.53}$$

$$H(z_1, z_2, z_3) := \frac{1}{3}[(z_1 + z_2 + z_3)^3 + (z_1 + \omega z_2 + \omega^2 z_3)^3 + (z_1 + \omega^2 z_2 + \omega z_3)^3]. \tag{1.54}$$

$$K(z_1, z_2, z_3) := (z_1 + z_2 + z_3)(z_1 + \omega z_2 + \omega^2 z_3)(z_1 + \omega^2 z_2 + \omega z_3). \tag{1.55}$$

The Hessians of G, K and H are:

$$\text{Hessian}(G) = 0, \quad \text{Hessian}(K) = -54K, \quad \text{Hessian}(H) = -216K. \tag{1.56}$$

THEOREM 1.6 (MAIN THEOREM 4). *The invariants G, H and K satisfy the following algebraic equations, which are the form-theoretic resolvents* (*algebraic resolvents*) *of G, H, K*:

$$\begin{cases} 4G^3 + H^2 G - C_6 G - 4C_9 = 0, \\ H(H^3 + 8K^3) - 9C_{12} = 0, \\ K(K^3 - H^3) - 27\mathfrak{C}_{12} = 0. \end{cases} \tag{1.57}$$

Let

$$r_1 = \frac{G^2}{C_6}, \quad r_2 = \frac{H^2}{C_6}, \quad r_3 = \frac{K^2}{C_6}. \tag{1.58}$$

PROPOSITION 1.7. *The following identities hold*:

$$\begin{cases} Z_1 = 27 r_1 (4r_1 + r_2 - 1)^2, \\ Z_2 = \frac{1}{3} r_2^2 + \frac{8}{3} r_3 \cdot \sqrt{r_2 r_3}, \\ \sqrt[3]{\frac{1}{4}(Z_1 + Z_2 - 1)^2 - \frac{1}{27} Z_2^3} = \frac{4}{9}(r_3^2 - r_2 \cdot \sqrt{r_2 r_3}). \end{cases} \tag{1.59}$$

THEOREM 1.8 (MAIN THEOREM 5). *The function-theoretic resolvents* (*analytic resolvents*) *of Z_1 and Z_2 are given by*:

$$\begin{aligned} & Z_1^2 : Z_2^3 : (Z_1 + Z_2 - 1)^2 : [27(Z_1 + Z_2 - 1)^2 - 4Z_2^3] : 1 \\ = & 729^2 r_1^2 (4r_1 + r_2 - 1)^4 : 27(r_2^2 + 8r_3 \cdot \sqrt{r_2 r_3})^3 \\ & : [256(r_3^2 - r_2 \cdot \sqrt{r_2 r_3})^3 + 4(r_2^2 + 8r_3 \cdot \sqrt{r_2 r_3})^3] \\ & : 6912(r_3^2 - r_2 \cdot \sqrt{r_2 r_3})^3 : 729. \end{aligned} \tag{1.60}$$

For the Hessian polyhedral equations, the corresponding J-invariants (J_1, J_2) satisfy:

$$\begin{aligned} &J_1^2 : J_2^3 : (J_1 + J_2 - 1)^2 : [27(J_1 + J_2 - 1)^2 - 4J_2^3] : 1 \\ =&729^2 v_1^2 (4v_1 + v_2 - 1)^4 : 27(v_2^2 + 8v_3 \cdot \sqrt{v_2 v_3})^3 \\ &:[256(v_3^2 - v_2 \cdot \sqrt{v_2 v_3})^3 + 4(v_2^2 + 8v_3 \cdot \sqrt{v_2 v_3})^3] \\ &:6912(v_3^2 - v_2 \cdot \sqrt{v_2 v_3})^3 : 729, \end{aligned} \tag{1.61}$$

where

$$v_1 = v_1(\omega_1, \omega_2), \quad v_2 = v_2(\omega_1, \omega_2), \quad v_3 = v_3(\omega_1, \omega_2)$$

are three Picard modular functions with $(\omega_1, \omega_2) \in \mathfrak{S}_2 = \{(z_1, z_2) \in \mathbb{C}^2 : z_1 + \overline{z_1} - z_2 \overline{z_2} > 0\}$.

Put

$$u = \frac{C_6 G}{C_9}, \quad v = \frac{C_9 H}{C_{12}}, \quad w = \frac{C_9 K}{\mathfrak{C}_{12}}. \tag{1.62}$$

We will denote the next three equations as the resolvents of the u, v and w's.

Theorem 1.9 (Main Theorem 6). *The functions u, v and w satisfy the following algebraic equations, which are the analytic and algebraic resolvents of u, v and w:*

$$\begin{cases} Z_1^2 u^3 + 5184 Z_2^2 u v^2 - 108 Z_1 u - 432 Z_1 = 0, \\ 6912 Z_2^3 v^4 + 8[27(Z_1 + Z_2 - 1)^2 - 4Z_2^3] v w^3 - 9 Z_1^2 = 0, \\ [27(Z_1 + Z_2 - 1)^2 - 4Z_2^3] w^4 - 6912 Z_2^3 v^3 w - 27 Z_1^2 = 0. \end{cases} \tag{1.63}$$

Theorem 1.10 (Main Theorem 7). *The functions u, v and w can be expressed as the algebraic functions of r_1, r_2 and r_3:*

$$\begin{cases} u = \dfrac{4}{4r_1 + r_2 - 1}, \\ v = \dfrac{9}{4} \dfrac{(4r_1 + r_2 - 1)\sqrt{r_1}}{r_2\sqrt{r_2} + 8r_3\sqrt{r_3}}, \\ w = \dfrac{27}{4} \dfrac{(4r_1 + r_2 - 1)\sqrt{r_1}}{r_3\sqrt{r_3} - r_2\sqrt{r_2}}. \end{cases} \tag{1.64}$$

Set

$$M := M_\infty = 81\sqrt{-3}\varphi^3, \quad N := N_\infty = 3\sqrt{-3}\psi^3. \tag{1.65}$$

$$M_0 := M(E(z_1, z_2, z_3)) = K^3, \quad N_0 := N(E(z_1, z_2, z_3)) = H^3. \tag{1.66}$$

They satisfy:

$$N_0(N_0+8M_0)^3=9^3C_{12}^3, \quad M_0(M_0-N_0)^3=27^3\mathfrak{C}_{12}^3. \tag{1.67}$$

Let

$$M_i:=M(ED^i(z_1,z_2,z_3))=(\psi-3\omega^{2i}\varphi)^3, \quad N_i:=N(ED^i(z_1,z_2,z_3))=(\psi+6\omega^{2i}\varphi)^3. \tag{1.68}$$

Then

$$M_\infty M_0M_1M_2=-81\sqrt{-3}\mathfrak{C}_{12}^3, \quad N_\infty N_0N_1N_2=3\sqrt{-3}C_{12}^3. \tag{1.69}$$

Let

$$L(z_1,z_2,z_3):=(z_1-z_2)^3(z_2-z_3)^3(z_3-z_1)^3. \tag{1.70}$$

Set

$$\begin{cases} L_{1,0,0}:=L(E(z_1,z_2,z_3))=(z_2^3-z_3^3)^3, \\ L_{1,1,0}:=L(EA(z_1,z_2,z_3))=(z_3^3-z_1^3)^3, \\ L_{1,2,0}:=L(EA^2(z_1,z_2,z_3))=(z_1^3-z_2^3)^3, \end{cases} \tag{1.71}$$

and

$$\begin{cases} L_{0,0,0}:=L(z_1,z_2,z_3)=(z_1-z_2)^3(z_2-z_3)^3(z_3-z_1)^3, \\ L_{0,0,1}:=L(C(z_1,z_2,z_3))=(z_1-\omega z_2)^3(\omega z_2-\omega^2z_3)^3(\omega^2z_3-z_1)^3, \\ L_{0,0,2}:=L(C^2(z_1,z_2,z_3))=(z_1-\omega^2z_2)^3(\omega^2z_2-\omega z_3)^3(\omega z_3-z_1)^3. \end{cases} \tag{1.72}$$

Then

$$L_{1,0,0}L_{1,1,0}L_{1,2,0}=L_{0,0,0}L_{0,0,1}L_{0,0,2}=C_9^3. \tag{1.73}$$

Put

$$\zeta_1=\frac{M_0}{C_9}, \quad \zeta_2=\frac{N_0}{C_9}. \tag{1.74}$$

THEOREM 1.11 (MAIN THEOREM 8). *The functions ζ_1 and ζ_2 satisfy the following equations*:

$$\begin{cases} Z_1^2\zeta_2(\zeta_2+8\zeta_1)^3=2^8\cdot3^9Z_2^3, \\ Z_1^2\zeta_1(\zeta_1-\zeta_2)^3=27^3[27(Z_1+Z_2-1)^2-4Z_2^3]. \end{cases} \tag{1.75}$$

Put

$$\xi_1 := \frac{M_0^{\frac{1}{3}}}{C_6^{\frac{1}{2}}}, \quad \xi_2 := \frac{N_0^{\frac{1}{3}}}{C_6^{\frac{1}{2}}}. \tag{1.76}$$

Theorem 1.12 (Main Theorem 9). *The functions ξ_1 and ξ_2 satisfy the following equations*:

$$\begin{cases} \xi_2^3(\xi_2^3+8\xi_1^3)^3 = 27Z_2^3, \\ 256\xi_1^3(\xi_1^3-\xi_2^3)^3 = 27[27(Z_1+Z_2-1)^2-4Z_2^3]. \end{cases} \tag{1.77}$$

The functions ξ_1, ξ_2 and ζ_1, ζ_2 satisfy the following relations:

$$\begin{cases} \frac{4}{729}\zeta_1^4[\xi_2^3(\xi_2^3+8\xi_1^3)^3+64\xi_1^3(\xi_1^3-\xi_2^3)^3] = [432\xi_1^6+\frac{1}{3}\zeta_1^2\xi_2(\xi_2^3+8\xi_1^3)-\zeta_1^2]^2, \\ \frac{4}{729}\zeta_2^4[\xi_2^3(\xi_2^3+8\xi_1^3)^3+64\xi_1^3(\xi_1^3-\xi_2^3)^3] = [432\xi_2^6+\frac{1}{3}\zeta_2^2\xi_2(\xi_2^3+8\xi_1^3)-\zeta_2^2]^2. \end{cases} \tag{1.78}$$

Theorem 1.13 (Main Theorem 10). *The function-theoretic resolvents* (*analytic resolvents*) *of Z_1 and Z_2 are also given by*:

$$\begin{aligned} &Z_1^2 : Z_2^3 : (Z_1+Z_2-1)^2 : [27(Z_1+Z_2-1)^2-4Z_2^3] : 1 \\ =&[2(\xi_2^3(\xi_2^3+8\xi_1^3)^3+64\xi_1^3(\xi_1^3-\xi_2^3)^3)^{\frac{1}{2}}-9\xi_2(\xi_2^3+8\xi_1^3)+27]^2 \\ &:27\xi_2^3(\xi_2^3+8\xi_1^3)^3 : 4[\xi_2^3(\xi_2^3+8\xi_1^3)^3+64\xi_1^3(\xi_1^3-\xi_2^3)^3] \\ &:6912\xi_1^3(\xi_1^3-\xi_2^3)^3 : 729. \end{aligned} \tag{1.79}$$

In fact,

$$\begin{cases} \sqrt[3]{M_0}+\sqrt[3]{M_1}+\sqrt[3]{M_2} = 3\psi, \\ \sqrt{-3}\sqrt[3]{M_\infty}+\sqrt[3]{M_0}+\omega\sqrt[3]{M_1}+\omega^2\sqrt[3]{M_2} = 0, \\ \sqrt[3]{M_0}+\omega^2\sqrt[3]{M_1}+\omega\sqrt[3]{M_2} = 0. \end{cases} \tag{1.80}$$

$$\begin{cases} -\sqrt{-3}\sqrt[3]{N_\infty}+\sqrt[3]{N_0}+\sqrt[3]{N_1}+\sqrt[3]{N_2} = 0, \\ \sqrt[3]{N_0}+\omega\sqrt[3]{N_1}+\omega^2\sqrt[3]{N_2} = 18\varphi, \\ \sqrt[3]{N_0}+\omega^2\sqrt[3]{N_1}+\omega\sqrt[3]{N_2} = 0. \end{cases} \tag{1.81}$$

$$\left\{\begin{aligned}&\sqrt[3]{L_{1,0,0}}+\sqrt[3]{L_{1,1,0}}+\sqrt[3]{L_{1,2,0}}=0,\\&\sqrt[3]{L_{0,0,0}}+\sqrt[3]{L_{0,0,1}}+\sqrt[3]{L_{0,0,2}}=0.\end{aligned}\right.\tag{1.82}$$

$$\left\{\begin{aligned}&L_{1,0,0}+L_{1,1,0}+L_{1,2,0}=3C_9,\\&L_{0,0,0}+L_{0,0,1}+L_{0,0,2}=3C_9.\end{aligned}\right.\tag{1.83}$$

$$\sqrt{-3}(M_0+M_1+M_2)=N_\infty-M_\infty,\quad \sqrt{-3}(N_0+N_1+N_2)=N_\infty+8M_\infty.\tag{1.84}$$

$$\begin{aligned}(L_{1,0,0}+L_{1,1,0}+L_{1,2,0})^3&=27L_{1,0,0}L_{1,1,0}L_{1,2,0},\\(L_{0,0,0}+L_{0,0,1}+L_{0,0,2})^3&=27L_{0,0,0}L_{0,0,1}L_{0,0,2}.\end{aligned}\tag{1.85}$$

Put

$$X(z_1,z_2,z_3)=z_1^3,\quad Y(z_1,z_2,z_3)=z_2^3,\quad Z(z_1,z_2,z_3)=z_3^3.\tag{1.86}$$

$$Q_1(z_1,z_2,z_3)=z_1z_2^2+z_2z_3^2+z_3z_1^2,\quad Q_2(z_1,z_2,z_3)=z_1^2z_2+z_2^2z_3+z_3^2z_1.\tag{1.87}$$

The functions φ, ψ, χ, X, Y, Z and Q_1, Q_2 satisfy the relations:

$$\left\{\begin{aligned}\varphi^3&=XYZ,\\\psi&=X+Y+Z,\\\chi&=XY+YZ+ZX,\\\chi+3\varphi^2+\varphi\psi&=Q_1Q_2,\\\psi\chi+6\varphi\chi+6\varphi^2\psi+9\varphi^3&=Q_1^3+Q_2^3,\\(X-Y)(Y-Z)(Z-X)&=Q_1^3-Q_2^3.\end{aligned}\right.\tag{1.88}$$

Let

$$\left\{\begin{aligned}&W_2=(X+Y+Z)^2-12(XY+YZ+ZX),\\&W_3=(X-Y)(Y-Z)(Z-X),\\&\mathfrak{W}_3=XYZ-\varphi^3,\\&W_4=(X+Y+Z)[(X+Y+Z)^3+216\varphi^3],\\&\mathfrak{W}_4=\varphi[27\varphi^3-(X+Y+Z)^3],\\&W_6=(X+Y+Z)^6-540\varphi^3(X+Y+Z)^3-5832\varphi^6.\end{aligned}\right.\tag{1.89}$$

We will regard (φ, X, Y, Z) as a point in the complex projective space $\mathbb{CP}^3$. In fact,

$$1728\mathfrak{W}_4^3 = W_6^2 - W_4^3 \tag{1.90}$$

and

$$\frac{\partial(\mathfrak{W}_3, W_4, \mathfrak{W}_4, W_6)}{\partial(\varphi, X, Y, Z)} = 0. \tag{1.91}$$

For the ternary cubic form

$$F_1(x_1, x_2, x_3) = Xx_1^3 + Yx_2^3 + Zx_3^3 - 3\varphi x_1 x_2 x_3, \tag{1.92}$$

we have

$$\begin{cases} \sqrt[3]{M_{\infty}} = -3\sqrt{-3}\varphi, \\ \sqrt[3]{M_0} = -3\varphi + X + Y + Z, \\ \sqrt[3]{M_1} = -3\omega^2\varphi + X + Y + Z, \\ \sqrt[3]{M_2} = -3\omega\varphi + X + Y + Z. \end{cases} \tag{1.93}$$

Correspondingly,

$$\begin{cases} \sqrt[3]{M_{\infty}}: \quad (\sqrt{-3}, 0, 0, 0) \in \mathbb{CP}^3, \\ \sqrt[3]{M_0}: \quad (1, 1, 1, 1) \in \mathbb{CP}^3, \\ \sqrt[3]{M_1}: \quad (\omega^2, 1, 1, 1) \in \mathbb{CP}^3, \\ \sqrt[3]{M_2}: \quad (\omega, 1, 1, 1) \in \mathbb{CP}^3. \end{cases} \tag{1.94}$$

For the ternary cubic form

$$F_2(x_1, x_2, x_3) = Xx_1^3 + Yx_2^3 + Zx_3^3 + 6\varphi x_1 x_2 x_3, \tag{1.95}$$

we have

$$\begin{cases} \sqrt[3]{N_{\infty}} = -\sqrt{-3}(X + Y + Z), \\ \sqrt[3]{N_0} = 6\varphi + X + Y + Z, \\ \sqrt[3]{N_1} = 6\omega^2\varphi + X + Y + Z, \\ \sqrt[3]{N_2} = 6\omega\varphi + X + Y + Z. \end{cases} \tag{1.96}$$

Correspondingly,

$$(1.97)\qquad \begin{cases} \sqrt[3]{N_\infty}: & (0,-\sqrt{-3},-\sqrt{-3},-\sqrt{-3})\in\mathbb{CP}^3,\\ \sqrt[3]{N_0}: & (1,1,1,1)\in\mathbb{CP}^3,\\ \sqrt[3]{N_1}: & (\omega^2,1,1,1)\in\mathbb{CP}^3,\\ \sqrt[3]{N_2}: & (\omega,1,1,1)\in\mathbb{CP}^3. \end{cases}$$

When $\mathfrak{W}_3=0$, we set

$$(1.98)\qquad \begin{cases} W_2=(X+Y+Z)^2-12(XY+YZ+ZX),\\ W_3=(X-Y)(Y-Z)(Z-X),\\ W_4=(X+Y+Z)[(X+Y+Z)^3+216XYZ],\\ W_6=(X+Y+Z)^6-540XYZ(X+Y+Z)^3-5832X^2Y^2Z^2,\\ W_{12}=XYZ[27XYZ-(X+Y+Z)^3]^3. \end{cases}$$

The invariants W_2, W_3, W_4, W_6 and W_{12} satisfy the relations:

$$(1.99)\qquad \begin{cases} 432W_3^2=W_2^3-3W_2W_4+2W_6,\\ 1728W_{12}=W_6^2-W_4^3. \end{cases}$$

The algebraic curves in $\mathbb{CP}^2$: $W_2=0$, $W_4=0$, $W_6=0$, $W_{12}=0$ intersect $W_3=0$ in 6, 12, 18, 15 points respectively.

$$\{W_2=0\}\cap\{W_3=0\}=\{(\alpha_i,1,1),(1,\alpha_i,1),(1,1,\alpha_i),i=1,2\},$$

where α_1 and α_2 are two roots of the quadratic equation:

$$\alpha^2-20\alpha-8=0.$$

$$\begin{aligned}&\{W_4=0\}\cap\{W_3=0\}\\ =&\{(-2,1,1),(1,-2,1),(1,1,-2),(\beta_i,1,1),(1,\beta_i,1),(1,1,\beta_i),i=1,2,3\},\end{aligned}$$

where β_1, β_2 and β_3 are the roots of the cubic equation:

$$\beta^3+6\beta^2+228\beta+8=0.$$

$$\{W_6=0\}\cap\{W_3=0\}=\{(\gamma_i,1,1),(1,\gamma_i,1),(1,1,\gamma_i),1\le i\le 6\},$$

where γ_i $(1\le i\le 6)$ are the roots of the equation of the sixth degree:

$$[\gamma^3+6\gamma^2-(258+162\sqrt{3})\gamma+8][\gamma^3+6\gamma^2-(258-162\sqrt{3})\gamma+8]=0.$$

$$\{W_{12}=0\}\cap\{W_3=0\}=\{(1,0,0),(0,1,0),(0,0,1),(1,1,0),(1,0,1),(0,1,1),\\ (\delta_i,1,1),(1,\delta_i,1),(1,1,\delta_i), i=1,2,3\},$$

where δ_1, δ_2 and δ_3 are the roots of the cubic equation:

$$\delta^3+6\delta^2-15\delta+8=0.$$

In fact,

$$\begin{aligned}&\mathbb{C}[W_2,W_3,W_4,W_6,W_{12}]/(W_2^3-3W_2W_4-432W_3^2+2W_6, W_6^2-W_4^3-1728W_{12})\\ \cong&\mathbb{C}[W_2,W_3,W_4].\end{aligned}\tag{1.100}$$

We find that

$$\begin{cases}(\sqrt{-3})^3\varphi(E(z_1,z_2,z_3))=\psi-3\varphi,\\ -\sqrt{-3}\psi(E(z_1,z_2,z_3))=\psi+6\varphi,\\ (\sqrt{-3})^3X(E(z_1,z_2,z_3))=\psi+6\varphi+3Q_1+3Q_2,\\ (\sqrt{-3})^3Y(E(z_1,z_2,z_3))=\psi+6\varphi+3\omega^2Q_1+3\omega Q_2,\\ (\sqrt{-3})^3Z(E(z_1,z_2,z_3))=\psi+6\varphi+3\omega Q_1+3\omega^2Q_2,\\ -\sqrt{-3}Q_1(E(z_1,z_2,z_3))=X+\omega^2Y+\omega Z,\\ -\sqrt{-3}Q_2(E(z_1,z_2,z_3))=X+\omega Y+\omega^2Z.\end{cases}\tag{1.101}$$

$$H_1:=\text{Hessian}(Q_1)=\begin{vmatrix}\frac{\partial^2Q_1}{\partial z_1^2}&\frac{\partial^2Q_1}{\partial z_1\partial z_2}&\frac{\partial^2Q_1}{\partial z_1\partial z_3}\\ \frac{\partial^2Q_1}{\partial z_1\partial z_2}&\frac{\partial^2Q_1}{\partial z_2^2}&\frac{\partial^2Q_1}{\partial z_2\partial z_3}\\ \frac{\partial^2Q_1}{\partial z_1\partial z_3}&\frac{\partial^2Q_1}{\partial z_2\partial z_3}&\frac{\partial^2Q_1}{\partial z_3^2}\end{vmatrix}=-8(\psi-3\varphi).\tag{1.102}$$

$$H_2:=\text{Hessian}(Q_2)=\begin{vmatrix}\frac{\partial^2Q_2}{\partial z_1^2}&\frac{\partial^2Q_2}{\partial z_1\partial z_2}&\frac{\partial^2Q_2}{\partial z_1\partial z_3}\\ \frac{\partial^2Q_2}{\partial z_1\partial z_2}&\frac{\partial^2Q_2}{\partial z_2^2}&\frac{\partial^2Q_2}{\partial z_2\partial z_3}\\ \frac{\partial^2Q_2}{\partial z_1\partial z_3}&\frac{\partial^2Q_2}{\partial z_2\partial z_3}&\frac{\partial^2Q_2}{\partial z_3^2}\end{vmatrix}=-8(\psi-3\varphi).\tag{1.103}$$

$$H_3:=\text{Hessian}(\varphi)=2\varphi.\tag{1.104}$$

$$H_4:=\text{Hessian}(\psi)=216\varphi.\tag{1.105}$$

We have

$$J_1 := \begin{vmatrix} \frac{\partial^2 Q_1}{\partial z_1^2} & \frac{\partial^2 Q_1}{\partial z_1 \partial z_2} & \frac{\partial^2 Q_1}{\partial z_1 \partial z_3} & \frac{\partial H_1}{\partial z_1} \\ \frac{\partial^2 Q_1}{\partial z_1 \partial z_2} & \frac{\partial^2 Q_1}{\partial z_2^2} & \frac{\partial^2 Q_1}{\partial z_2 \partial z_3} & \frac{\partial H_1}{\partial z_2} \\ \frac{\partial^2 Q_1}{\partial z_1 \partial z_3} & \frac{\partial^2 Q_1}{\partial z_2 \partial z_3} & \frac{\partial^2 Q_1}{\partial z_3^2} & \frac{\partial H_1}{\partial z_3} \\ \frac{\partial H_1}{\partial z_1} & \frac{\partial H_1}{\partial z_2} & \frac{\partial H_1}{\partial z_3} & 0 \end{vmatrix} = 6912[3Q_1^2 - (\psi + 6\varphi)Q_2]. \tag{1.106}$$

$$J_2 := \begin{vmatrix} \frac{\partial^2 Q_2}{\partial z_1^2} & \frac{\partial^2 Q_2}{\partial z_1 \partial z_2} & \frac{\partial^2 Q_2}{\partial z_1 \partial z_3} & \frac{\partial H_2}{\partial z_1} \\ \frac{\partial^2 Q_2}{\partial z_1 \partial z_2} & \frac{\partial^2 Q_2}{\partial z_2^2} & \frac{\partial^2 Q_2}{\partial z_2 \partial z_3} & \frac{\partial H_2}{\partial z_2} \\ \frac{\partial^2 Q_2}{\partial z_1 \partial z_3} & \frac{\partial^2 Q_2}{\partial z_2 \partial z_3} & \frac{\partial^2 Q_2}{\partial z_3^2} & \frac{\partial H_2}{\partial z_3} \\ \frac{\partial H_2}{\partial z_1} & \frac{\partial H_2}{\partial z_2} & \frac{\partial H_2}{\partial z_3} & 0 \end{vmatrix} = 6912[3Q_2^2 - (\psi + 6\varphi)Q_1]. \tag{1.107}$$

$$J_3 := \begin{vmatrix} \frac{\partial^2 \varphi}{\partial z_1^2} & \frac{\partial^2 \varphi}{\partial z_1 \partial z_2} & \frac{\partial^2 \varphi}{\partial z_1 \partial z_3} & \frac{\partial H_3}{\partial z_1} \\ \frac{\partial^2 \varphi}{\partial z_1 \partial z_2} & \frac{\partial^2 \varphi}{\partial z_2^2} & \frac{\partial^2 \varphi}{\partial z_2 \partial z_3} & \frac{\partial H_3}{\partial z_2} \\ \frac{\partial^2 \varphi}{\partial z_1 \partial z_3} & \frac{\partial^2 \varphi}{\partial z_2 \partial z_3} & \frac{\partial^2 \varphi}{\partial z_3^2} & \frac{\partial H_3}{\partial z_3} \\ \frac{\partial H_3}{\partial z_1} & \frac{\partial H_3}{\partial z_2} & \frac{\partial H_3}{\partial z_3} & 0 \end{vmatrix} = -12\varphi^2. \tag{1.108}$$

$$J_4 := \begin{vmatrix} \frac{\partial^2 \psi}{\partial z_1^2} & \frac{\partial^2 \psi}{\partial z_1 \partial z_2} & \frac{\partial^2 \psi}{\partial z_1 \partial z_3} & \frac{\partial H_4}{\partial z_1} \\ \frac{\partial^2 \psi}{\partial z_1 \partial z_2} & \frac{\partial^2 \psi}{\partial z_2^2} & \frac{\partial^2 \psi}{\partial z_2 \partial z_3} & \frac{\partial H_4}{\partial z_2} \\ \frac{\partial^2 \psi}{\partial z_1 \partial z_3} & \frac{\partial^2 \psi}{\partial z_2 \partial z_3} & \frac{\partial^2 \psi}{\partial z_3^2} & \frac{\partial H_4}{\partial z_3} \\ \frac{\partial H_4}{\partial z_1} & \frac{\partial H_4}{\partial z_2} & \frac{\partial H_4}{\partial z_3} & 0 \end{vmatrix} = -36^4\chi. \tag{1.109}$$

$$K_1 := \begin{vmatrix} \frac{\partial Q_1}{\partial z_1} & \frac{\partial Q_1}{\partial z_2} & \frac{\partial Q_1}{\partial z_3} \\ \frac{\partial H_1}{\partial z_1} & \frac{\partial H_1}{\partial z_2} & \frac{\partial H_1}{\partial z_3} \\ \frac{\partial J_1}{\partial z_1} & \frac{\partial J_1}{\partial z_2} & \frac{\partial J_1}{\partial z_3} \end{vmatrix} = 6912 \cdot 24[(\psi + 6\varphi)^3 - 27Q_2^3]. \tag{1.110}$$

$$K_2 := \begin{vmatrix} \frac{\partial Q_2}{\partial z_1} & \frac{\partial Q_2}{\partial z_2} & \frac{\partial Q_2}{\partial z_3} \\ \frac{\partial H_2}{\partial z_1} & \frac{\partial H_2}{\partial z_2} & \frac{\partial H_2}{\partial z_3} \\ \frac{\partial J_2}{\partial z_1} & \frac{\partial J_2}{\partial z_2} & \frac{\partial J_2}{\partial z_3} \end{vmatrix} = -6912 \cdot 24[(\psi + 6\varphi)^3 - 27Q_1^3]. \tag{1.111}$$

$$K_3 := \begin{vmatrix} \frac{\partial \varphi}{\partial z_1} & \frac{\partial \varphi}{\partial z_2} & \frac{\partial \varphi}{\partial z_3} \\ \frac{\partial H_3}{\partial z_1} & \frac{\partial H_3}{\partial z_2} & \frac{\partial H_3}{\partial z_3} \\ \frac{\partial J_3}{\partial z_1} & \frac{\partial J_3}{\partial z_2} & \frac{\partial J_3}{\partial z_3} \end{vmatrix} = 0. \tag{1.112}$$

$$K_4 := \begin{vmatrix} \frac{\partial \psi}{\partial z_1} & \frac{\partial \psi}{\partial z_2} & \frac{\partial \psi}{\partial z_3} \\ \frac{\partial H_4}{\partial z_1} & \frac{\partial H_4}{\partial z_2} & \frac{\partial H_4}{\partial z_3} \\ \frac{\partial J_4}{\partial z_1} & \frac{\partial J_4}{\partial z_2} & \frac{\partial J_4}{\partial z_3} \end{vmatrix} = -36^5 \cdot 54(Q_1^3 - Q_2^3). \tag{1.113}$$

$$(G, H, K) := \begin{vmatrix} \frac{\partial G}{\partial z_1} & \frac{\partial G}{\partial z_2} & \frac{\partial G}{\partial z_3} \\ \frac{\partial H}{\partial z_1} & \frac{\partial H}{\partial z_2} & \frac{\partial H}{\partial z_3} \\ \frac{\partial K}{\partial z_1} & \frac{\partial K}{\partial z_2} & \frac{\partial K}{\partial z_3} \end{vmatrix} = 27[3(Q_1^2 + Q_2^2) - (\psi + 6\varphi)(Q_1 + Q_2)]. \tag{1.114}$$

Put

$$U_1 = z_1 z_2^2, \quad U_2 = z_2 z_3^2, \quad U_3 = z_3 z_1^2, \tag{1.115}$$

$$V_1 = z_1^2 z_2, \quad V_2 = z_2^2 z_3, \quad V_3 = z_3^2 z_1. \tag{1.116}$$

Then

$$U_1 V_1 = XY, \quad U_2 V_2 = YZ, \quad U_3 V_3 = ZX. \tag{1.117}$$

$$U_1 U_2 U_3 = V_1 V_2 V_3 = XYZ = \varphi^3. \tag{1.118}$$

$$Q_1 = U_1 + U_2 + U_3, \quad Q_2 = V_1 + V_2 + V_3, \quad \psi = X + Y + Z. \tag{1.119}$$

The ternary cubic form

$$\begin{aligned} F(x_1, x_2, x_3) =& X x_1^3 + Y x_2^3 + Z x_3^3 + 6\varphi x_1 x_2 x_3 + \\ &+ 3(U_1 x_1 x_2^2 + U_2 x_2 x_3^2 + U_3 x_3 x_1^2) + 3(V_1 x_1^2 x_2 + V_2 x_2^2 x_3 + V_3 x_3^2 x_1) \\ =& (z_1 x_1 + z_2 x_2 + z_3 x_3)^3. \end{aligned} \tag{1.120}$$

$$\begin{cases} (\sqrt{-3})^3 U_1(E(z_1, z_2, z_3)) = X + \omega^2 Y + \omega Z + (\omega - 1) U_1 + (1 - \omega^2) U_2 + \\ \qquad + (\omega^2 - \omega) U_3 + (\omega - \omega^2) V_1 + (1 - \omega) V_2 + (\omega^2 - 1) V_3, \\ (\sqrt{-3})^3 U_2(E(z_1, z_2, z_3)) = X + \omega^2 Y + \omega Z + (1 - \omega^2) U_1 + (\omega^2 - \omega) U_2 + \\ \qquad + (\omega - 1) U_3 + (\omega^2 - 1) V_1 + (\omega - \omega^2) V_2 + (1 - \omega) V_3, \\ (\sqrt{-3})^3 U_3(E(z_1, z_2, z_3)) = X + \omega^2 Y + \omega Z + (\omega^2 - \omega) U_1 + (\omega - 1) U_2 + \\ \qquad + (1 - \omega^2) U_3 + (1 - \omega) V_1 + (\omega^2 - 1) V_2 + (\omega - \omega^2) V_3. \end{cases} \tag{1.121}$$

(1.122)
$$\begin{cases}(\sqrt{-3})^3 V_1(E(z_1, z_2, z_3)) = X+\omega Y+\omega^2 Z+(\omega-\omega^2)U_1+(\omega^2-1)U_2+\\ \qquad +(1-\omega)U_3+(1-\omega^2)V_1+(\omega-1)V_2+(\omega^2-\omega)V_3,\\ (\sqrt{-3})^3 V_2(E(z_1, z_2, z_3)) = X+\omega Y+\omega^2 Z+(1-\omega)U_1+(\omega-\omega^2)U_2+\\ \qquad +(\omega^2-1)U_3+(\omega-1)V_1+(\omega^2-\omega)V_2+(1-\omega^2)V_3,\\ (\sqrt{-3})^3 V_3(E(z_1, z_2, z_3)) = X+\omega Y+\omega^2 Z+(\omega^2-1)U_1+(1-\omega)U_2+\\ \qquad +(\omega-\omega^2)U_3+(\omega^2-\omega)V_1+(1-\omega^2)V_2+(\omega-1)V_3.\end{cases}$$

Let us define three rational functions (invariants) associated to the complex projective plane $\mathbb{CP}^2$:

$$\begin{cases}f_1^{\text{Hessian}}(\xi, \eta) := 432\dfrac{C_9(\xi, \eta)^2}{C_6(\xi, \eta)^3},\\ f_2^{\text{Hessian}}(\xi, \eta) := 3\dfrac{C_{12}(\xi, \eta)}{C_6(\xi, \eta)^2},\\ f_3^{\text{Hessian}}(\xi, \eta) := \dfrac{C_{18}(\xi, \eta)^2}{1728\mathfrak{C}_{12}(\xi, \eta)^3}.\end{cases} \tag{1.123}$$

Put

(1.124)
$$\begin{cases}F_1^{\text{Hessian}}(\xi, \eta) := (f_1^{\text{Hessian}}(\xi, \eta), f_3^{\text{Hessian}}(\xi, \eta)) = \left(\dfrac{432C_9(\xi, \eta)^2}{C_6(\xi, \eta)^3}, \dfrac{C_{18}(\xi, \eta)^2}{1728\mathfrak{C}_{12}(\xi, \eta)^3}\right),\\ F_2^{\text{Hessian}}(\xi, \eta) := (f_2^{\text{Hessian}}(\xi, \eta), f_3^{\text{Hessian}}(\xi, \eta)) = \left(\dfrac{3C_{12}(\xi, \eta)}{C_6(\xi, \eta)^2}, \dfrac{C_{18}(\xi, \eta)^2}{1728\mathfrak{C}_{12}(\xi, \eta)^3}\right).\end{cases}$$

The function $F_1^{\text{Hessian}}(\xi, \eta)$ is ramified at the points $(0, 0)$, $(\infty, 0)$, $(0, 1)$, $(\infty, 1)$, $(0, \infty)$, (∞, ∞), with degrees 162, 108, 108, 72, 108, 72, respectively. Their least common multiple is 648. The function $F_2^{\text{Hessian}}(\xi, \eta)$ is ramified at the points $(0, 0)$, $(\infty, 0)$, $(0, 1)$, $(\infty, 1)$, $(0, \infty)$, (∞, ∞), with degrees 216, 108, ∞, 72, 144, 72, respectively. Their least common multiple is 1296. Thus, the rational invariants F_1^{Hessian} and F_2^{Hessian} correspond to the Hessian groups G_{648} and G_{1296}, respectively.

This paper consists of seven sections. In section two, we give the reciprocity law about the Appell hypergeometric partial differential equations. In section three, we study the algebraic solutions of Appell hypergeometric partial differential equations. In section four, we give the Hessian polyhedral equations and study their properties. In section five, we study the invariant theory for the system of algebraic equations which can be solved by our Hessian polyhedral equations. In section six, we study the ternary cubic forms associated to the Hessian polyhedra. In section seven, we find some rational invariants on $\mathbb{CP}^2$.

2. The reciprocity law about the Appell hypergeometric functions

Let us consider the Appell hypergeometric partial differential equations (see [AK] for more details):

$$\begin{cases} x(1-x)\dfrac{\partial^2 z}{\partial x^2}+y(1-x)\dfrac{\partial^2 z}{\partial x \partial y}+[c-(a+b+1)x]\dfrac{\partial z}{\partial x}-by\dfrac{\partial z}{\partial y}-abz=0, \\ y(1-y)\dfrac{\partial^2 z}{\partial y^2}+x(1-y)\dfrac{\partial^2 z}{\partial x \partial y}+[c-(a+b'+1)y]\dfrac{\partial z}{\partial y}-b'x\dfrac{\partial z}{\partial x}-ab'z=0, \end{cases}$$

which are equivalent to the following three equations:

$$\begin{cases} \dfrac{\partial^2 z}{\partial x^2}+\left(\dfrac{c-b'}{x}+\dfrac{a+b-c+1}{x-1}+\dfrac{b'}{x-y}\right)\dfrac{\partial z}{\partial x}-\dfrac{by(y-1)}{x(x-1)(x-y)}\dfrac{\partial z}{\partial y}+\dfrac{abz}{x(x-1)}=0, \\ \dfrac{\partial^2 z}{\partial y^2}+\left(\dfrac{c-b}{y}+\dfrac{a+b'-c+1}{y-1}+\dfrac{b}{y-x}\right)\dfrac{\partial z}{\partial y}-\dfrac{b'x(x-1)}{y(y-1)(y-x)}\dfrac{\partial z}{\partial x}+\dfrac{ab'z}{y(y-1)}=0, \\ \dfrac{\partial^2 z}{\partial x \partial y}=\dfrac{b'}{x-y}\dfrac{\partial z}{\partial x}-\dfrac{b}{x-y}\dfrac{\partial z}{\partial y}. \end{cases}$$

There are three linearly independent solutions z_1, z_2 and z_3. Set $u_1=\frac{z_1}{z_3}$ and $u_2=\frac{z_2}{z_3}$.

We find that (see [Y])

$$\begin{cases} \{u_1,u_2;x,y\}_x=\dfrac{by(y-1)}{x(x-1)(x-y)}, \\ \{u_1,u_2;x,y\}_y=\dfrac{b'x(x-1)}{y(y-1)(y-x)}, \\ [u_1,u_2;x,y]_x=\dfrac{c-b'}{x}+\dfrac{a+b-c+1}{x-1}+\dfrac{b'-2b}{x-y}, \\ [u_1,u_2;x,y]_y=\dfrac{c-b}{y}+\dfrac{a+b'-c+1}{y-1}+\dfrac{b-2b'}{y-x}. \end{cases}$$

Put

$$\Delta=\begin{vmatrix} \frac{\partial z_1}{\partial x} & \frac{\partial z_1}{\partial y} & z_1 \\ \frac{\partial z_2}{\partial x} & \frac{\partial z_2}{\partial y} & z_2 \\ \frac{\partial z_3}{\partial x} & \frac{\partial z_3}{\partial y} & z_3 \end{vmatrix}. \tag{2.1}$$

We have

$$\begin{aligned}\frac{\partial\Delta}{\partial x} &= \begin{vmatrix} \frac{\partial^2 z_1}{\partial x^2} & \frac{\partial z_1}{\partial y} & z_1 \\ \frac{\partial^2 z_2}{\partial x^2} & \frac{\partial z_2}{\partial y} & z_2 \\ \frac{\partial^2 z_3}{\partial x^2} & \frac{\partial z_3}{\partial y} & z_3 \end{vmatrix} + \begin{vmatrix} \frac{\partial z_1}{\partial x} & \frac{\partial^2 z_1}{\partial x\partial y} & z_1 \\ \frac{\partial z_2}{\partial x} & \frac{\partial^2 z_2}{\partial x\partial y} & z_2 \\ \frac{\partial z_3}{\partial x} & \frac{\partial^2 z_3}{\partial x\partial y} & z_3 \end{vmatrix} \\ &= -\left(\frac{c-b'}{x} + \frac{a+b-c+1}{x-1} + \frac{b'}{x-y}\right)\Delta - \frac{b}{x-y}\Delta \\ &= -\left(\frac{c-b'}{x} + \frac{a+b-c+1}{x-1} + \frac{b+b'}{x-y}\right)\Delta. \end{aligned} \tag{2.2}$$

Similarly,

$$\frac{\partial\Delta}{\partial y} = -\left(\frac{c-b}{y} + \frac{a+b'-c+1}{y-1} + \frac{b+b'}{y-x}\right)\Delta. \tag{2.3}$$

Hence,

$$\Delta = Cx^{b'-c}y^{b-c}(x-1)^{c-a-b-1}(y-1)^{c-a-b'-1}(x-y)^{-b-b'}. \tag{2.4}$$

On the other hand,

$$\Delta = z_3^2 \begin{vmatrix} \frac{\partial(z_1/z_3)}{\partial x} & \frac{\partial(z_1/z_3)}{\partial y} & z_1 \\ \frac{\partial(z_2/z_3)}{\partial x} & \frac{\partial(z_2/z_3)}{\partial y} & z_2 \\ 0 & 0 & z_3 \end{vmatrix} = z_3^3 \frac{\partial(\xi,\eta)}{\partial(x,y)}, \tag{2.5}$$

where $\xi = z_1/z_3$ and $\eta = z_2/z_3$.

Therefore,

$$z_3^3 \frac{\partial(\xi,\eta)}{\partial(x,y)} = Cx^{b'-c}y^{b-c}(x-1)^{c-a-b-1}(y-1)^{c-a-b'-1}(x-y)^{-b-b'}. \tag{2.6}$$

When ξ, η and $x^{b'-c}y^{b-c}(x-1)^{c-a-b-1}(y-1)^{c-a-b'-1}(x-y)^{-b-b'}$ are algebraic functions, (2.6) implies that z_3 is an algebraic function. Hence, z_1, z_2 and z_3 are all algebraic functions.

Theorem 2.1 (Main Theorem 1). *For the system of linear partial differential equations*:

$$\left\{\begin{aligned} &\frac{\partial^2 z}{\partial v_1^2} + \frac{1}{3}P_1\frac{\partial z}{\partial v_1} - F_1\frac{\partial z}{\partial v_2} + \left(\frac{\partial F_1}{\partial v_2} - \frac{1}{3}\frac{\partial P_1}{\partial v_1} - \frac{2}{9}P_1^2 - \frac{2}{3}F_1P_2\right)z = 0, \\ &\frac{\partial^2 z}{\partial v_1\partial v_2} - \frac{1}{3}P_2\frac{\partial z}{\partial v_1} - \frac{1}{3}P_1\frac{\partial z}{\partial v_2} + \left(\frac{1}{3}\frac{\partial P_2}{\partial v_1} + \frac{1}{3}\frac{\partial P_1}{\partial v_2} + \frac{1}{9}P_1P_2 - F_1F_2\right)z = 0, \\ &\frac{\partial^2 z}{\partial v_2^2} + \frac{1}{3}P_2\frac{\partial z}{\partial v_2} - F_2\frac{\partial z}{\partial v_1} + \left(\frac{\partial F_2}{\partial v_1} - \frac{1}{3}\frac{\partial P_2}{\partial v_2} - \frac{2}{9}P_2^2 - \frac{2}{3}F_2P_1\right)z = 0, \end{aligned}\right. \tag{2.7}$$

put

$$\begin{cases} P_1 = \dfrac{c-b'}{v_1} + \dfrac{a+b-c+1}{v_1-1} + \dfrac{b'-2b}{v_1-v_2}, \\ P_2 = \dfrac{c-b}{v_2} + \dfrac{a+b'-c+1}{v_2-1} + \dfrac{b-2b'}{v_2-v_1}, \\ F_1 = \dfrac{bv_2(v_2-1)}{v_1(v_1-1)(v_1-v_2)}, \\ F_2 = \dfrac{b'v_1(v_1-1)}{v_2(v_2-1)(v_2-v_1)}. \end{cases} \tag{2.8}$$

Set

$$w = v_1^{-\frac{1}{3}(c-b')} v_2^{-\frac{1}{3}(c-b)} (v_1-1)^{-\frac{1}{3}(a+b-c+1)} (v_2-1)^{-\frac{1}{3}(a+b'-c+1)} (v_1-v_2)^{-\frac{1}{3}(b+b')} z, \tag{2.9}$$

then the function $w = w(v_1, v_2)$ *satisfies the following equations*:

$$\begin{cases} \dfrac{\partial^2 w}{\partial v_1^2} + \left(\dfrac{c-b'}{v_1} + \dfrac{a+b-c+1}{v_1-1} + \dfrac{b'}{v_1-v_2}\right)\dfrac{\partial w}{\partial v_1} - \dfrac{bv_2(v_2-1)}{v_1(v_1-1)(v_1-v_2)}\dfrac{\partial w}{\partial v_2} + \\ + \dfrac{ab}{v_1(v_1-1)} w = 0, \\ \dfrac{\partial^2 w}{\partial v_2^2} + \left(\dfrac{c-b}{v_2} + \dfrac{a+b'-c+1}{v_2-1} + \dfrac{b}{v_2-v_1}\right)\dfrac{\partial w}{\partial v_2} - \dfrac{b'v_1(v_1-1)}{v_2(v_2-1)(v_2-v_1)}\dfrac{\partial w}{\partial v_1} + \\ + \dfrac{ab'}{v_2(v_2-1)} w = 0, \\ \dfrac{\partial^2 w}{\partial v_1 \partial v_2} = \dfrac{b'}{v_1-v_2}\dfrac{\partial w}{\partial v_1} - \dfrac{b}{v_1-v_2}\dfrac{\partial w}{\partial v_2}. \end{cases} \tag{2.10}$$

A solution is given by

$$w = F_1(a; b, b'; c; v_1, v_2), \tag{2.11}$$

where $F_1 = F_1(a; b, b'; c; x, y)$ *is the Appell hypergeometric function* (*see* [AK]). *Consequently,*

$$\begin{aligned} z =& v_1^{\frac{1}{3}(c-b')} v_2^{\frac{1}{3}(c-b)} (v_1-1)^{\frac{1}{3}(a+b-c+1)} (v_2-1)^{\frac{1}{3}(a+b'-c+1)} (v_1-v_2)^{\frac{1}{3}(b+b')} \times \\ & \times F_1(a; b, b'; c; v_1, v_2). \end{aligned} \tag{2.12}$$

Proof. Note that

$$z = v_1^{\frac{1}{3}(c-b')} v_2^{\frac{1}{3}(c-b)} (v_1-1)^{\frac{1}{3}(a+b-c+1)} (v_2-1)^{\frac{1}{3}(a+b'-c+1)} (v_1-v_2)^{\frac{1}{3}(b+b')} w.$$

We have

$$\begin{aligned}\frac{\partial z}{\partial v_1} =&v_1^{\frac{1}{3}(c-b')}v_2^{\frac{1}{3}(c-b)}(v_1-1)^{\frac{1}{3}(a+b-c+1)}(v_2-1)^{\frac{1}{3}(a+b'-c+1)}(v_1-v_2)^{\frac{1}{3}(b+b')}\times\\ &\times\left[\frac{\partial w}{\partial v_1}+\left(\frac{1}{3}P_1+\frac{b}{v_1-v_2}\right)w\right].\end{aligned}$$

$$\begin{aligned}\frac{\partial z}{\partial v_2} =&v_1^{\frac{1}{3}(c-b')}v_2^{\frac{1}{3}(c-b)}(v_1-1)^{\frac{1}{3}(a+b-c+1)}(v_2-1)^{\frac{1}{3}(a+b'-c+1)}(v_1-v_2)^{\frac{1}{3}(b+b')}\times\\ &\times\left[\frac{\partial w}{\partial v_2}+\left(\frac{1}{3}P_2+\frac{b'}{v_2-v_1}\right)w\right].\end{aligned}$$

$$\begin{aligned}&\frac{\partial^2 z}{\partial v_1^2} = v_1^{\frac{1}{3}(c-b')}v_2^{\frac{1}{3}(c-b)}(v_1-1)^{\frac{1}{3}(a+b-c+1)}(v_2-1)^{\frac{1}{3}(a+b'-c+1)}(v_1-v_2)^{\frac{1}{3}(b+b')}\times\\ &\times\left[\frac{\partial^2 w}{\partial v_1^2}+2(\frac{1}{3}P_1+\frac{b}{v_1-v_2})\frac{\partial w}{\partial v_1}+((\frac{1}{3}P_1+\frac{b}{v_1-v_2})^2+\frac{1}{3}\frac{\partial P_1}{\partial v_1}+\frac{-b}{(v_1-v_2)^2})w\right].\end{aligned}$$

$$\begin{aligned}&\frac{\partial^2 z}{\partial v_2^2} = v_1^{\frac{1}{3}(c-b')}v_2^{\frac{1}{3}(c-b)}(v_1-1)^{\frac{1}{3}(a+b-c+1)}(v_2-1)^{\frac{1}{3}(a+b'-c+1)}(v_1-v_2)^{\frac{1}{3}(b+b')}\times\\ &\times\left[\frac{\partial^2 w}{\partial v_2^2}+2(\frac{1}{3}P_2+\frac{b'}{v_2-v_1})\frac{\partial w}{\partial v_2}+((\frac{1}{3}P_2+\frac{b'}{v_2-v_1})^2+\frac{1}{3}\frac{\partial P_2}{\partial v_2}+\frac{-b'}{(v_2-v_1)^2})w\right].\end{aligned}$$

$$\begin{aligned}\frac{\partial^2 z}{\partial v_1\partial v_2} =&v_1^{\frac{1}{3}(c-b')}v_2^{\frac{1}{3}(c-b)}(v_1-1)^{\frac{1}{3}(a+b-c+1)}(v_2-1)^{\frac{1}{3}(a+b'-c+1)}(v_1-v_2)^{\frac{1}{3}(b+b')}\times\\ &\times[\frac{\partial^2 w}{\partial v_1\partial v_2}+(\frac{1}{3}P_2+\frac{b'}{v_2-v_1})\frac{\partial w}{\partial v_1}+(\frac{1}{3}P_1+\frac{b}{v_1-v_2})\frac{\partial w}{\partial v_2}+\\ &+((\frac{1}{3}P_1+\frac{b}{v_1-v_2})(\frac{1}{3}P_2+\frac{b'}{v_2-v_1})+\frac{1}{3}\frac{\partial P_1}{\partial v_2}+\frac{b}{(v_1-v_2)^2})w].\end{aligned}$$

The equation

$$\frac{\partial^2 z}{\partial v_1^2}+\frac{1}{3}P_1\frac{\partial z}{\partial v_1}-F_1\frac{\partial z}{\partial v_2}+\left(\frac{\partial F_1}{\partial v_2}-\frac{1}{3}\frac{\partial P_1}{\partial v_1}-\frac{2}{9}P_1^2-\frac{2}{3}F_1P_2\right)z=0$$

gives that

$$\begin{aligned}&\frac{\partial^2 w}{\partial v_1^2}+\left(P_1+\frac{2b}{v_1-v_2}\right)\frac{\partial w}{\partial v_1}-F_1\frac{\partial w}{\partial v_2}+\\ &+\left[\frac{b}{v_1-v_2}P_1+\frac{b^2-b}{(v_1-v_2)^2}-F_1P_2-\frac{b'}{v_2-v_1}F_1+\frac{\partial F_1}{\partial v_2}\right]w=0.\end{aligned}$$

Here,

$$\frac{\partial F_1}{\partial v_2}=\frac{b}{(v_1-v_2)^2}-\frac{b}{v_1(v_1-1)}.$$

Thus,

$$\begin{aligned}&\frac{b}{v_1-v_2}P_1+\frac{b^2-b}{(v_1-v_2)^2}-F_1P_2-\frac{b'}{v_2-v_1}F_1+\frac{\partial F_1}{\partial v_2}\\=&\frac{b}{v_1-v_2}\left(P_1+\frac{b}{v_1-v_2}\right)-F_1\left(P_2+\frac{b'}{v_2-v_1}\right)+\frac{-b}{(v_1-v_2)^2}+\frac{\partial F_1}{\partial v_2}\\=&\frac{b}{v_1-v_2}\left(\frac{c-b'}{v_1}+\frac{a+b-c+1}{v_1-1}+\frac{b'-b}{v_1-v_2}\right)+\\&-\frac{bv_2(v_2-1)}{v_1(v_1-1)(v_1-v_2)}\left(\frac{c-b}{v_2}+\frac{a+b'-c+1}{v_2-1}+\frac{b-b'}{v_2-v_1}\right)-\frac{b}{v_1(v_1-1)}\\=&\frac{(a+1)b}{v_1(v_1-1)}-\frac{b}{v_1(v_1-1)}\\=&\frac{ab}{v_1(v_1-1)}.\end{aligned}$$

Hence, we get the following equation:

$$\begin{aligned}&\frac{\partial^2 w}{\partial v_1^2}+\left(\frac{c-b'}{v_1}+\frac{a+b-c+1}{v_1-1}+\frac{b'}{v_1-v_2}\right)\frac{\partial w}{\partial v_1}-\frac{bv_2(v_2-1)}{v_1(v_1-1)(v_1-v_2)}\frac{\partial w}{\partial v_2}+\\&+\frac{ab}{v_1(v_1-1)}w=0.\end{aligned}$$

Exchange the position of v_1 and v_2, we get the other equation.

The equation

$$\frac{\partial^2 z}{\partial v_1\partial v_2}-\frac{1}{3}P_2\frac{\partial z}{\partial v_1}-\frac{1}{3}P_1\frac{\partial z}{\partial v_2}+\left(\frac{1}{3}\frac{\partial P_2}{\partial v_1}+\frac{1}{3}\frac{\partial P_1}{\partial v_2}+\frac{1}{9}P_1P_2-F_1F_2\right)z=0$$

gives that

$$\frac{\partial^2 w}{\partial v_1\partial v_2}+\frac{b'}{v_2-v_1}\frac{\partial w}{\partial v_1}+\frac{b}{v_1-v_2}\frac{\partial w}{\partial v_2}+\left[\frac{b-bb'}{(v_1-v_2)^2}+\frac{2}{3}\frac{\partial P_1}{\partial v_2}+\frac{1}{3}\frac{\partial P_2}{\partial v_1}-F_1F_2\right]w=0.$$

Note that

$$\frac{\partial P_1}{\partial v_2}=\frac{b'-2b}{(v_1-v_2)^2},\quad\frac{\partial P_2}{\partial v_1}=\frac{b-2b'}{(v_2-v_1)^2},\quad F_1F_2=-\frac{bb'}{(v_1-v_2)^2},$$

we find that

$$\frac{b-bb'}{(v_1-v_2)^2}+\frac{2}{3}\frac{\partial P_1}{\partial v_2}+\frac{1}{3}\frac{\partial P_2}{\partial v_1}-F_1F_2=0.$$

Thus,

$$\frac{\partial^2 w}{\partial v_1\partial v_2}=\frac{b'}{v_1-v_2}\frac{\partial w}{\partial v_1}-\frac{b}{v_1-v_2}\frac{\partial w}{\partial v_2}.$$

It is known that the equations

$$F_1:\begin{cases} x(1-x)\dfrac{\partial^2 z}{\partial x^2}+y(1-x)\dfrac{\partial^2 z}{\partial x\partial y}+[c-(a+b+1)x]\dfrac{\partial z}{\partial x}-by\dfrac{\partial z}{\partial y}-abz=0,\\ y(1-y)\dfrac{\partial^2 z}{\partial y^2}+x(1-y)\dfrac{\partial^2 z}{\partial x\partial y}+[c-(a+b'+1)y]\dfrac{\partial z}{\partial y}-b'x\dfrac{\partial z}{\partial x}-ab'z=0\end{cases}$$

are equivalent to the following three equations:

$$\begin{cases}\dfrac{\partial^2 z}{\partial x^2}+\left(\dfrac{c-b'}{x}+\dfrac{a+b-c+1}{x-1}+\dfrac{b'}{x-y}\right)\dfrac{\partial z}{\partial x}-\dfrac{by(y-1)}{x(x-1)(x-y)}\dfrac{\partial z}{\partial y}+\dfrac{abz}{x(x-1)}=0,\\ \dfrac{\partial^2 z}{\partial y^2}+\left(\dfrac{c-b}{y}+\dfrac{a+b'-c+1}{y-1}+\dfrac{b}{y-x}\right)\dfrac{\partial z}{\partial y}-\dfrac{b'x(x-1)}{y(y-1)(y-x)}\dfrac{\partial z}{\partial x}+\dfrac{ab'z}{y(y-1)}=0,\\ \dfrac{\partial^2 z}{\partial x\partial y}=\dfrac{b'}{x-y}\dfrac{\partial z}{\partial x}-\dfrac{b}{x-y}\dfrac{\partial z}{\partial y},\end{cases}$$

where $F_1=F_1(a;b,b';c;x,y)$ is the Appell hypergeometric function (see [AK]). A solution is given by

$$w=F_1(a;b,b';c;v_1,v_2).$$

Consequently,

$$\begin{aligned}z=&v_1^{\frac{1}{3}(c-b')}v_2^{\frac{1}{3}(c-b)}(v_1-1)^{\frac{1}{3}(a+b-c+1)}(v_2-1)^{\frac{1}{3}(a+b'-c+1)}(v_1-v_2)^{\frac{1}{3}(b+b')}\times\\ &\times F_1(a;b,b';c;v_1,v_2).\end{aligned}$$

□

3. The algebraic solutions of Appell hypergeometric partial differential equations

Let us recall some basic facts about Appell-Lauricella functions (see [CW1], [CW2] or [DM1], [DM2], [M]). With the usual notation $(a,0):=1$ and $(a,n):=a(a+1)\cdots(a+n-1)$ for $a\in\mathbb{C}$, $n\in\mathbb{N}$, let us define the Appell-Lauricella function F_1 of N complex variables $x_2,\cdots,x_{N+1}$ with parameters $a,b_2,\cdots,b_{N+1}$ and $c\in\mathbb{C}$, $c\neq 0,-1,-2,\cdots$, by the series:

$$F_1(a,b_2,\cdots,b_{N+1};c;x_2,\cdots,x_{N+1})=\sum\frac{(a,\sum_j n_j)\prod_j(b_j,n_j)}{(c,\sum_j n_j)\prod_j(1,n_j)}\prod_j x_j^{n_j},$$

where j runs from 2 to $N+1$, each n_j runs from 0 to ∞, and the series converges if all $|x_j| < 1$. Almost everywhere we will use instead of this series its integral representation:

$$\frac{1}{B(1-\mu_1, 1-\mu_{N+2})} \int u^{-\mu_0}(u-1)^{-\mu_1} \prod_{j=2}^{N+1} (u-x_j)^{-\mu_j}\, du,$$

where the exponential parameters $\mu_0, \mu_1, \cdots, \mu_{N+2}$, are related to a, $b_2, \cdots, b_{N+1}$, c by

$$\begin{aligned} &\mu_j = b_j \quad \text{for} \quad j = 2, \cdots, N+1, \\ &\mu_0 = c - \sum b_j, \quad \mu_1 = 1 + a - c, \\ &\sum_{j=0}^{N+2} \mu_j = 2, \quad \text{i.e.} \quad \mu_{N+2} = 1 - a. \end{aligned}$$

THEOREM (see [CW1], Theorem 1). *Assume all $\mu_0, \cdots, \mu_{N+2} \in \mathbb{Q} - \mathbb{Z}$. Then there are no algebraic F_1 in more than three variables. The function $F_1(x, y)$ is algebraic if and only if $\mu_0, \cdots, \mu_4$ are up to permutations and a common change of the sign $\equiv \frac{1}{3}, \frac{1}{6}, \frac{1}{6}, \frac{1}{6}, \frac{1}{6}$ mod $\mathbb{Z}$. Then Δ is the symmetry group of the extended Hesse polytope of order* 1296*, projectively isomorphic to a subgroup of order* 216 *of the group $PSL(3, \mathbb{F}_3)$. The function $F_1(x, y, z)$ is algebraic if and only if $\mu_0, \cdots, \mu_5$ are all $\equiv \frac{1}{6}$ mod $\mathbb{Z}$ or all $\equiv -\frac{1}{6}$ mod $\mathbb{Z}$. Then Δ is the symmetry group of the Witting polytope of order* 155520*, projectively isomorphic to the group $PSp(4, \mathbb{F}_3)$ of order* 25920.

In the case $N = 2$ this means that $F_1(a, b, b'; c; x, y)$ is algebraic if and only if up to a common change of sign

$$\begin{array}{ccrrrrrl} a & \equiv & -\frac{1}{6} & -\frac{1}{6} & -\frac{1}{6} & -\frac{1}{6} & -\frac{1}{3} & \text{mod } \mathbb{Z} \\ b & \equiv & \frac{1}{6} & \frac{1}{6} & \frac{1}{3} & \frac{1}{6} & \frac{1}{6} & \text{mod } \mathbb{Z} \\ b' & \equiv & \frac{1}{6} & \frac{1}{6} & \frac{1}{6} & \frac{1}{3} & \frac{1}{6} & \text{mod } \mathbb{Z} \\ c & \equiv & -\frac{1}{3} & \frac{1}{2} & -\frac{1}{3} & -\frac{1}{3} & \frac{1}{2} & \text{mod } \mathbb{Z} \end{array}$$

For

$$(3.1) \qquad \begin{cases} P_1 = \dfrac{c-b'}{v_1} + \dfrac{a+b-c+1}{v_1-1} + \dfrac{b'-2b}{v_1-v_2}, \\ P_2 = \dfrac{c-b}{v_2} + \dfrac{a+b'-c+1}{v-2-1} + \dfrac{b-2b'}{v_2-v_1}, \\ F_1 = \dfrac{bv_2(v_2-1)}{v_1(v_1-1)(v_1-v_2)}, \\ F_2 = \dfrac{b'v_1(v_1-1)}{v_2(v_2-1)(v_2-v_1)}, \end{cases}$$

set

$$\alpha_1 = c - b', \quad \beta_1 = a + b - c + 1, \quad \gamma_1 = b' - 2b,$$
$$\alpha_2 = c - b, \quad \beta_2 = a + b' - c + 1, \quad \gamma_2 = b - 2b'.$$

We find that

$$\alpha_1 - \alpha_2 = b - b', \quad \beta_1 - \beta_2 = b - b', \quad \gamma_1 - \gamma_2 = -3(b - b').$$

This leads us to consider the double-triangular s-functions:

$$s_1 = s_1(\alpha_1, \beta_1, \gamma_1; \alpha_2, \beta_2, \gamma_2; v_1, v_2), \quad s_2 = s_2(\alpha_1, \beta_1, \gamma_1; \alpha_2, \beta_2, \gamma_2; v_1, v_2), \tag{3.2}$$

where

$$\alpha_1 - \alpha_2 = \beta_1 - \beta_2 = -\frac{1}{3}(\gamma_1 - \gamma_2).$$

In particular, when $\alpha_1 = \alpha_2$ or $\beta_1 = \beta_2$ or $\gamma_1 = \gamma_2$, they reduce the triangular s-functions (see [Y]):

$$s_1 = s_1(\alpha, \beta, \gamma; v_1, v_2), \quad s_2 = s_2(\alpha, \beta, \gamma; v_1, v_2). \tag{3.3}$$

This gives the following five cases:

(1) $a = -\frac{1}{6}$, $b = \frac{1}{6}$, $b' = \frac{1}{6}$, $c = -\frac{1}{3}$.

$$\begin{cases} P_1(v_1, v_2) = \dfrac{-\frac{1}{2}}{v_1} + \dfrac{\frac{4}{3}}{v_1 - 1} + \dfrac{-\frac{1}{6}}{v_1 - v_2}, \\ P_2(v_1, v_2) = \dfrac{-\frac{1}{2}}{v_2} + \dfrac{\frac{4}{3}}{v_2 - 1} + \dfrac{-\frac{1}{6}}{v_2 - v_1}, \\ F_1(v_1, v_2) = \dfrac{\frac{1}{6} v_2(v_2 - 1)}{v_1(v_1 - 1)(v_1 - v_2)}, \\ F_2(v_1, v_2) = \dfrac{\frac{1}{6} v_1(v_1 - 1)}{v_2(v_2 - 1)(v_2 - v_1)}. \end{cases} \tag{3.4}$$

Correspondingly,

$$z = v_1^{-\frac{1}{6}} v_2^{-\frac{1}{6}} (v_1 - 1)^{\frac{4}{9}} (v_2 - 1)^{\frac{4}{9}} (v_1 - v_2)^{\frac{1}{9}} F_1\left(-\frac{1}{6}; \frac{1}{6}, \frac{1}{6}; -\frac{1}{3}; v_1, v_2\right).$$

The triangular s-functions are:

$$s_1 = s_1\left(-\frac{1}{2}, \frac{4}{3}, -\frac{1}{6}; v_1, v_2\right), \quad s_2 = s_2\left(-\frac{1}{2}, \frac{4}{3}, -\frac{1}{6}; v_1, v_2\right).$$

(2) $a=-\frac{1}{6}$, $b=\frac{1}{6}$, $b'=\frac{1}{6}$, $c=\frac{1}{2}$.

$$\begin{cases} P_1(v_1,v_2)=\dfrac{\frac{1}{3}}{v_1}+\dfrac{\frac{1}{2}}{v_1-1}+\dfrac{-\frac{1}{6}}{v_1-v_2}, \\ P_2(v_1,v_2)=\dfrac{\frac{1}{3}}{v_2}+\dfrac{\frac{1}{2}}{v_2-1}+\dfrac{-\frac{1}{6}}{v_2-v_1}, \\ F_1(v_1,v_2)=\dfrac{\frac{1}{6}v_2(v_2-1)}{v_1(v_1-1)(v_1-v_2)}, \\ F_2(v_1,v_2)=\dfrac{\frac{1}{6}v_1(v_1-1)}{v_2(v_2-1)(v_2-v_1)}. \end{cases} \tag{3.5}$$

Correspondingly,

$$z=v_1^{\frac{1}{9}}v_2^{\frac{1}{9}}(v_1-1)^{\frac{1}{6}}(v_2-1)^{\frac{1}{6}}(v_1-v_2)^{\frac{1}{9}}F_1\left(-\frac{1}{6};\frac{1}{6},\frac{1}{6};\frac{1}{2};v_1,v_2\right).$$

The triangular s-functions are:

$$s_1=s_1\left(\frac{1}{3},\frac{1}{2},-\frac{1}{6};v_1,v_2\right),\quad s_2=s_2\left(\frac{1}{3},\frac{1}{2},-\frac{1}{6};v_1,v_2\right).$$

(3) $a=-\frac{1}{6}$, $b=\frac{1}{3}$, $b'=\frac{1}{6}$, $c=-\frac{1}{3}$.

$$\begin{cases} P_1(v_1,v_2)=\dfrac{-\frac{1}{2}}{v_1}+\dfrac{\frac{3}{2}}{v_1-1}+\dfrac{-\frac{1}{2}}{v_1-v_2}, \\ P_2(v_1,v_2)=\dfrac{-\frac{2}{3}}{v_2}+\dfrac{\frac{4}{3}}{v_2-1}, \\ F_1(v_1,v_2)=\dfrac{\frac{1}{3}v_2(v_2-1)}{v_1(v_1-1)(v_1-v_2)}, \\ F_2(v_1,v_2)=\dfrac{\frac{1}{6}v_1(v_1-1)}{v_2(v_2-1)(v_2-v_1)}. \end{cases} \tag{3.6}$$

Correspondingly,

$$z=v_1^{-\frac{1}{6}}v_2^{-\frac{2}{9}}(v_1-1)^{\frac{1}{2}}(v_2-1)^{\frac{4}{9}}(v_1-v_2)^{\frac{1}{6}}F_1\left(-\frac{1}{6};\frac{1}{3},\frac{1}{6};-\frac{1}{3};v_1,v_2\right).$$

The double-triangular s-functions are:

$$s_1=s_1\left(-\frac{1}{2},\frac{3}{2},-\frac{1}{2};-\frac{2}{3},\frac{4}{3},0;v_1,v_2\right),\quad s_2=s_2\left(-\frac{1}{2},\frac{3}{2},-\frac{1}{2};-\frac{2}{3},\frac{4}{3},0;v_1,v_2\right).$$

(4) $a = -\frac{1}{6}$, $b = \frac{1}{6}$, $b' = \frac{1}{3}$, $c = -\frac{1}{3}$.

$$\begin{cases} P_1(v_1, v_2) = \dfrac{-\frac{2}{3}}{v_1} + \dfrac{\frac{4}{3}}{v_1 - 1}, \\ P_2(v_1, v_2) = \dfrac{-\frac{1}{2}}{v_2} + \dfrac{\frac{3}{2}}{v_2 - 1} + \dfrac{-\frac{1}{2}}{v_2 - v_1}, \\ F_1(v_1, v_2) = \dfrac{\frac{1}{6}v_2(v_2 - 1)}{v_1(v_1 - 1)(v_1 - v_2)}, \\ F_2(v_1, v_2) = \dfrac{\frac{1}{3}v_1(v_1 - 1)}{v_2(v_2 - 1)(v_2 - v_1)}. \end{cases} \tag{3.7}$$

Correspondingly,

$$z = v_1^{-\frac{2}{9}} v_2^{-\frac{1}{6}} (v_1 - 1)^{\frac{4}{9}} (v_2 - 1)^{\frac{1}{2}} (v_1 - v_2)^{\frac{1}{6}} F_1\left(-\frac{1}{6}; \frac{1}{6}, \frac{1}{3}; -\frac{1}{3}; v_1, v_2\right).$$

The double-triangular s-functions are:

$$s_1 = s_1\left(-\frac{2}{3}, \frac{4}{3}, 0; -\frac{1}{2}, \frac{3}{2}, -\frac{1}{2}; v_1, v_2\right), \quad s_2 = s_2\left(-\frac{2}{3}, \frac{4}{3}, 0; -\frac{1}{2}, \frac{3}{2}, -\frac{1}{2}; v_1, v_2\right).$$

(5) $a = -\frac{1}{3}$, $b = \frac{1}{6}$, $b' = \frac{1}{6}$, $c = \frac{1}{2}$.

$$\begin{cases} P_1(v_1, v_2) = \dfrac{\frac{1}{3}}{v_1} + \dfrac{\frac{1}{3}}{v_1 - 1} + \dfrac{-\frac{1}{6}}{v_1 - v_2}, \\ P_2(v_1, v_2) = \dfrac{\frac{1}{3}}{v_2} + \dfrac{\frac{1}{3}}{v_2 - 1} + \dfrac{-\frac{1}{6}}{v_2 - v_1}, \\ F_1(v_1, v_2) = \dfrac{\frac{1}{6}v_2(v_2 - 1)}{v_1(v_1 - 1)(v_1 - v_2)}, \\ F_2(v_1, v_2) = \dfrac{\frac{1}{6}v_1(v_1 - 1)}{v_2(v_2 - 1)(v_2 - v_1)}. \end{cases} \tag{3.8}$$

Correspondingly,

$$z = v_1^{\frac{1}{9}} v_2^{\frac{1}{9}} (v_1 - 1)^{\frac{1}{9}} (v_2 - 1)^{\frac{1}{9}} (v_1 - v_2)^{\frac{1}{9}} F_1\left(-\frac{1}{3}; \frac{1}{6}, \frac{1}{6}; \frac{1}{2}; v_1, v_2\right).$$

The triangular s-functions are:

$$s_1 = s_1\left(\frac{1}{3}, \frac{1}{3}, -\frac{1}{6}; v_1, v_2\right), \quad s_2 = s_2\left(\frac{1}{3}, \frac{1}{3}, -\frac{1}{6}; v_1, v_2\right).$$

We will study the transformation between two Appell hypergeometric partial differential equations:

$$\begin{cases} x(1-x)\dfrac{\partial^2 z}{\partial x^2}+y(1-x)\dfrac{\partial^2 z}{\partial x\partial y}+[c-(a+b+1)x]\dfrac{\partial z}{\partial x}-by\dfrac{\partial z}{\partial y}-abz=0, \\ y(1-y)\dfrac{\partial^2 z}{\partial y^2}+x(1-y)\dfrac{\partial^2 z}{\partial x\partial y}+[c-(a+b'+1)y]\dfrac{\partial z}{\partial y}-b'x\dfrac{\partial z}{\partial x}-ab'z=0, \end{cases}$$

and

$$\begin{cases} u_1(1-u_1)\dfrac{\partial^2 v}{\partial u_1^2}+u_2(1-u_1)\dfrac{\partial^2 v}{\partial u_1\partial u_2}+[\gamma-(\alpha+\beta+1)u_1]\dfrac{\partial v}{\partial u_1}-\beta u_2\dfrac{\partial v}{\partial u_2}+ \\ -\alpha\beta v=0, \\ u_2(1-u_2)\dfrac{\partial^2 v}{\partial u_2^2}+u_1(1-u_2)\dfrac{\partial^2 v}{\partial u_1\partial u_2}+[\gamma-(\alpha+\beta'+1)u_2]\dfrac{\partial v}{\partial u_2}-\beta' u_1\dfrac{\partial v}{\partial u_1}+ \\ -\alpha\beta' v=0. \end{cases}$$

Here, $u_1=u_1(x,y)$, $u_2=u_2(x,y)$ and $z=wv$ with $w=w(x,y)$.

We have

$$\frac{\partial z}{\partial x}=\frac{\partial w}{\partial x}v+w\left(\frac{\partial v}{\partial u_1}\frac{\partial u_1}{\partial x}+\frac{\partial v}{\partial u_2}\frac{\partial u_2}{\partial x}\right),\quad \frac{\partial z}{\partial y}=\frac{\partial w}{\partial y}v+w\left(\frac{\partial v}{\partial u_1}\frac{\partial u_1}{\partial y}+\frac{\partial v}{\partial u_2}\frac{\partial u_2}{\partial y}\right).$$

For $z_1=wv_1$, $z_2=wv_2$ and $z_3=wv_3$, we find that

$$\begin{vmatrix} \frac{\partial z_1}{\partial x} & \frac{\partial z_1}{\partial y} & z_1 \\ \frac{\partial z_2}{\partial x} & \frac{\partial z_2}{\partial y} & z_2 \\ \frac{\partial z_3}{\partial x} & \frac{\partial z_3}{\partial y} & z_3 \end{vmatrix} = w^3\frac{\partial(u_1,u_2)}{\partial(x,y)}\begin{vmatrix} \frac{\partial v_1}{\partial u_1} & \frac{\partial v_1}{\partial u_2} & v_1 \\ \frac{\partial v_2}{\partial u_1} & \frac{\partial v_2}{\partial u_2} & v_2 \\ \frac{\partial v_3}{\partial u_1} & \frac{\partial v_3}{\partial u_2} & v_3 \end{vmatrix}. \tag{3.9}$$

On the other hand,

$$\begin{vmatrix} \frac{\partial z_1}{\partial x} & \frac{\partial z_1}{\partial y} & z_1 \\ \frac{\partial z_2}{\partial x} & \frac{\partial z_2}{\partial y} & z_2 \\ \frac{\partial z_3}{\partial x} & \frac{\partial z_3}{\partial y} & z_3 \end{vmatrix} = C_1x^{b'-c}y^{b-c}(x-1)^{c-a-b-1}(y-1)^{c-a-b'-1}(x-y)^{-b-b'}. \tag{3.10}$$

(3.11)

$$\begin{vmatrix} \frac{\partial v_1}{\partial u_1} & \frac{\partial v_1}{\partial u_2} & v_1 \\ \frac{\partial v_2}{\partial u_1} & \frac{\partial v_2}{\partial u_2} & v_2 \\ \frac{\partial v_3}{\partial u_1} & \frac{\partial v_3}{\partial u_2} & v_3 \end{vmatrix} = C_2u_1^{\beta'-\gamma}u_2^{\beta-\gamma}(u_1-1)^{\gamma-\alpha-\beta-1}(u_2-1)^{\gamma-\alpha-\beta'-1}(u_1-u_2)^{-\beta-\beta'}.$$

Thus,

$$w^3 = C\frac{x^{b'-c}y^{b-c}(x-1)^{c-a-b-1}(y-1)^{c-a-b'-1}(x-y)^{-b-b'}}{u_1^{\beta'-\gamma}u_2^{\beta-\gamma}(u_1-1)^{\gamma-\alpha-\beta-1}(u_2-1)^{\gamma-\alpha-\beta'-1}(u_1-u_2)^{-\beta-\beta'}}\frac{1}{\frac{\partial(u_1,u_2)}{\partial(x,y)}}. \tag{3.12}$$

Here, C_1, C_2 and C are three constants.

Note that

$$\frac{\partial(\xi,\eta)}{\partial(x,y)} = \frac{1}{z_3^3}Cx^{b'-c}y^{b-c}(x-1)^{c-a-b-1}(y-1)^{c-a-b'-1}(x-y)^{-b-b'}. \tag{3.13}$$

We will study the following equations:

$$\frac{u_1(\xi,\eta)}{u_3(\xi,\eta)} = R_1(x,y), \quad \frac{u_2(\xi,\eta)}{u_3(\xi,\eta)} = R_2(x,y). \tag{3.14}$$

We find that

$$\begin{aligned} \frac{\partial(R_1,R_2)}{\partial(x,y)} &= \frac{\partial(u_1/u_3,u_2/u_3)}{\partial(\xi,\eta)}\frac{\partial(\xi,\eta)}{\partial(x,y)} \\ &= \frac{\partial(u_1/u_3,u_2/u_3)}{\partial(\xi,\eta)}\frac{1}{z_3^3}Cx^{b'-c}y^{b-c}(x-1)^{c-a-b-1}(y-1)^{c-a-b'-1}(x-y)^{-b-b'}. \end{aligned} \tag{3.15}$$

4. The Hessian polyhedral equations

A polytope is a geometrical figure bounded by portions of lines, planes, or hyperplanes; e.g., in two dimensions it is a polygon, in three a polyhedron. We define a polytope (see [Co1]) as a finite convex region of n-dimensional space enclosed by a finite number of hyperplanes. The part of the polytope that lies in one of the hyperplanes is called a cell. A polytope Π_n $(n > 2)$ is said to be regular if its cells are regular and there is a regular vertex figure at every vertex.

In his great monograph in 1852, Schläfli gave the n-dimensional Schläfli symbol (see [Co1]), which enables us to read off many properties of a regular polytope at a glance; e.g., the elements of $\{p,q,r,\cdots\}$, besides vertices and edges, are plane faces $\{p\}$, solid faces $\{p,q\}$, and so on. The general regular polytope $\{p,q,\cdots,v,w\}$ has cells $\{p,q,\cdots,v\}$ and vertex figures $\{q,\cdots,v,w\}$. In particular, for any $n > 1$,

$$\alpha_n = \{3,3,\cdots,3,3\} = \{3^{n-1}\},$$

$$\beta_n = \{3,3,\cdots,3,4\} = \{3^{n-2},4\},$$

$$\gamma_n = \{4, 3, \cdots, 3, 3\} = \{4, 3^{n-2}\}.$$

The Schläfli's criterion for the existence of a regular figure corresponding to a given symbol $\{p, q, \cdots, v, w\}$ is given by (see [Co1]):

$$\Delta_{p,q,\cdots,v,w} > 0.$$

Here,

$$\Delta_{p,q,\cdots,v,w} = \begin{vmatrix} 1 & c_1 & 0 & \cdots & 0 & 0 & 0 \\ c_1 & 1 & c_2 & \cdots & 0 & 0 & 0 \\ & \cdots & & & & \cdots & \\ 0 & 0 & 0 & \cdots & c_{n-2} & 1 & c_{n-1} \\ 0 & 0 & 0 & \cdots & 0 & c_{n-1} & 1 \end{vmatrix},$$

where

$$c_1 = \cos\frac{\pi}{p}, \quad c_2 = \cos\frac{\pi}{q}, \cdots, c_{n-2} = \cos\frac{\pi}{v}, \quad c_{n-1} = \cos\frac{\pi}{w}.$$

When $n = 3$, we have $\Delta_{p,q} > 0$, or

$$\frac{1}{p} + \frac{1}{q} > \frac{1}{2},$$

which admits the Platonic solids

$$\{3, 3\}, \quad \{3, 4\}, \quad \{4, 3\}, \quad \{3, 5\}, \quad \{5, 3\}.$$

When $n = 4$, we have $\Delta_{p,q,r} > 0$, or

$$\sin\frac{\pi}{p}\sin\frac{\pi}{r} > \cos\frac{\pi}{q},$$

which admits the six polytopes

$$\{3, 3, 3\}, \quad \{3, 3, 4\}, \quad \{4, 3, 3\}, \quad \{3, 4, 3\}, \quad \{3, 3, 5\}, \quad \{5, 3, 3\}.$$

When $n = 5$, we have $\Delta_{p,q,r,s} > 0$, or

$$\frac{\cos^2\frac{\pi}{q}}{\sin^2\frac{\pi}{p}} + \frac{\cos^2\frac{\pi}{r}}{\sin^2\frac{\pi}{s}} < 1,$$

which admits the three polytopes:

$$\alpha_5 = \{3, 3, 3, 3\}, \quad \beta_5 = \{3, 3, 3, 4\}, \quad \gamma_5 = \{4, 3, 3, 3\}.$$

Since the only regular polytopes in five dimensions are α_5, β_5, γ_5, it follows by induction that in more than five dimensions the only regular polytopes are α_n, β_n, γ_n.

It is well-known that the regular polyhedra are connected with the Gauss hypergeometric differential equation (see [Kl9] and [Kl10]). In contrast with the real regular polyhedra, we will give a story on the complex regular polyhedra, which are intimately connected with the Appell hypergeometric partial differential equations.

According to [Co2], in the complex affine plane with a unitary metric, let us define a polygon to be a finite figure consisting of points (not all on one line) called vertices, and lines (not all through one point) called edges, satisfying the following two conditions:

(1) Every edge is incident with at least two vertices, and every vertex with at least two edges.
(2) Any two vertices are connected by a chain of successively incident edges and vertices.

The group of all unitary transformations that preserve the incidences is called the symmetry group of the polygon. The figure consisting of a vertex and an incident edge is called a flag. The polygon is said to be regular if its symmetry group is transitive on its flags.

Extending the above ideas to unitary n-space, let us consider a finite non-empty set of subspaces, μ-flats ($0 \leq \mu < n$), with a relation of incidence, defined by saying that a μ-flat and a ν-flat are incident if one of them is a proper subspace of the other. A subset of this set of flats is said to be connected if any two of its flats can be joined by a chain of successively incident flats. If $\lambda < \nu - 1$, any incident λ-flat and ν-flat determine a medial figure consisting of all the μ-flats ($\lambda < \mu < \nu$) that are incident with both.

Such a set of flats is called a polytope if it satisfies the following two conditions:

(1) Every medial figure includes at least two μ-flats for each μ ($-1 \leq \lambda < \mu < \nu \leq n$).
(2) Every medial figure with $\lambda < \nu - 2$ is connected.

The group of all unitary transformations that permute the μ-flats for each μ and preserve the incidences is called the symmetry group of the polytope.

In an n-dimensional polytope Π_n, a μ-flat and all its incident λ-flats ($\lambda < \mu$) form a sub-polytope Π_μ, called a μ-dimensional element of Π_n. In particular there are vertices Π_0, edges Π_1, faces Π_2, $\cdots$, and cells Π_{n-1}. The figure consisting of one element of each dimension number, all mutually incident, is called a flag. The polytope is called regular if its symmetry group is transitive on its flags.

In fact, the symmetry group of a regular complex polytope is generated by a sequence of unitary reflections of periods p_1, p_2, $\cdots$, p_n, such that any two non-consecutive reflections are commutative. This leads to the generalized Schläfli symbol

$$p_1\{q_1\}p_2\{q_2\}\cdots p_{n-1}\{q_{n-1}\}p_n.$$

Let us recall some basic facts about the Hessian polyhedron (see [Co2]). In the case of the Hessian polyhedron 3{3}3{3}3, the Hermitian form is

$$x^1\overline{x^1} - \sqrt{\frac{1}{3}}(x^1\overline{x^2} + x^2\overline{x^1}) + x^2\overline{x^2} - \sqrt{\frac{1}{3}}(x^2\overline{x^3} + x^3\overline{x^2}) + x^3\overline{x^3} = u_1\overline{u_1} + u_2\overline{u_2} + u_3\overline{u_3},$$

where

$$u_1 = -x^1 + \frac{x^2}{\sqrt{3}}, \quad u_2 = \frac{x^2}{\sqrt{3}}, \quad u_3 = \frac{x^2}{\sqrt{3}} - x^3.$$

Thus the three mirrors are

$$u_1 = 0, \quad \frac{u_1 + u_2 + u_3}{\sqrt{3}} = 0, \quad u_3 = 0.$$

R_1 and R_3 multiply u_1 and u_3 (respectively) by $\omega = \exp(2\pi i/3)$ and R_2 is

$$u'_\lambda = u_\lambda - \frac{i\omega^2}{\sqrt{3}}(u_1 + u_2 + u_3).$$

In fact, 3{3}3{3}3 has just the 27 vertices:

$$(0, \omega^\mu, -\omega^\nu), \quad (-\omega^\nu, 0, \omega^\mu), \quad (\omega^\mu, -\omega^\nu, 0),$$

where μ and ν take (independently) the three values 0, 1, 2 or (equally well) 1, 2, 3. Making the latter choice, we may conveniently use the concise symbols:

$$0\mu\nu, \quad \nu 0\mu, \quad \mu\nu 0$$

for these 27 points.

Since the symbol 3{3}3{3}3 is a palindrome, this polyhedron is self-reciprocal. Having 27 vertices, it also has 27 faces 3{3}3. Since the vertex figure is another 3{3}3, there are 8 edges at each vertex, and 72 edges altogether.

Two vertices are naturally said to be adjacent if they belong to the same edge; their symbols agree in an even number of places, that is, in just two places or nowhere. The 72 edges occur in 36 pairs of opposites, such that any two vertices belonging to opposite edges are non-adjacent; their symbols agree in just one place.

Although the edges occur in pairs of opposites, the vertices obviously do not (as 27 is an odd number). In fact, 3{3}3{3}3 is not 2-symmetric but 3-symmetric: the 27 vertices lie by threes on nine diameters. The three vertices on any diameter are derived from one another by cyclically permuting the digits 1, 2, 3 wherever they occur.

Similarly, the 72 edges lie by sixes on twelve planes of symmetry

$$u_1 = 0, u_2 = 0, u_3 = 0, \omega^\lambda u_1 + \omega^\mu u_2 + \omega^\nu u_3 = 0 (\lambda, \mu, \nu = 0, 1, 2 \text{ independently}).$$

These planes include the mirrors for the generating reflections R_1, R_2, R_3; in fact, they are the mirrors for all the reflections that occur in 3[3]3[3]3.

Since

$$R_1R_2R_1 = R_2R_1R_2, \quad R_2R_3R_2 = R_3R_2R_3,$$

the generators R_1, R_2, R_3 are mutually conjugate; and every reflection in the group is conjugate to some $R_\nu^{\pm 1}$. Thus the section of 3{3}3{3}3 by any plane of symmetry is a polygon 3{4}2.

The nine diameters and twelve planes of symmetry are related in a remarkable way: every two diameters determine a plane of symmetry which contains a third diameter.

3{3}3{3}3 is called the Hessian polyhedron because the incidences of the diameters and planes of symmetry are the same as the incidences of the points and lines of the Hessian configuration

$$\begin{pmatrix} 9 & 4 \\ 3 & 12 \end{pmatrix}$$

or $(9_4, 12_3)$ (see [MBD]): nine points lying by threes on twelve lines, with four of the lines through each point. To make this transition from complex affine 3-space to the complex projective plane, we have to interpret the affine coordinates $(0, \omega^\mu, -\omega^\nu)$, $(-\omega^\nu, 0, \omega^\mu)$, $(\omega^\mu, -\omega^\nu, 0)$ for the 27 points $0\mu\nu$, $\nu 0\mu$, $\mu\nu 0$ as homogeneous coordinates:

$$\begin{array}{ccc} (0,1,-1) & (-1,0,1) & (1,-1,0) \\ (0,1,-\omega) & (-\omega,0,1) & (1,-\omega,0) \\ (0,1,-\omega^2) & (-\omega^2,0,1) & (1,-\omega^2,0) \end{array}$$

for the nine points given by

$$x^3 + y^3 + z^3 = xyz = 0.$$

These are the nine points of inflexion of the plane cubic curve

$$x^3 + y^3 + z^3 - 3axyz = 0 \quad (a \neq 0, 1)$$

whose Hessian is

$$x^3 + y^3 + z^3 + (a - 4a^{-2})xyz = 0.$$

In fact, the 27 lines on the cubic surface represent the 27 vertices of 3{3}3{3}3 in such a way that two of the lines are intersecting or skew according as the corresponding vertices are non-adjacent or adjacent. This leads to the question why the group of automorphisms of the 27 lines has order 51840 whereas the order of 3[3]3[3]3 is only 648. The explanation is that, among the pairs of non-adjacent vertices of 3{3}3{3}3, we have included pairs that belong to a diameter. In fact, there are altogether $9 + 36$ triads of mutually non-adjacent vertices: 9 lying on diameters, and 36, such as 120, 210, 330, forming equilateral triangles, three inscribed in each of the twelve γ_2^3's. However, on the cubic surface there is no corresponding distinction: the 45 triads of intersecting lines are all alike; their planes are just the 45 tritangent planes. The nine diameters correspond to one of the forty ways in which the 27 lines can be distributed into nine triads, each triad forming a tritangent plane. The representation of the nine diameters by the points of the configuration $(9_4, 12_3)$ corresponds to the fact that there are 12 ways of distributing these 9 tritangent planes into 3 Steiner trihedra so that each plane belongs to 4 of the trihedra.

Thus the group of automorphisms of the Hessian polyhedron 3{3}3{3}3 is a subgroup of index 40 in the group of automorphisms of the 27 lines.

When the three complex coordinates u_ν in unitary 3-space are replaced by six Cartesian coordinates in Euclidean 6-space, the polyhedron 3{3}3{3}3 yields the uniform polytope 2_{21} whose 27 vertices faithfully represent the 27 lines.

A polyhedron closely related to 3{3}3{3}3 is 2{4}3{3}3, for which the Hermitian form is

$$x^1\overline{x^1} - \sqrt{\frac{1}{2}}(x^1\overline{x^2} + x^2\overline{x^1}) + x^2\overline{x^2} - \sqrt{\frac{1}{3}}(x^2\overline{x^3} + x^3\overline{x^2}) + x^3\overline{x^3} = u_1\overline{u_1} + u_2\overline{u_2} + u_3\overline{u_3},$$

where

$$u_1 = -\frac{i\omega^2 x^1}{\sqrt{2}} + \frac{x^2}{\sqrt{3}}, \quad u_2 = \frac{i\omega x^1}{\sqrt{2}} + \frac{x^2}{\sqrt{3}}, \quad u_3 = \frac{x^2}{\sqrt{3}} - x^3.$$

The three mirrors are now

$$\frac{u_1 - \omega u_2}{\sqrt{2}} = 0, \quad \frac{u_1 + u_2 + u_3}{\sqrt{3}} = 0, \quad u_3 = 0.$$

R_2 and R_3 (of period 3) are the same as above, but R_1 (of period 2) is

$$u_1' = \omega u_2, \quad u_2' = \omega^2 u_1, \quad u_3' = u_3.$$

In fact, 2{4}3{3}3 has 54 vertices

$$(0, \pm\omega^\mu, \mp\omega^\nu), \quad (\mp\omega^\nu, 0, \pm\omega^\mu), \quad (\pm\omega^\mu, \mp\omega^\nu, 0),$$

namely the vertices of two reciprocal 3{3}3{3}3's of the same size, each derivable from the other by the central inversion

$$u_1' = -u_1, \quad u_2' = -u_2, \quad u_3' = -u_3.$$

There are 216 edges, each joining a vertex of one 3{3}3{3}3 to one of the 8 nearest vertices of the other. Since the vertex figure 3{3}3 has 8 vertices, the polyhedron has 8 edges at each vertex, 216 edges altogether. There are 72 faces 2{4}3.

The reciprocal polyhedron 3{3}3{4}2 has 72 vertices, 216 edges and 54 faces 3{3}3. Here, the vertices are:

$$(\omega^\lambda, \omega^\mu, \omega^\nu), \quad (-\omega^\lambda, -\omega^\mu, -\omega^\nu), \quad \text{and} \quad (\pm i\omega^\lambda\sqrt{3}, 0, 0), \quad \text{permuted,}$$

where $\lambda, \mu, \nu = 0, 1, 2$, independently.

Notice that 2{4}3{3}3 and 3{3}3{4}2 are 6-symmetric. In the case of 3{3}3{4}2, the six vertices $(\pm i\omega^\lambda\sqrt{3}, 0, 0)$ form a γ_1^6 on one of the twelve diameters.

These diameters distribute themselves into four triads of mutually orthogonal diameters, namely, lines joining the origin to the following triads of the points:

$$\begin{array}{ccc} (1,0,0), & (0,1,0), & (0,0,1); \\ (1,1,1), & (\omega,\omega^2,1), & (\omega^2,\omega,1); \\ (\omega,1,1), & (\omega^2,\omega^2,1), & (1,\omega,1); \\ (\omega^2,1,1), & (1,\omega^2,1), & (\omega,\omega,1). \end{array}$$

The twelve planes of symmetry of 3{3}3{3}3 play the same role for 2{4}3{3}3 and 3{3}3{4}2; but these two reciprocal polyhedra have nine additional planes of symmetry:

$$u_\mu - \omega^\lambda u_\nu = 0 \quad (\mu \neq \nu).$$

The section of 3{3}3{4}2 by any plane of symmetry is either 3{4}3 or 2{4}6.

Now, we need some general results about finite complex reflection groups. In 1954, Shephard and Todd ([ST]) published a list of all finite irreducible complex reflection groups up to conjugacy. In their classification they separately studied the imprimitive groups and the primitive groups. In the latter case they used the classification of finite collineation groups containing homologies. Furthermore, Shephard and Todd determined the degrees of the reflection groups, using the invariant theory of the corresponding collineation groups in the primitive case. In 1967, Coxeter ([Co3]) presented a number of graphs connected with complex reflection groups to systematize the results of Shephard and Todd. In 1976, Cohen ([C]) gave another approach to obtain a systematization of the same results.

According to [ST], let $\mathfrak{G}$ be a finite unitary group generated by reflections in a unitary space U_n of n dimensions. The matrices which correspond to the operations of $\mathfrak{G}$ can be regarded as collineation matrices in projective space S_{n-1} of $n-1$ dimensions, and the corresponding collineations form a group $\mathfrak{G}'$ which is isomorphic to the quotient group of $\mathfrak{G}$ by the cyclic subgroup $\mathfrak{Z}$ which consists of the elements of $\mathfrak{G}$ represented by scalar matrices. To the m-fold reflections of $\mathfrak{G}$ correspond collineations of finite period m leaving fixed all points of a prime of S_{n-1}. Such collineations are known as homologies, and thus $\mathfrak{G}'$ is generated by homologies.

Now we consider the primitive finite unitary groups generated by reflections in U_3. The groups $\mathfrak{G}'$ generated by homologies are of orders 60, 168, 216 or 360 (see also [MBD]).

$\mathfrak{G}'_{60}$ is isomorphic with the alternating group on five symbols, and the corresponding group $\mathfrak{G}$, of order 120, is $[3,5]$, the symmetry group of the icosahedron.

$\mathfrak{G}'_{216}$ is the Hessian group, which leaves invariant the configuration of inflections of a cubic curve. This group is generated by homologies of period 3, belonging to 12 cyclic subgroups, and in addition contains nine homologies of period 2 which generate an imprimitive subgroup. A finite unitary group generated by reflections corresponding to this collineation group must contain 3-fold reflections, and may, in addition, contain 2-fold reflections corresponding to the homologies of period 2. There are in fact two such groups; one, of order 648, contains only 3-fold reflections and is the symmetry group of the complex regular polyhedron $3\{3\}3\{3\}3$ and the other, of order 1296, containing both 3-fold and 2-fold reflections, is the symmetry group of the regular complex polyhedron $2\{4\}3\{3\}3$ or its reciprocal $3\{3\}3\{4\}2$.

When $n = 4$, the groups $\mathfrak{G}'$ generated by homologies are of orders 576, 1920, 7200, 11520 and 25920.

$\mathfrak{G}'_{25920}$ is unique among the quaternary groups in that all its homologies are of period three, and it gives rise to a finite unitary group generated by reflections of order 155520, the symmetry group of the complex regular polytope $3\{3\}3\{3\}3\{3\}3$.

The primitive groups generated by homologies in more than four variables consist of five cases (see [ST]), which fall into two sets. The first set comprises groups in S_5, S_6 and S_7 respectively of orders $72 \cdot 6!$, $4 \cdot 9!$ and $96 \cdot 10!$ to which correspond Euclidean groups of orders $72 \cdot 6!$, $8 \cdot 9!$ and $192 \cdot 10!$ which are the symmetry groups of the polytopes 2_{21}, 3_{21} and 4_{21} respectively. The remaining two groups, of orders $36 \cdot 6!$ and $18 \cdot 9!$ in S_4 and S_5 correspond to finite unitary groups generated by reflections (in U_5 and U_6 respectively) of orders $72 \cdot 6!$ and $108 \cdot 9!$, they are the groups $[21;2]^3$ and $[21;3]^3$.

Following [ST], the symmetry group of the polyhedron $3\{3\}3\{3\}3$ contains 3-fold reflections in the 12 planes:

$$x_i = 0, \quad x_1 + \omega^j x_2 + \omega^k x_3 = 0 \quad (i,j,k = 1,2,3).$$

It is generated by the reflections

$$\mathbf{R}_1 : \begin{pmatrix} 1 & & \\ & 1 & \\ & & \omega^2 \end{pmatrix}, \quad \mathbf{R}_2 : \frac{1}{\sqrt{-3}} \begin{pmatrix} \omega & \omega^2 & \omega^2 \\ \omega^2 & \omega & \omega^2 \\ \omega^2 & \omega^2 & \omega \end{pmatrix}, \quad \mathbf{R}_3 : \begin{pmatrix} 1 & & \\ & \omega^2 & \\ & & 1 \end{pmatrix}$$

in the planes $x_3 = 0$, $x_1 + x_2 + x_3 = 0$, $x_2 = 0$ respectively. The characteristic roots of $\mathbf{R}_1$ and $\mathbf{R}_3$ are $(1,1,\omega^2)$ and of $\mathbf{R}_2$ are $(1,1,\omega)$. The matrix of the product $R_1 R_2^{-1} R_3$ is

$$\frac{-1}{\sqrt{-3}} \begin{pmatrix} \omega^2 & 1 & \omega \\ \omega & \omega & \omega \\ 1 & \omega^2 & \omega \end{pmatrix}.$$

Its characteristic equation is

$$\lambda^3 - \omega^2\lambda^2 + \omega\lambda - 1 = 0,$$

and its characteristic roots are ζ^5, ζ^8, ζ^{11} where $\zeta = \exp(2\pi i/12)$.

The symmetry group of the regular polyhedron 2{4}3{3}3 or its reciprocal contains in addition to the above 3-fold reflections, 2-fold reflections in the nine planes

$$x_i - \omega^k x_j = 0, \quad (i, j, k = 1, 2, 3).$$

It is generated by the 3-fold reflections $\mathbf{R}_1$, $\mathbf{R}_2$ above together with the 2-fold reflection

$$\mathbf{S} : \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}$$

in the plane $x_2 - x_3 = 0$. The matrix of the product $SR_1R_2^{-1}$ is

$$\frac{-1}{\sqrt{-3}} \begin{pmatrix} \omega^2 & \omega & \omega \\ 1 & 1 & \omega \\ \omega & \omega^2 & \omega \end{pmatrix}.$$

Its characteristic equation is

$$\lambda^3 + \omega = 0,$$

and its characteristic roots are ζ^5, ζ^{11}, ζ^{17} where $\zeta = \exp(2\pi i/18)$.

According to [Hu], when we study the 27 lines on a general cubic surface, all the geometry is closely tied up with the automorphism group of the lines, the Weyl group of E_6. The group $W(E_6)$ is particularly interesting because it can be expressed in terms of different Chevalley groups (see [CC] for more details):

$$G_{25920} \cong PU(3, 1; \mathbb{F}_4) \cong PSp(4, \mathbb{Z}/3\mathbb{Z}),$$

where $G_{25920} \subset W(E_6)$ is the simple subgroup of index two consisting of all even elements.

It is well-known that the group of permutations (see [Hu] or [Ma]), $\mathrm{Aut}(\mathcal{L})$, of the 27 lines (or of the 45 planes), by which we mean the permutations of the lines preserving the intersection behavior of the lines,

$$|\mathrm{Aut}(\mathcal{L})| = |S_6| \cdot 2 \cdot 36 = 51840.$$

In fact, $\mathrm{Aut}(\mathcal{L})$ is nothing but the Weyl group $W(E_6)$, and the 36 double sixes correspond to the positive roots. The 27 lines correspond to the 27 fundamental weights of

E_6, and the many other sets of objects (lines, tritangents, etc.) correspond to natural sets of objects (roots, weights, etc.) of E_6.

Let us recall some fundamental facts about the unitary reflection groups of order 25920 (see [Hu]). We have actions of G_{25920} on $\mathbb{P}^3$ and on $\mathbb{P}^4$, both of which are in fact generated by unitary reflections. We will be interested in the arrangement in $\mathbb{P}^3$. The arrangement induced in each of the 40 planes of this arrangement is the extended Hessian pencil, the arrangement of 21 lines which are the 12 lines of the Hessian pencil together with the nine lines joining corners of the four triangles. The 12 lines of the Hessian pencil are the 12 two-fold lines lying in the plane, the nine other lines are the four-fold lines lying in the plane. We also remark that the five-fold points of the arrangement split into two different types of singular points in the planes, namely 36 two-fold points and nine four-fold points which are the base points of the Hessian pencil. The 12-fold points of the arrangement lying in one of the planes are five-fold points of the induced arrangement. For each 12-fold point we can also speak of the induced arrangement, by blowing up the point in $\mathbb{P}^3$ and considering the proper transforms of the 12 planes intersecting the point in the exceptional $\mathbb{P}^2$. In our case, we get the Hessian pencil itself, i.e., the 12 lines of the four degenerate cubics.

The invariants forms under the action of G_{25920} on $\mathbb{P}^3$ were calculated by Maschke (see [Mas2] and [Hu]). This is done essentially by reducing the problem to that of the invariants of the Hessian group of order 648 acting on $\mathbb{P}^2$. The 40 planes of the arrangement in $\mathbb{P}^3$ are given explicitly in homogeneous coordinates $(z_0 : z_1 : z_2 : z_3)$:

$$\begin{aligned}
&(4) \quad z_i = 0 \quad (i = 0, 1, 2, 3).\\
&(9) \quad (z_1^3 + z_2^3 + z_3^3)^3 - 27 z_1^3 z_2^3 z_3^3 = 0.\\
&(9) \quad (z_0^3 + z_1^3 + z_3^3)^3 - 27 z_0^3 z_1^3 z_3^3 = 0.\\
&(9) \quad (z_0^3 + z_1^3 + z_2^3)^3 - 27 z_0^3 z_1^3 z_2^3 = 0.\\
&(9) \quad (z_0^3 + z_2^3 + z_3^3)^3 - 27 z_0^3 z_2^3 z_3^3 = 0.
\end{aligned}$$

In the $\mathbb{P}^2$ given by $z_0 = 0$ with homogeneous coordinates (z_1, z_2, z_3) the action of G_{648} is generated by five collineations:

$$\tag{4.1} \begin{cases}
A(z_1, z_2, z_3) = (z_2, z_3, z_1),\\
B(z_1, z_2, z_3) = (z_1, z_3, z_2),\\
C(z_1, z_2, z_3) = (z_1, \omega z_2, \omega^2 z_3),\\
D(z_1, z_2, z_3) = (z_1, \omega z_2, \omega z_3),\\
E(z_1, z_2, z_3) = \left(\dfrac{z_1 + z_2 + z_3}{\sqrt{-3}}, \dfrac{z_1 + \omega z_2 + \omega^2 z_3}{\sqrt{-3}}, \dfrac{z_1 + \omega^2 z_2 + \omega z_3}{\sqrt{-3}} \right).
\end{cases}$$

Writing in the form of matrices:

$$A = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}, \quad B = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix},$$

$$C = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \omega & 0 \\ 0 & 0 & \omega^2 \end{pmatrix}, \quad D = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \omega & 0 \\ 0 & 0 & \omega \end{pmatrix},$$

$$E = \frac{1}{\sqrt{-3}} \begin{pmatrix} 1 & 1 & 1 \\ 1 & \omega & \omega^2 \\ 1 & \omega^2 & \omega \end{pmatrix}.$$

Note that

$$AA^* = BB^* = CC^* = DD^* = EE^* = I.$$

This implies that

$$A, B, C, D, E \in U(3).$$

In fact, the Hessian group $G_{648} = \langle A, B, C, D, E \rangle$, where

$$A^3 = B^2 = C^3 = D^3 = E^4 = I.$$

A complete system of invariants for G_{648} has degrees 6, 9, 12, 12 and 18 and can be given explicitly by the following forms:

$$(4.2) \qquad \begin{cases} C_6 = z_1^6 + z_2^6 + z_3^6 - 10(z_1^3 z_2^3 + z_2^3 z_3^3 + z_3^3 z_1^3), \\ C_9 = (z_1^3 - z_2^3)(z_2^3 - z_3^3)(z_3^3 - z_1^3), \\ C_{12} = (z_1^3 + z_2^3 + z_3^3)[(z_1^3 + z_2^3 + z_3^3)^3 + 216 z_1^3 z_2^3 z_3^3], \\ \mathfrak{C}_{12} = z_1 z_2 z_3 [27 z_1^3 z_2^3 z_3^3 - (z_1^3 + z_2^3 + z_3^3)^3], \\ C_{18} = (z_1^3 + z_2^3 + z_3^3)^6 - 540 z_1^3 z_2^3 z_3^3 (z_1^3 + z_2^3 + z_3^3)^3 - 5832 z_1^6 z_2^6 z_3^6. \end{cases}$$

C_{18} is the functional determinant of C_{12} and $\mathfrak{C}_{12}$. $\mathfrak{C}_{12}$ is just the product of the 12 lines of the Hessian arrangement. C_{12} is the Hessian of $\mathfrak{C}_{12}$ and vise versa. C_9 is the so-called difference product of z_1^3, z_2^3, z_3^3. The two relations among these invariants are

$$(4.3) \qquad \begin{cases} 432 C_9^2 = C_6^3 - 3 C_6 C_{12} + 2 C_{18}, \\ 1728 \mathfrak{C}_{12}^3 = C_{18}^2 - C_{12}^3. \end{cases}$$

Following [ST], the Hessian group of order 216 in S_2 is the collineation group that leaves invariant the nine inflections of a pencil of cubic curves in the plane. The invariant forms of this collineation group consist of a sextic C_6, a form C_9 of degree 9 representing the harmonic polars of the inflections, and two forms $\mathfrak{C}_{12}$, C_{12} representing respectively the four degenerate cubics and the four equianharmonic cubics in the pencil. A syzygy connects $\mathfrak{C}_{12}^3$ with C_6, C_9 and C_{12}:

$$6912\mathfrak{C}_{12}^3 = (432C_9^2 - C_6^3 + 3C_6C_{12})^2 - 4C_{12}^3. \tag{4.4}$$

For the two corresponding finite unitary groups generated by reflections in U_3 of order 648 and 1296 respectively, the invariant forms may be taken to be C_6, C_9, C_{12} and C_6, C_{12}, C_{18}, respectively. The Jacobian of C_6, C_9, C_{12} is a multiple of the square of $\mathfrak{C}_{12}$; that of C_6, C_{12}, C_{18} is a multiple of the product $C_9\mathfrak{C}_{12}^2$.

Maschke proved that the action of G_{25920} in $\mathbb{P}^3$ is generated by the action of a G_{648} acting on one of the 40 planes and a tetrahedral group of order 24 consisting of the permutation group acting on the z_i. This tetrahedral group stabilizes the tetrahedron consisting of the first four forms in the 40 planes, and G_{25290} is generated by it and the G_{648} acting on $z_0 = 0$ as above. From this one deduces that each invariant form of G_{25920} can be written as a polynomial in z_0 with coefficients which are polynomial expressions in the invariants of G_{648}.

A complete system of invariants for G_{25920} has degrees 12, 18, 24, 30 and 40 and can be given by the following forms:

$$\left\{\begin{aligned}
F_{12} &= 6z_0^{12} + 6\cdot 22C_6z_0^6 + 6\cdot 220C_9z_0^3 + C_6^2 + 5C_{12},\\
F_{18} &= -54z_0^{18} + 17\cdot 54C_6z_0^{12} + 54\cdot 1870C_9z_0^9\\
&\quad + \frac{1}{2}\cdot 17\cdot 27(19C_6^2 - 15C_{12})z_0^6 + 54\cdot 170C_6C_9z_0^3 + C_6^3\\
&\quad - 30C_6C_{12} - 25C_{18},\\
F_{24} &= 1728C_6z_0^{18} - 36\cdot 1728C_9z_0^{15} + 15\cdot 144(7C_{12} + C_6^2)z_0^{12}\\
&\quad - 10\cdot 1728C_6C_9z_0^9 + 36(178C_{18} - 135C_6C_{12} + 5C_6^3)z_0^6\\
&\quad + 432(41C_{12} - C_6^2)C_9z_0^3 + C_6^4 + 6C_6^2C_{12} - 16C_6C_{18} + 9C_{12}^2,\\
F_{30} &= -2\cdot 6^4C_6z_0^{24} + 312\cdot 6^4C_9z_0^{21} + 216(715C_{12} - 127C_6^2)z_0^{18}\\
&\quad + 272\cdot 6^4C_6C_9z_0^{15} + 18(1306C_{18} + 6045C_6C_{12} - 295C_6^3)z_0^{12}\\
&\quad + 216(73C_6^2 - 5473C_{12})C_9z_0^9\\
&\quad + \frac{1}{2}\cdot 3(16648C_6C_{18} + 2334C_6^2C_{12} - 20709C_{12}^2 - C_6^4)z_0^6\\
&\quad - 36(1370C_{18} - 657C_6C_{12} + 7C_6^3)C_9z_0^3 + C_6^5 - 19C_6^3C_{12}\\
&\quad + 29C_6^2C_{18} - 6C_6C_{12}^2 - 5C_{12}C_{18}.
\end{aligned}\right.$$

F_{40} is just the product of the 40 planes defining the arrangement in $\mathbb{P}^3$, i.e., it represents the primes of the 40 homologies of period three contained in the group.

The forms F_{12}, F_{18}, F_{24}, F_{30} and F_{40} satisfy the relation:

$$\begin{aligned} &2^{28}\cdot 3^{15}\cdot 5^{15}\cdot F_{40}^3\\ &=\begin{vmatrix} \Phi_{30} & 2\Phi_{24}^2 & F_{18}\Phi_{24} & F_{12}\Phi_{24}\\ 2\Phi_{24} & 27F_{12}\Phi_{30}-11F_{18}\Phi_{24} & 3F_{12}\Phi_{24}-4F_{18}^2 & 3\Phi_{30}-4F_{12}F_{18}\\ F_{18} & 3F_{12}\Phi_{24}-4F_{18}^2 & 13F_{12}F_{18}-3\Phi_{30} & 9F_{12}^2-2\Phi_{24}\\ F_{12} & 3\Phi_{30}-4F_{12}F_{18} & 9F_{12}^2-2\Phi_{24} & F_{18} \end{vmatrix}, \end{aligned}$$

where

$$4\Phi_{24}=25F_{24}-9F_{12}^2,$$

$$6\Phi_{30}=25F_{30}-F_{12}F_{18}.$$

There are two simpler relations involving F_{40}. Firstly, F_{40} is the Hessian determinant of F_{12}:

$$\text{Hessian}(F_{12})=2^{14}\cdot 3^8\cdot 5^3\cdot 11^4\cdot F_{40}.$$

Secondly, the Jacobian determinant

$$\frac{\partial(F_{12},F_{18},F_{24},F_{30})}{\partial(z_0,z_1,z_2,z_3)}=2^{26}\cdot 3^{15}\cdot 5^6\cdot F_{40}^2.$$

The action of G_{25920} on $\mathbb{P}^3$ in terms of the hyperelliptic functions $Z_{\alpha,\beta}$ was first treated by Witting (see [Wi]). That is why the arrangement defined by the action of G_{25920} on $\mathbb{P}^3$ is called a Witting configuration.

In [Y], we gave the modular equation associated to the Picard curves:

THEOREM (see [Y], Main Theorem 3). *The following identity of differential forms holds*:

$$\frac{dw_1\wedge dw_2}{\sqrt[3]{w_1^2w_2^2(1-w_1)(1-\lambda_1^3w_1)(1-\lambda_2^3w_1)}}=-\frac{5t_1}{t_2^2}\frac{dt_1\wedge dt_2}{\sqrt[3]{t_1^2t_2^2(1-t_1)(1-\kappa_1^3t_1)(1-\kappa_2^3t_1)}},$$

where

$$w_1=\frac{(\beta+\gamma)t_1+\alpha}{\beta t_1+(\alpha+\gamma)},\quad w_2=\frac{t_1[(\beta+\gamma)t_1+\alpha]^2[\beta t_1+(\alpha+\gamma)]}{t_2^5},$$

which is a rational transformation of order five. Here,

$$\begin{cases} \alpha=\dfrac{(v_1-1)(v_2-1)(v_1+v_2-2)-(u_1-1)(u_2-1)(u_1+u_2-2)}{2(u_1-1)(u_2-1)(v_1-1)(v_2-1)},\\ \beta=\dfrac{-(v_1-1)(v_2-1)(2v_1v_2-v_1-v_2)+(u_1-1)(u_2-1)(2u_1u_2-u_1-u_2)}{2(u_1-1)(u_2-1)(v_1-1)(v_2-1)},\\ \gamma=\dfrac{(v_1-1)(v_2-1)}{(u_1-1)(u_2-1)}, \end{cases}$$

with

$$u_1 = \kappa_1^3, \quad u_2 = \kappa_2^3, \quad v_1 = \lambda_1^3, \quad v_2 = \lambda_2^3.$$

Moreover, the moduli (κ_1, κ_2) *and* (λ_1, λ_2) *satisfy the modular equation*:

$$(\kappa_1^3 - 1)(\kappa_2^3 - 1)(\kappa_1^3 - \kappa_2^3) = (\lambda_1^3 - 1)(\lambda_2^3 - 1)(\lambda_1^3 - \lambda_2^3),$$

which is an algebraic variety of dimension three. The corresponding two algebraic surfaces are:

$$\begin{cases} w_3^3 = w_1^2 w_2^2 (1 - w_1)(1 - \lambda_1^3 w_1)(1 - \lambda_2^3 w_1), \\ t_3^3 = t_1^2 t_2^2 (1 - t_1)(1 - \kappa_1^3 t_1)(1 - \kappa_2^3 t_1), \end{cases}$$

which give the homogeneous algebraic equations of degree seven:

$$\begin{cases} w_3^3 w_4^4 = w_1^2 w_2^2 (w_4 - w_1)(w_4 - \lambda_1^3 w_1)(w_4 - \lambda_2^3 w_1), \\ t_3^3 t_4^4 = t_1^2 t_2^2 (t_4 - t_1)(t_4 - \kappa_1^3 t_1)(t_4 - \kappa_2^3 t_1). \end{cases}$$

In fact, the above modular equation can be written as follows:

$$C_9(\kappa_1, \kappa_2, 1) = C_9(\lambda_1, \lambda_2, 1). \tag{4.5}$$

This shows that our modular equation is intimately connected with the Hessian polyhedra.

Set $X = z_1^3$, $Y = z_2^3$ and $Z = z_3^3$. Put

$$\begin{cases} \sigma_1 = X + Y + Z, \\ \sigma_2 = XY + YZ + ZX, \\ \sigma_3 = XYZ. \end{cases} \tag{4.6}$$

Then

$$\begin{cases} C_6 = \sigma_1^2 - 12\sigma_2, \\ C_{12} = \sigma_1(\sigma_1^3 + 216\sigma_3), \\ C_{18} = \sigma_1^6 - 540\sigma_1^3\sigma_3 - 5832\sigma_3^2, \\ \mathfrak{C}_{12}^3 = \sigma_3(27\sigma_3 - \sigma_1^3)^3. \end{cases} \tag{4.7}$$

Note that

$$C_9 = (X - Y)(Y - Z)(Z - X) = (XY^2 + YZ^2 + ZX^2) - (X^2Y + Y^2Z + Z^2X),$$

we have

$$\begin{aligned}C_9^2 = &(XY^2+YZ^2+ZX^2+X^2Y+Y^2Z+Z^2X)^2+\\ &-4(XY^2+YZ^2+ZX^2)(X^2Y+Y^2Z+Z^2X).\end{aligned}$$

Here,

$$\begin{aligned}&XY^2+YZ^2+ZX^2+X^2Y+Y^2Z+Z^2X\\ =&(X+Y+Z)(XY+YZ+ZX)-3XYZ\\ =&\sigma_1\sigma_2-3\sigma_3.\end{aligned}$$

$$\begin{aligned}&(XY^2+YZ^2+ZX^2)(X^2Y+Y^2Z+Z^2X)\\ =&X^3Y^3+Y^3Z^3+Z^3X^3+XYZ(X^3+Y^3+Z^3)+3X^2Y^2Z^2,\end{aligned}$$

where

$$\begin{aligned}&X^3Y^3+Y^3Z^3+Z^3X^3\\ =&(XY+YZ+ZX)[(XY+YZ+ZX)^2-3XYZ(X+Y+Z)]+3(XYZ)^2\\ =&\sigma_2(\sigma_2^2-3\sigma_1\sigma_3)+3\sigma_3^2,\end{aligned}$$

$$\begin{aligned}&X^3+Y^3+Z^3\\ =&(X+Y+Z)[(X+Y+Z)^2-3(XY+YZ+ZX)]+3XYZ\\ =&\sigma_1(\sigma_1^2-3\sigma_2)+3\sigma_3.\end{aligned}$$

So,

$$(XY^2+YZ^2+ZX^2)(X^2Y+Y^2Z+Z^2X)=\sigma_2^3-6\sigma_1\sigma_2\sigma_3+9\sigma_3^2+\sigma_1^3\sigma_3.$$

Thus,

$$C_9^2=\sigma_1^2\sigma_2^2-4\sigma_1^3\sigma_3-4\sigma_2^3-27\sigma_3^2+18\sigma_1\sigma_2\sigma_3. \tag{4.8}$$

In the affine coordinates $\xi = z_1/z_3$ and $\eta = z_2/z_3$, we have

$$\begin{cases} C_6 = C_6(\xi,\eta) = \xi^6+\eta^6+1-10(\xi^3\eta^3+\xi^3+\eta^3),\\ C_9 = C_9(\xi,\eta) = (\xi^3-\eta^3)(\eta^3-1)(1-\xi^3),\\ C_{12} = C_{12}(\xi,\eta) = (\xi^3+\eta^3+1)[(\xi^3+\eta^3+1)^3+216\xi^3\eta^3],\\ \mathfrak{C}_{12} = \mathfrak{C}_{12}(\xi,\eta) = \xi\eta[27\xi^3\eta^3-(\xi^3+\eta^3+1)^3],\\ C_{18} = C_{18}(\xi,\eta) = (\xi^3+\eta^3+1)^6-540\xi^3\eta^3(\xi^3+\eta^3+1)^3-5832\xi^6\eta^6.\end{cases} \tag{4.9}$$

The relation

$$432C_9^2 = C_6^3-3C_6C_{12}+2C_{18}$$

gives that

$$432\frac{C_9^2}{C_6^3}=1-3\frac{C_{12}}{C_6^2}+2\frac{C_{18}}{C_6^3}. \tag{4.10}$$

We define the Hessian polyhedral equations as follows:

$$\begin{cases} 432\dfrac{C_9^2}{C_6^3}=R_1(x,y), \\ 3\dfrac{C_{12}}{C_6^2}=R_2(x,y), \end{cases} \tag{4.11}$$

where R_1 and R_2 are rational functions of x and y.

Theorem 4.1 (Main Theorem 2). *The relations between Hessian polyhedra and Appell hypergeometric partial differential equations are given by the following identities*:

$$\begin{aligned} &z_1^6+z_2^6+z_3^6-10(z_1^3z_2^3+z_2^3z_3^3+z_3^3z_1^3) \\ =&2^6\cdot 3^7\cdot C^2\cdot R_1\left[\frac{1}{4}(R_1+R_2-1)^2-\frac{1}{27}R_2^3\right]^{\frac{4}{3}}\left[\frac{\partial(R_1,R_2)}{\partial(x,y)}\right]^{-2}H(x,y)^2. \end{aligned} \tag{4.12}$$

$$\begin{aligned} &(z_1^3-z_2^3)(z_2^3-z_3^3)(z_3^3-z_1^3) \\ =&2^7\cdot 3^9\cdot C^3\cdot R_1^2\left[\frac{1}{4}(R_1+R_2-1)^2-\frac{1}{27}R_2^3\right]^2\left[\frac{\partial(R_1,R_2)}{\partial(x,y)}\right]^{-3}H(x,y)^3. \end{aligned} \tag{4.13}$$

$$\begin{aligned} &(z_1^3+z_2^3+z_3^3)[(z_1^3+z_2^3+z_3^3)^3+216z_1^3z_2^3z_3^3] \\ =&2^{12}\cdot 3^{13}\cdot C^4\cdot R_1^2R_2\left[\frac{1}{4}(R_1+R_2-1)^2-\frac{1}{27}R_2^3\right]^{\frac{8}{3}}\left[\frac{\partial(R_1,R_2)}{\partial(x,y)}\right]^{-4}H(x,y)^4. \end{aligned} \tag{4.14}$$

$$\begin{aligned} &z_1z_2z_3[27z_1^3z_2^3z_3^3-(z_1^3+z_2^3+z_3^3)^3] \\ =&2^{10}\cdot 3^{13}\cdot C^4\cdot R_1^2\left[\frac{1}{4}(R_1+R_2-1)^2-\frac{1}{27}R_2^3\right]^3\left[\frac{\partial(R_1,R_2)}{\partial(x,y)}\right]^{-4}H(x,y)^4. \end{aligned} \tag{4.15}$$

$$\begin{aligned} &(z_1^3+z_2^3+z_3^3)^6-540z_1^3z_2^3z_3^3(z_1^3+z_2^3+z_3^3)^3-5832z_1^6z_2^6z_3^6 \\ =&2^{17}\cdot 3^{21}\cdot C^6\cdot R_1^3(R_1+R_2-1)\left[\frac{1}{4}(R_1+R_2-1)^2-\frac{1}{27}R_2^3\right]^4 \\ &\times\left[\frac{\partial(R_1,R_2)}{\partial(x,y)}\right]^{-6}H(x,y)^6. \end{aligned} \tag{4.16}$$

Here,

$$H(x,y) := x^{b'-c}y^{b-c}(x-1)^{c-a-b-1}(y-1)^{c-a-b'-1}(x-y)^{-b-b'}. \tag{4.17}$$

Proof. We have

$$\frac{\partial(R_1,R_2)}{\partial(\xi,\eta)} = 1296\frac{\partial\left(\frac{C_9^2}{C_6^3},\frac{C_{12}}{C_6^2}\right)}{\partial(\xi,\eta)}.$$

Here, we find that

$$\frac{\partial\left(\frac{C_9^2}{C_6^3},\frac{C_{12}}{C_6^2}\right)}{\partial(\xi,\eta)} = \frac{C_9}{C_6^6}\left[2C_6\frac{\partial(C_9,C_{12})}{\partial(\xi,\eta)} + 3C_9\frac{\partial(C_{12},C_6)}{\partial(\xi,\eta)} + 4C_{12}\frac{\partial(C_6,C_9)}{\partial(\xi,\eta)}\right].$$

Now, we need the following proposition:

Proposition 4.2 (Main Proposition 1). *The following identities hold*:

$$\begin{vmatrix} \frac{\partial C_6}{\partial\xi} & \frac{\partial C_6}{\partial\eta} & 2C_6 \\ \frac{\partial C_9}{\partial\xi} & \frac{\partial C_9}{\partial\eta} & 3C_9 \\ \frac{\partial C_{12}}{\partial\xi} & \frac{\partial C_{12}}{\partial\eta} & 4C_{12} \end{vmatrix} = -864\mathfrak{C}_{12}^2. \tag{4.18}$$

$$\begin{vmatrix} \frac{\partial C_6}{\partial\xi} & \frac{\partial C_6}{\partial\eta} & 2C_6 \\ \frac{\partial C_9}{\partial\xi} & \frac{\partial C_9}{\partial\eta} & 3C_9 \\ \frac{\partial C_{18}}{\partial\xi} & \frac{\partial C_{18}}{\partial\eta} & 6C_{18} \end{vmatrix} = -1296C_6\mathfrak{C}_{12}^2. \tag{4.19}$$

$$\begin{vmatrix} \frac{\partial C_6}{\partial\xi} & \frac{\partial C_6}{\partial\eta} & 2C_6 \\ \frac{\partial C_{12}}{\partial\xi} & \frac{\partial C_{12}}{\partial\eta} & 4C_{12} \\ \frac{\partial C_{18}}{\partial\xi} & \frac{\partial C_{18}}{\partial\eta} & 6C_{18} \end{vmatrix} = 432\cdot 864C_9\mathfrak{C}_{12}^2. \tag{4.20}$$

$$\begin{vmatrix} \frac{\partial C_9}{\partial\xi} & \frac{\partial C_9}{\partial\eta} & 3C_9 \\ \frac{\partial C_{12}}{\partial\xi} & \frac{\partial C_{12}}{\partial\eta} & 4C_{12} \\ \frac{\partial C_{18}}{\partial\xi} & \frac{\partial C_{18}}{\partial\eta} & 6C_{18} \end{vmatrix} = 1296(C_6^2 - C_{12})\mathfrak{C}_{12}^2. \tag{4.21}$$

$$\begin{vmatrix} \frac{\partial C_6}{\partial\xi} & \frac{\partial C_6}{\partial\eta} & 2C_6 \\ \frac{\partial C_9}{\partial\xi} & \frac{\partial C_9}{\partial\eta} & 3C_9 \\ \frac{\partial \mathfrak{C}_{12}}{\partial\xi} & \frac{\partial \mathfrak{C}_{12}}{\partial\eta} & 4\mathfrak{C}_{12} \end{vmatrix} = \frac{1}{2}(C_{12}^2 - C_6C_{18}). \tag{4.22}$$

$$\begin{vmatrix} \frac{\partial C_6}{\partial \xi} & \frac{\partial C_6}{\partial \eta} & 2C_6 \\ \frac{\partial C_{12}}{\partial \xi} & \frac{\partial C_{12}}{\partial \eta} & 4C_{12} \\ \frac{\partial \mathfrak{C}_{12}}{\partial \xi} & \frac{\partial \mathfrak{C}_{12}}{\partial \eta} & 4\mathfrak{C}_{12} \end{vmatrix} = 144 C_9 C_{18}. \tag{4.23}$$

$$\begin{vmatrix} \frac{\partial C_9}{\partial \xi} & \frac{\partial C_9}{\partial \eta} & 3C_9 \\ \frac{\partial C_{12}}{\partial \xi} & \frac{\partial C_{12}}{\partial \eta} & 4C_{12} \\ \frac{\partial \mathfrak{C}_{12}}{\partial \xi} & \frac{\partial \mathfrak{C}_{12}}{\partial \eta} & 4\mathfrak{C}_{12} \end{vmatrix} = \frac{1}{2}(C_6^2 - C_{12}) C_{18}. \tag{4.24}$$

$$\begin{vmatrix} \frac{\partial C_6}{\partial \xi} & \frac{\partial C_6}{\partial \eta} & 2C_6 \\ \frac{\partial C_{18}}{\partial \xi} & \frac{\partial C_{18}}{\partial \eta} & 6C_{18} \\ \frac{\partial \mathfrak{C}_{12}}{\partial \xi} & \frac{\partial \mathfrak{C}_{12}}{\partial \eta} & 4\mathfrak{C}_{12} \end{vmatrix} = 216 C_9 C_{12}^2. \tag{4.25}$$

$$\begin{vmatrix} \frac{\partial C_9}{\partial \xi} & \frac{\partial C_9}{\partial \eta} & 3C_9 \\ \frac{\partial C_{18}}{\partial \xi} & \frac{\partial C_{18}}{\partial \eta} & 6C_{18} \\ \frac{\partial \mathfrak{C}_{12}}{\partial \xi} & \frac{\partial \mathfrak{C}_{12}}{\partial \eta} & 4\mathfrak{C}_{12} \end{vmatrix} = \frac{3}{4}(C_6^2 - C_{12}) C_{12}^2. \tag{4.26}$$

$$\begin{vmatrix} \frac{\partial C_{12}}{\partial \xi} & \frac{\partial C_{12}}{\partial \eta} & 4C_{12} \\ \frac{\partial C_{18}}{\partial \xi} & \frac{\partial C_{18}}{\partial \eta} & 6C_{18} \\ \frac{\partial \mathfrak{C}_{12}}{\partial \xi} & \frac{\partial \mathfrak{C}_{12}}{\partial \eta} & 4\mathfrak{C}_{12} \end{vmatrix} = 0. \tag{4.27}$$

Proof. Put $X = \xi^3$ and $Y = \eta^3$. We have

$$\frac{\partial C_6}{\partial \xi} = 6\xi^2 (X - 5Y - 5),$$

$$\frac{\partial C_6}{\partial \eta} = 6\eta^2 (Y - 5X - 5),$$

$$\frac{\partial C_9}{\partial \xi} = 3\xi^2 (Y - 1)(1 - 2X + Y),$$

$$\frac{\partial C_9}{\partial \eta} = 3\eta^2 (1 - X)(1 - 2Y + X),$$

$$\frac{\partial C_{12}}{\partial \xi} = 12\xi^2 [(X + Y + 1)^3 + 54XY + 54Y(X + Y + 1)],$$

$$\frac{\partial C_{12}}{\partial \eta} = 12\eta^2[(X+Y+1)^3 + 54XY + 54X(X+Y+1)].$$

Hence,

$$\frac{\partial(C_6, C_9)}{\partial(\xi, \eta)} = 18\xi^2\eta^2[-(X+Y)^3 + 15(X^2+Y^2) - 24XY + 6(X+Y) - 10].$$

$$\begin{aligned}\frac{\partial(C_9, C_{12})}{\partial(\xi, \eta)} =& 36\xi^2\eta^2\{[(X+Y+1)^3 + 54XY][X^2+Y^2-4XY+2(X+Y)-2]+ \\ &+54(X+Y+1)[-XY(X+Y)+2(X^2+Y^2)-(X+Y)]\}.\end{aligned}$$

$$\frac{\partial(C_{12}, C_6)}{\partial(\xi, \eta)} = 432\xi^2\eta^2(Y-X)[(X+Y+1)^3 + 54XY + 9(X+Y+1)(X+Y-5)].$$

Note that

$$\begin{cases} C_6 = X^2+Y^2+1-10(XY+X+Y), \\ C_9 = (X-Y)(Y-1)(1-X), \\ C_{12} = (X+Y+1)[(X+Y+1)^3 + 216XY]. \end{cases}$$

Put

$$\sigma_1 = X+Y+1, \quad \sigma_2 = XY.$$

Then

$$\begin{cases} C_6 = \sigma_1^2 - 12\sigma_2 - 12\sigma_1 + 12, \\ C_9 = (Y-X)(\sigma_2 - \sigma_1 + 2), \\ C_{12} = \sigma_1(\sigma_1^3 + 216\sigma_2), \end{cases}$$

and

$$\frac{\partial(C_9, C_{12})}{\partial(\xi, \eta)} = 36\xi^2\eta^2(\sigma_1^5 - 6\sigma_1^3\sigma_2 - 324\sigma_2^2 + 105\sigma_1^3 - 162\sigma_1\sigma_2 - 270\sigma_1^2 - 162\sigma_2 + 162\sigma_1),$$

$$\frac{\partial(C_{12}, C_6)}{\partial(\xi, \eta)} = 432\xi^2\eta^2(Y-X)(\sigma_1^3 + 9\sigma_1^2 + 54\sigma_2 - 54\sigma_1),$$

$$\frac{\partial(C_6, C_9)}{\partial(\xi, \eta)} = 18\xi^2\eta^2(-\sigma_1^3 + 18\sigma_1^2 - 54\sigma_2 - 27\sigma_1).$$

Therefore,

$$\begin{aligned}
&2C_6\frac{\partial(C_9,C_{12})}{\partial(\xi,\eta)}+3C_9\frac{\partial(C_{12},C_6)}{\partial(\xi,\eta)}+4C_{12}\frac{\partial(C_6,C_9)}{\partial(\xi,\eta)}\\
=&72\xi^2\eta^2[(\sigma_1^2-12\sigma_2-12\sigma_1+12)\times\\
&\times(\sigma_1^5-6\sigma_1^3\sigma_2-324\sigma_2^2+105\sigma_1^3-162\sigma_1\sigma_2-270\sigma_1^2-162\sigma_2+162\sigma_1)+\\
&+18(\sigma_1^2-4\sigma_2-2\sigma_1+1)(\sigma_2-\sigma_1+2)(\sigma_1^3+9\sigma_1^2+54\sigma_2-54\sigma_1)+\\
&+\sigma_1(\sigma_1^3+216\sigma_2)(-\sigma_1^3+18\sigma_1^2-54\sigma_2-27\sigma_1)]\\
=&72\xi^2\eta^2\cdot(-12)(27\sigma_2-\sigma_1^3)^2\\
=&-864\xi^2\eta^2(27\sigma_2-\sigma_1^3)^2.
\end{aligned}$$

On the other hand, note that

$$\mathfrak{C}_{12}=\xi\eta[27XY-(X+Y+1)^3]=\xi\eta(27\sigma_2-\sigma_1^3).$$

Hence, we find that

$$2C_6\frac{\partial(C_9,C_{12})}{\partial(\xi,\eta)}+3C_9\frac{\partial(C_{12},C_6)}{\partial(\xi,\eta)}+4C_{12}\frac{\partial(C_6,C_9)}{\partial(\xi,\eta)}=-864\mathfrak{C}_{12}^2.$$

By the identity $432C_9^2=C_6^3-3C_6C_{12}+2C_{18}$, we have

$$C_{18}=\frac{1}{2}(C_{12}-C_6^2)C_6+216C_9^2+C_6C_{12}.$$

$$\frac{\partial C_{18}}{\partial\xi}=\frac{3}{2}(C_{12}-C_6^2)\frac{\partial C_6}{\partial\xi}+432C_9\frac{\partial C_9}{\partial\xi}+\frac{3}{2}C_6\frac{\partial C_{12}}{\partial\xi}.$$

$$\frac{\partial C_{18}}{\partial\eta}=\frac{3}{2}(C_{12}-C_6^2)\frac{\partial C_6}{\partial\eta}+432C_9\frac{\partial C_9}{\partial\eta}+\frac{3}{2}C_6\frac{\partial C_{12}}{\partial\eta}.$$

Thus,

$$\begin{vmatrix}\frac{\partial C_6}{\partial\xi}&\frac{\partial C_6}{\partial\eta}&2C_6\\\frac{\partial C_9}{\partial\xi}&\frac{\partial C_9}{\partial\eta}&3C_9\\\frac{\partial C_{18}}{\partial\xi}&\frac{\partial C_{18}}{\partial\eta}&6C_{18}\end{vmatrix}=\frac{3}{2}C_6\begin{vmatrix}\frac{\partial C_6}{\partial\xi}&\frac{\partial C_6}{\partial\eta}&2C_6\\\frac{\partial C_9}{\partial\xi}&\frac{\partial C_9}{\partial\eta}&3C_9\\\frac{\partial C_{12}}{\partial\xi}&\frac{\partial C_{12}}{\partial\eta}&4C_{12}\end{vmatrix}=-1296C_6\mathfrak{C}_{12}^2.$$

$$\begin{vmatrix}\frac{\partial C_6}{\partial\xi}&\frac{\partial C_6}{\partial\eta}&2C_6\\\frac{\partial C_{12}}{\partial\xi}&\frac{\partial C_{12}}{\partial\eta}&4C_{12}\\\frac{\partial C_{18}}{\partial\xi}&\frac{\partial C_{18}}{\partial\eta}&6C_{18}\end{vmatrix}=-432C_9\begin{vmatrix}\frac{\partial C_6}{\partial\xi}&\frac{\partial C_6}{\partial\eta}&2C_6\\\frac{\partial C_9}{\partial\xi}&\frac{\partial C_9}{\partial\eta}&3C_9\\\frac{\partial C_{12}}{\partial\xi}&\frac{\partial C_{12}}{\partial\eta}&4C_{12}\end{vmatrix}=432\cdot 864C_9\mathfrak{C}_{12}^2.$$

$$\begin{vmatrix} \frac{\partial C_9}{\partial\xi} & \frac{\partial C_9}{\partial\eta} & 3C_9 \\ \frac{\partial C_{12}}{\partial\xi} & \frac{\partial C_{12}}{\partial\eta} & 4C_{12} \\ \frac{\partial C_{18}}{\partial\xi} & \frac{\partial C_{18}}{\partial\eta} & 6C_{18} \end{vmatrix} = \frac{3}{2}(C_{12}-C_6^2)\begin{vmatrix} \frac{\partial C_6}{\partial\xi} & \frac{\partial C_6}{\partial\eta} & 2C_6 \\ \frac{\partial C_9}{\partial\xi} & \frac{\partial C_9}{\partial\eta} & 3C_9 \\ \frac{\partial C_{12}}{\partial\xi} & \frac{\partial C_{12}}{\partial\eta} & 4C_{12} \end{vmatrix} = 1296(C_6^2-C_{12})\mathfrak{C}_{12}^2.$$

By the identity $1728\mathfrak{C}_{12}^3 = C_{18}^2 - C_{12}^3$, we have

$$\mathfrak{C}_{12} = \frac{C_{18}^2}{1728\mathfrak{C}_{12}^2} - \frac{C_{12}^3}{1728\mathfrak{C}_{12}^2}.$$

$$\frac{\partial\mathfrak{C}_{12}}{\partial\xi} = \frac{1}{2592}\frac{C_{18}}{\mathfrak{C}_{12}^2}\frac{\partial C_{18}}{\partial\xi} - \frac{1}{1728}\frac{C_{12}^2}{\mathfrak{C}_{12}^2}\frac{\partial C_{12}}{\partial\xi}.$$

$$\frac{\partial\mathfrak{C}_{12}}{\partial\eta} = \frac{1}{2592}\frac{C_{18}}{\mathfrak{C}_{12}^2}\frac{\partial C_{18}}{\partial\eta} - \frac{1}{1728}\frac{C_{12}^2}{\mathfrak{C}_{12}^2}\frac{\partial C_{12}}{\partial\eta}.$$

Thus,

$$\begin{aligned} &\begin{vmatrix} \frac{\partial C_6}{\partial\xi} & \frac{\partial C_6}{\partial\eta} & 2C_6 \\ \frac{\partial C_9}{\partial\xi} & \frac{\partial C_9}{\partial\eta} & 3C_9 \\ \frac{\partial \mathfrak{C}_{12}}{\partial\xi} & \frac{\partial \mathfrak{C}_{12}}{\partial\eta} & 4\mathfrak{C}_{12} \end{vmatrix} \\ =&\frac{1}{2592}\frac{C_{18}}{\mathfrak{C}_{12}^2}\begin{vmatrix} \frac{\partial C_6}{\partial\xi} & \frac{\partial C_6}{\partial\eta} & 2C_6 \\ \frac{\partial C_9}{\partial\xi} & \frac{\partial C_9}{\partial\eta} & 3C_9 \\ \frac{\partial C_{18}}{\partial\xi} & \frac{\partial C_{18}}{\partial\eta} & 6C_{18} \end{vmatrix} - \frac{1}{1728}\frac{C_{12}^2}{\mathfrak{C}_{12}^2}\begin{vmatrix} \frac{\partial C_6}{\partial\xi} & \frac{\partial C_6}{\partial\eta} & 2C_6 \\ \frac{\partial C_9}{\partial\xi} & \frac{\partial C_9}{\partial\eta} & 3C_9 \\ \frac{\partial C_{12}}{\partial\xi} & \frac{\partial C_{12}}{\partial\eta} & 4C_{12} \end{vmatrix} \\ =&\frac{1}{2}(C_{12}^2 - C_6C_{18}). \end{aligned}$$

$$\begin{vmatrix} \frac{\partial C_6}{\partial\xi} & \frac{\partial C_6}{\partial\eta} & 2C_6 \\ \frac{\partial C_{12}}{\partial\xi} & \frac{\partial C_{12}}{\partial\eta} & 4C_{12} \\ \frac{\partial \mathfrak{C}_{12}}{\partial\xi} & \frac{\partial \mathfrak{C}_{12}}{\partial\eta} & 4\mathfrak{C}_{12} \end{vmatrix} = \frac{1}{2592}\frac{C_{18}}{\mathfrak{C}_{12}^2}\begin{vmatrix} \frac{\partial C_6}{\partial\xi} & \frac{\partial C_6}{\partial\eta} & 2C_6 \\ \frac{\partial C_{12}}{\partial\xi} & \frac{\partial C_{12}}{\partial\eta} & 4C_{12} \\ \frac{\partial C_{18}}{\partial\xi} & \frac{\partial C_{18}}{\partial\eta} & 6C_{18} \end{vmatrix} = 144C_9C_{18}.$$

$$\begin{vmatrix} \frac{\partial C_9}{\partial\xi} & \frac{\partial C_9}{\partial\eta} & 3C_9 \\ \frac{\partial C_{12}}{\partial\xi} & \frac{\partial C_{12}}{\partial\eta} & 4C_{12} \\ \frac{\partial \mathfrak{C}_{12}}{\partial\xi} & \frac{\partial \mathfrak{C}_{12}}{\partial\eta} & 4\mathfrak{C}_{12} \end{vmatrix} = \frac{1}{2592}\frac{C_{18}}{\mathfrak{C}_{12}^2}\begin{vmatrix} \frac{\partial C_9}{\partial\xi} & \frac{\partial C_9}{\partial\eta} & 3C_9 \\ \frac{\partial C_{12}}{\partial\xi} & \frac{\partial C_{12}}{\partial\eta} & 4C_{12} \\ \frac{\partial C_{18}}{\partial\xi} & \frac{\partial C_{18}}{\partial\eta} & 6C_{18} \end{vmatrix} = \frac{1}{2}(C_6^2 - C_{12})C_{18}.$$

$$\begin{vmatrix} \frac{\partial C_6}{\partial\xi} & \frac{\partial C_6}{\partial\eta} & 2C_6 \\ \frac{\partial C_{18}}{\partial\xi} & \frac{\partial C_{18}}{\partial\eta} & 6C_{18} \\ \frac{\partial \mathfrak{C}_{12}}{\partial\xi} & \frac{\partial \mathfrak{C}_{12}}{\partial\eta} & 4\mathfrak{C}_{12} \end{vmatrix} = \frac{1}{1728}\frac{C_{12}^2}{\mathfrak{C}_{12}^2}\begin{vmatrix} \frac{\partial C_6}{\partial\xi} & \frac{\partial C_6}{\partial\eta} & 2C_6 \\ \frac{\partial C_{12}}{\partial\xi} & \frac{\partial C_{12}}{\partial\eta} & 4C_{12} \\ \frac{\partial C_{18}}{\partial\xi} & \frac{\partial C_{18}}{\partial\eta} & 6C_{18} \end{vmatrix} = 216C_9C_{12}^2.$$

$$\begin{vmatrix} \frac{\partial C_9}{\partial\xi} & \frac{\partial C_9}{\partial\eta} & 3C_9 \\ \frac{\partial C_{18}}{\partial\xi} & \frac{\partial C_{18}}{\partial\eta} & 6C_{18} \\ \frac{\partial \mathfrak{C}_{12}}{\partial\xi} & \frac{\partial \mathfrak{C}_{12}}{\partial\eta} & 4\mathfrak{C}_{12} \end{vmatrix} = \frac{1}{1728}\frac{C_{12}^2}{\mathfrak{C}_{12}^2}\begin{vmatrix} \frac{\partial C_9}{\partial\xi} & \frac{\partial C_9}{\partial\eta} & 3C_9 \\ \frac{\partial C_{12}}{\partial\xi} & \frac{\partial C_{12}}{\partial\eta} & 4C_{12} \\ \frac{\partial C_{18}}{\partial\xi} & \frac{\partial C_{18}}{\partial\eta} & 6C_{18} \end{vmatrix} = \frac{3}{4}(C_6^2 - C_{12})C_{12}^2.$$

□

Now, we have

$$\frac{\partial(R_1,R_2)}{\partial(\xi,\eta)}=-2^9\cdot 3^7\frac{C_9}{C_6^6}\mathfrak{C}_{12}^2.$$

By (3.15), this implies that

$$\frac{\partial(R_1,R_2)}{\partial(x,y)}=-2^9\cdot 3^7\frac{C_9}{C_6^6}\mathfrak{C}_{12}^2\frac{C}{z_3^3}H(x,y).$$

Hence,

$$-2^9\cdot 3^7\cdot C\frac{C_9\mathfrak{C}_{12}^2}{C_6^6\frac{\partial(R_1,R_2)}{\partial(x,y)}}H(x,y)=z_3^3.$$

By (4.3) and (4.11), we have

$$\begin{cases}\dfrac{C_{18}}{C_6^3}=\dfrac{1}{2}(R_1+R_2-1),\\ \dfrac{\mathfrak{C}_{12}^3}{C_6^6}=\dfrac{1}{1728}\left[\dfrac{1}{4}(R_1+R_2-1)^2-\dfrac{1}{27}R_2^3\right].\end{cases}$$

Note that

$$\sqrt{C_6}\frac{C_9\mathfrak{C}_{12}^2}{C_6^6}=\left(\frac{C_9^2}{C_6^3}\right)^{\frac{1}{2}}\left(\frac{\mathfrak{C}_{12}^3}{C_6^6}\right)^{\frac{2}{3}}=\frac{1}{144}\left(\frac{R_1}{432}\right)^{\frac{1}{2}}\left[\frac{1}{4}(R_1+R_2-1)^2-\frac{1}{27}R_2^3\right]^{\frac{2}{3}}.$$

We have

$$\sqrt{C_6}z_3^3=-2^9\cdot 3^7\cdot C\cdot\frac{1}{144}\left(\frac{R_1}{432}\right)^{\frac{1}{2}}\left[\frac{1}{4}(R_1+R_2-1)^2-\frac{1}{27}R_2^3\right]^{\frac{2}{3}}\left[\frac{\partial(R_1,R_2)}{\partial(x,y)}\right]^{-1}H(x,y).$$

Hence, we find that

$$\begin{aligned}&z_1^6+z_2^6+z_3^6-10(z_1^3z_2^3+z_2^3z_3^3+z_3^3z_1^3)=C_6(\xi,\eta)\cdot z_3^6\\ =&2^6\cdot 3^7\cdot C^2\cdot R_1\left[\frac{1}{4}(R_1+R_2-1)^2-\frac{1}{27}R_2^3\right]^{\frac{4}{3}}\left[\frac{\partial(R_1,R_2)}{\partial(x,y)}\right]^{-2}H(x,y)^2.\end{aligned}$$

By

$$\begin{cases}C_9=\dfrac{1}{\sqrt{432}}\sqrt{R_1}C_6^{\frac{3}{2}},\\ C_{12}=\dfrac{1}{3}R_2C_6^2,\\ C_{18}=\dfrac{1}{2}(R_1+R_2-1)C_6^3,\\ \mathfrak{C}_{12}=\dfrac{1}{12}\sqrt[3]{\dfrac{1}{4}(R_1+R_2-1)^2-\dfrac{1}{27}R_2^3}\cdot C_6^2,\end{cases}$$

we have

$$\begin{aligned}&(z_1^3-z_2^3)(z_2^3-z_3^3)(z_3^3-z_1^3)\\=&2^7\cdot 3^9\cdot C^3\cdot R_1^2\left[\frac{1}{4}(R_1+R_2-1)^2-\frac{1}{27}R_2^3\right]^2\left[\frac{\partial(R_1,R_2)}{\partial(x,y)}\right]^{-3}H(x,y)^3.\end{aligned}$$

$$\begin{aligned}&(z_1^3+z_2^3+z_3^3)[(z_1^3+z_2^3+z_3^3)^3+216z_1^3z_2^3z_3^3]\\=&2^{12}\cdot 3^{13}\cdot C^4\cdot R_1^2R_2\left[\frac{1}{4}(R_1+R_2-1)^2-\frac{1}{27}R_2^3\right]^{\frac{8}{3}}\left[\frac{\partial(R_1,R_2)}{\partial(x,y)}\right]^{-4}H(x,y)^4.\end{aligned}$$

$$\begin{aligned}&z_1z_2z_3[27z_1^3z_2^3z_3^3-(z_1^3+z_2^3+z_3^3)^3]\\=&2^{10}\cdot 3^{13}\cdot C^4\cdot R_1^2\left[\frac{1}{4}(R_1+R_2-1)^2-\frac{1}{27}R_2^3\right]^3\left[\frac{\partial(R_1,R_2)}{\partial(x,y)}\right]^{-4}H(x,y)^4.\end{aligned}$$

$$\begin{aligned}&(z_1^3+z_2^3+z_3^3)^6-540z_1^3z_2^3z_3^3(z_1^3+z_2^3+z_3^3)^3-5832z_1^6z_2^6z_3^6\\=&2^{17}\cdot 3^{21}\cdot C^6\cdot R_1^3(R_1+R_2-1)\left[\frac{1}{4}(R_1+R_2-1)^2-\frac{1}{27}R_2^3\right]^4\\&\times\left[\frac{\partial(R_1,R_2)}{\partial(x,y)}\right]^{-6}H(x,y)^6.\end{aligned}$$

$\square$

Let us consider the particular Hessian polyhedral equations:

$$\begin{cases}432\dfrac{C_9^2}{C_6^3}=x,\\ 3\dfrac{C_{12}}{C_6^2}=y.\end{cases}\tag{4.28}$$

In the projective coordinates, they are given by

$$\begin{cases}432\dfrac{C_9(z_1,z_2,z_3)^2}{C_6(z_1,z_2,z_3)^3}=x,\\ 3\dfrac{C_{12}(z_1,z_2,z_3)}{C_6(z_1,z_2,z_3)^2}=y.\end{cases}$$

Writing explicitly, we have

$$\begin{cases}x=432\dfrac{(\xi^3-\eta^3)^2(\eta^3-1)^2(1-\xi^3)^2}{[\xi^6+\eta^6+1-10(\xi^3\eta^3+\xi^3+\eta^3)]^3},\\ y=3\dfrac{(\xi^3+\eta^3+1)[(\xi^3+\eta^3+1)^3+216\xi^3\eta^3]}{[\xi^6+\eta^6+1-10(\xi^3\eta^3+\xi^3+\eta^3)]^2}.\end{cases}$$

The Hessian polyhedral irrationality (ξ, η) is defined by the equations:

$$\begin{cases} 432\dfrac{C_9^2}{C_6^3} = J_1, \\ 3\dfrac{C_{12}}{C_6^2} = J_2, \end{cases} \tag{4.29}$$

where J_1 and J_2 are J-invariants (see [Y]):

$$\begin{cases} J_1 = J_1(\lambda_1, \lambda_2) = \dfrac{\lambda_2^2(\lambda_2 - 1)^2}{\lambda_1^2(\lambda_1 - 1)^2(\lambda_1 - \lambda_2)^2}, \\ J_2 = J_2(\lambda_1, \lambda_2) = \dfrac{\lambda_1^2(\lambda_1 - 1)^2}{\lambda_2^2(\lambda_2 - 1)^2(\lambda_2 - \lambda_1)^2}. \end{cases} \tag{4.30}$$

The corresponding Picard curve is

$$y^3 = x(x-1)(x-\lambda_1)(x-\lambda_2). \tag{4.31}$$

Here, $\lambda_1 = \lambda_1(\xi, \eta)$ and $\lambda_2 = \lambda_2(\xi, \eta)$ are Picard modular functions.

Denote by

$$(F, G, H) := \frac{\partial(F, G, H)}{\partial(z_1, z_2, z_3)}. \tag{4.32}$$

Proposition 4.3 (Main Proposition 2). *In the projective coordinates, the following duality formulas hold*:

$$\begin{cases} (C_6, C_9, C_{12}) = -2^5 \cdot 3^4 \cdot \mathfrak{C}_{12}^2, \\ (C_6, C_9, C_{18}) = -2^4 \cdot 3^5 \cdot C_6 \mathfrak{C}_{12}^2, \\ (C_6, C_{12}, C_{18}) = 2^9 \cdot 3^7 \cdot C_9 \mathfrak{C}_{12}^2, \\ (C_9, C_{12}, C_{18}) = 2^4 \cdot 3^5 \cdot (C_6^2 - C_{12}) \mathfrak{C}_{12}^2. \end{cases} \tag{4.33}$$

$$\begin{cases} (C_6, C_9, \mathfrak{C}_{12}) = \dfrac{3}{2}(C_{12}^2 - C_6 C_{18}), \\ (C_6, C_{12}, \mathfrak{C}_{12}) = 432 C_9 C_{18}, \\ (C_9, C_{12}, \mathfrak{C}_{12}) = \dfrac{3}{2} C_{18}(C_6^2 - C_{12}). \end{cases} \tag{4.34}$$

$$\begin{cases} (C_6, C_{18}, \mathfrak{C}_{12}) = 648 C_9 C_{12}^2, \\ (C_9, C_{18}, \mathfrak{C}_{12}) = \dfrac{9}{4} C_{12}^2 (C_6^2 - C_{12}), \\ (C_{12}, C_{18}, \mathfrak{C}_{12}) = 0. \end{cases} \tag{4.35}$$

Proof. Put $X = z_1^3$, $Y = z_2^3$ and $Z = z_3^3$. Then

$$\begin{cases} \dfrac{\partial C_6}{\partial z_1} = 6z_1^2(X - 5Y - 5Z), \\ \dfrac{\partial C_6}{\partial z_2} = 6z_2^2(Y - 5Z - 5X), \\ \dfrac{\partial C_6}{\partial z_3} = 6z_3^2(Z - 5X - 5Y). \end{cases}$$

$$\begin{cases} \dfrac{\partial C_9}{\partial z_1} = 3z_1^2(Y - Z)(Y + Z - 2X), \\ \dfrac{\partial C_9}{\partial z_2} = 3z_2^2(Z - X)(Z + X - 2Y), \\ \dfrac{\partial C_9}{\partial z_3} = 3z_3^2(X - Y)(X + Y - 2Z). \end{cases}$$

$$\begin{cases} \dfrac{\partial C_{12}}{\partial z_1} = 12z_1^2[(X + Y + Z)^3 + 54YZ(2X + Y + Z)], \\ \dfrac{\partial C_{12}}{\partial z_2} = 12z_2^2[(X + Y + Z)^3 + 54ZX(2Y + Z + X)], \\ \dfrac{\partial C_{12}}{\partial z_3} = 12z_3^2[(X + Y + Z)^3 + 54XY(2Z + X + Y)]. \end{cases}$$

We have

$$\frac{\partial(C_6, C_9, C_{12})}{\partial(z_1, z_2, z_3)} = 3 \cdot 6 \cdot 12 \cdot z_1^2 z_2^2 z_3^2 \cdot \Delta.$$

Here,

$$\begin{aligned} \Delta =& [(X + Y + Z)^3 + 54YZ(2X + Y + Z)]\times \\ & \times [-(X + Y + Z)^3 - 9X(X + Y + Z)^2 + 27X^2(Y + Z) + 27X(Y^2 + Z^2)]+ \\ & +[(X + Y + Z)^3 + 54ZX(2Y + Z + X)]\times \\ & \times [-(X + Y + Z)^3 - 9Y(X + Y + Z)^2 + 27Y^2(Z + X) + 27Y(Z^2 + X^2)]+ \\ & +[(X + Y + Z)^3 + 54XY(2Z + X + Y)]\times \\ & \times [-(X + Y + Z)^3 - 9Z(X + Y + Z)^2 + 27Z^2(X + Y) + 27Z(X^2 + Y^2)]. \end{aligned}$$

In fact,

$$\begin{aligned}\Delta =& -3(X+Y+Z)^6+\\ &-54(X+Y+Z)^3[YZ(2X+Y+Z)+ZX(2Y+Z+X)+XY(2Z+X+Y)]+\\ &-9(X+Y+Z)^5\cdot(X+Y+Z)+\\ &-486XYZ(X+Y+Z)^2[(2X+Y+Z)+(2Y+Z+X)+(2Z+X+Y)]+\\ &+27(X+Y+Z)^3[X^2(Y+Z)+Y^2(Z+X)+Z^2(X+Y)]+\\ &+1458XYZ[X(Y+Z)(2X+Y+Z)+Y(Z+X)(2Y+Z+X)+\\ &\qquad\qquad +Z(X+Y)(2Z+X+Y)]+\\ &+27(X+Y+Z)^3[X(Y^2+Z^2)+Y(Z^2+X^2)+Z(X^2+Y^2)]+\\ &+1458XYZ[(2X+Y+Z)(Y^2+Z^2)+(2Y+Z+X)(Z^2+X^2)+\\ &\qquad\qquad +(2Z+X+Y)(X^2+Y^2)].\end{aligned}$$

Hence,

$$\begin{aligned}\Delta =& -3(X+Y+Z)^6+\\ &-54(X+Y+Z)^3[(X+Y+Z)(XY+YZ+ZX)+3XYZ]+\\ &-9(X+Y+Z)^6+\\ &-486XYZ(X+Y+Z)^2\cdot 4(X+Y+Z)+\\ &+27(X+Y+Z)^3[(X+Y+Z)(XY+YZ+ZX)-3XYZ]+\\ &+1458XYZ[3(X+Y+Z)(XY+YZ+ZX)-3XYZ]+\\ &+27(X+Y+Z)^3[(X+Y+Z)(XY+YZ+ZX)-3XYZ]+\\ &+1458XYZ[2(X+Y+Z)^3-3(X+Y+Z)(XY+YZ+ZX)-3XYZ].\end{aligned}$$

Put

$$\sigma_1 = X+Y+Z, \quad \sigma_2 = XY+YZ+ZX, \quad \sigma_3 = XYZ.$$

Then

$$\Delta = -12\sigma_1^6 + 648\sigma_1^3\sigma_3 - 8748\sigma_3^2 = -12(27\sigma_3-\sigma_1^3)^2.$$

Thus,

$$\frac{\partial(C_6,C_9,C_{12})}{\partial(z_1,z_2,z_3)} = -3\cdot 6\cdot 12^2\cdot[z_1z_2z_3(27\sigma_3-\sigma_1^3)]^2 = -2^5\cdot 3^4\cdot \mathfrak{C}_{12}^2.$$

Note that

$$C_{18} = \frac{1}{2}(432C_9^2 - C_6^3 + 3C_6C_{12}).$$

This gives that

$$\frac{\partial C_{18}}{\partial z_j} = \frac{3}{2}(C_{12} - C_6^2)\frac{\partial C_6}{\partial z_j} + 432 C_9 \frac{\partial C_9}{\partial z_j} + \frac{3}{2} C_6 \frac{\partial C_{12}}{\partial z_j}, \quad (j = 1, 2, 3).$$

Therefore,

$$\frac{\partial(C_6, C_9, C_{18})}{\partial(z_1, z_2, z_3)} = \frac{3}{2} C_6 \frac{\partial(C_6, C_9, C_{12})}{\partial(z_1, z_2, z_3)} = -2^4 \cdot 3^5 \cdot C_6 \mathfrak{C}_{12}^2.$$

$$\frac{\partial(C_6, C_{12}, C_{18})}{\partial(z_1, z_2, z_3)} = -432 C_9 \frac{\partial(C_6, C_9, C_{12})}{\partial(z_1, z_2, z_3)} = 2^9 \cdot 3^7 \cdot C_9 \mathfrak{C}_{12}^2.$$

$$\frac{\partial(C_9, C_{12}, C_{18})}{\partial(z_1, z_2, z_3)} = \frac{3}{2}(C_{12} - C_6^2) \frac{\partial(C_6, C_9, C_{12})}{\partial(z_1, z_2, z_3)} = -2^4 \cdot 3^5 \cdot (C_{12} - C_6^2) \mathfrak{C}_{12}^2.$$

By

$$1728 \mathfrak{C}_{12}^3 = C_{18}^2 - C_{12}^3,$$

we find that

$$3 \cdot 1728 \mathfrak{C}_{12}^2 \frac{\partial \mathfrak{C}_{12}}{\partial z_j} = 2 C_{18} \frac{\partial C_{18}}{\partial z_j} - 3 C_{12}^2 \frac{\partial C_{12}}{\partial z_j}, \quad (j = 1, 2, 3).$$

This gives the other identities. □

COROLLARY 4.4. *The invariants C_6, C_9, C_{12}, $\mathfrak{C}_{12}^2$ and C_{18} can be represented by:*

(4.36) $$\mathfrak{C}_{12}^2 = -\frac{1}{2592}(C_6, C_9, C_{12}).$$

(4.37) $$C_6 = \frac{2}{3} \frac{(C_6, C_9, C_{18})}{(C_6, C_9, C_{12})}.$$

(4.38) $$C_9 = -\frac{1}{432} \frac{(C_6, C_{12}, C_{18})}{(C_6, C_9, C_{12})}.$$

(4.39) $$C_{12} = \frac{2}{3} \frac{(C_9, C_{12}, C_{18})}{(C_6, C_9, C_{12})} + \frac{4}{9} \frac{(C_6, C_9, C_{18})^2}{(C_6, C_9, C_{12})^2}.$$

(4.40) $$C_{18} = -\frac{(C_6, C_9, C_{12})(C_6, C_{12}, \mathfrak{C}_{12})}{(C_6, C_{12}, C_{18})}.$$

Proof. We find that

$$\frac{(C_6,C_9,C_{18})}{(C_6,C_9,C_{12})}=\frac{3}{2}C_6.$$

$$\frac{(C_6,C_{12},C_{18})}{(C_6,C_9,C_{12})}=-432C_9.$$

$$\frac{(C_9,C_{12},C_{18})}{(C_6,C_9,C_{12})}=\frac{3}{2}(C_{12}-C_6^2).$$

$$C_{18}=\frac{(C_6,C_{12},\mathfrak{C}_{12})}{432C_9}.$$

□

PROPOSITION 4.5. *The following algebraic equations of the sixth degree hold*:

$$\begin{aligned}&27(C_6,C_9,\mathfrak{C}_{12})(C_6,C_9,C_{12})^4(C_6,C_{12},C_{18})\\&=18(C_9,C_{12},C_{18})^2(C_6,C_9,C_{12})^2(C_6,C_{12},C_{18})+8(C_6,C_9,C_{18})^4(C_6,C_{12},C_{18})+\\&+24(C_9,C_{12},C_{18})(C_6,C_9,C_{18})^2(C_6,C_9,C_{12})(C_6,C_{12},C_{18})+\\&+27(C_6,C_9,C_{18})(C_6,C_{12},\mathfrak{C}_{12})(C_6,C_9,C_{12})^4.\end{aligned}\tag{4.41}$$

$$\begin{aligned}&27(C_6,C_{18},\mathfrak{C}_{12})(C_6,C_9,C_{12})^5+8(C_6,C_{12},C_{18})(C_6,C_9,C_{18})^4+\\&+18(C_6,C_{12},C_{18})(C_9,C_{12},C_{18})^2(C_6,C_9,C_{12})^2+\\&+24(C_6,C_{12},C_{18})(C_9,C_{12},C_{18})(C_6,C_9,C_{18})^2(C_6,C_9,C_{12})=0.\end{aligned}\tag{4.42}$$

$$\begin{aligned}&27(C_9,C_{18},\mathfrak{C}_{12})(C_6,C_9,C_{12})^5+18(C_9,C_{12},C_{18})^3(C_6,C_9,C_{12})^2+\\&+8(C_6,C_9,C_{18})^4(C_9,C_{12},C_{18})+24(C_9,C_{12},C_{18})^2(C_6,C_9,C_{18})^2(C_6,C_9,C_{12})=0.\end{aligned}\tag{4.43}$$

Proof. They are obtained from the identities:

$$(C_6,C_9,\mathfrak{C}_{12})=\frac{3}{2}(C_{12}^2-C_6C_{18}).$$

$$(C_6,C_{18},\mathfrak{C}_{12})=648C_9C_{12}^2.$$

$$(C_9,C_{18},\mathfrak{C}_{12})=\frac{9}{4}C_{12}^2(C_6^2-C_{12}).$$

□

Proposition 4.6. *The invariants* C_6, C_9, C_{12}, $\mathfrak{C}_{12}$ *and* C_{18} *satisfy the following duality formula*:

$$(C_9, C_{12}, \mathfrak{C}_{12})(C_6, C_{12}, C_{18}) = (C_6, C_{12}, \mathfrak{C}_{12})(C_9, C_{12}, C_{18}). \tag{4.44}$$

Proof. It is obtained from the equation

$$(C_9, C_{12}, \mathfrak{C}_{12}) = \frac{3}{2} C_{18}(C_6^2 - C_{12}).$$

□

Put

$$\begin{cases} Y_1 = (C_6, C_{12}, \mathfrak{C}_{12}), \\ Y_2 = (C_9, C_{12}, \mathfrak{C}_{12}), \\ Y_3 = (C_6, C_{12}, C_{18}), \\ Y_4 = (C_9, C_{12}, C_{18}). \end{cases} \tag{4.45}$$

$$\begin{cases} X_1 = (C_6, C_9, C_{12}), \\ X_2 = (C_6, C_9, \mathfrak{C}_{12}), \\ X_3 = (C_6, C_9, C_{18}), \\ X_4 = (C_6, C_{18}, \mathfrak{C}_{12}), \\ X_5 = (C_9, C_{18}, \mathfrak{C}_{12}). \end{cases} \tag{4.46}$$

Then Proposition 4.5 and Proposition 4.6 give that

$$18 X_1^2 Y_3 Y_4^2 + 24 X_1 X_3^2 Y_3 Y_4 + 27 X_1^4 X_3 Y_1 + (8 X_3^4 - 27 X_1^4 X_2) Y_3 = 0, \tag{4.47}$$

$$18 X_1^2 Y_3 Y_4^2 + 24 X_1 X_3^2 Y_3 Y_4 + 8 X_3^4 Y_3 + 27 X_1^5 X_4 = 0, \tag{4.48}$$

$$18 X_1^2 Y_4^3 + 24 X_1 X_3^2 Y_4^2 + 8 X_3^4 Y_4 + 27 X_1^5 X_5 = 0, \tag{4.49}$$

$$Y_2 Y_3 - Y_1 Y_4 = 0. \tag{4.50}$$

(4.47) – (4.48) gives that

$$X_3 Y_1 - X_2 Y_3 = X_1 X_4. \tag{4.51}$$

(4.47) × Y_4 − (4.49) × Y_3 gives that

$$X_3Y_1Y_4 - X_2Y_3Y_4 = X_1X_5Y_3. \tag{4.52}$$

By (4.51), we have

$$X_4Y_4 = X_5Y_3. \tag{4.53}$$

Thus, we get the following theorem:

THEOREM 4.7 (MAIN THEOREM 3). *The functions* X_1, X_2, X_3, X_4, X_5 *and* Y_1, Y_2, Y_3, Y_4 *satisfy the system of algebraic equations of the sixth degree*:

$$\begin{cases} 18X_1^2Y_4^3 + 24X_1X_3^2Y_4^2 + 8X_3^4Y_4 + 27X_1^5X_5 = 0, \\ Y_2Y_3 = Y_1Y_4, \\ X_4Y_4 = X_5Y_3, \\ X_3Y_1 - X_2Y_3 = X_1X_4. \end{cases} \tag{4.54}$$

The first equation in (4.52) is an algebraic hypersurface of the sixth degree in $\mathbb{C}^4$. The second and the third equations are the algebraic hypersurfaces of the second degree in $\mathbb{C}^4$. The last equation is an algebraic hypersurface of the second degree in $\mathbb{C}^6$.

Let

$$\mathcal{A} = \{(C_6, C_9, C_{12}, \mathfrak{C}_{12}, C_{18}) \in \mathbb{C}^5\} \tag{4.55}$$

be the configuration space of the Hessian invariants. Let

$$\mathcal{B} = \{(X_1, X_2, X_3, X_4, X_5, Y_1, Y_2, Y_3, Y_4) \in \mathbb{C}^9\} \tag{4.56}$$

be the dual configuration space of $\mathcal{A}$. Then

$$\dim\mathcal{A} = 3, \quad \dim\mathcal{B} = 5. \tag{4.57}$$

Let us consider the Hessian polyhedral equations:

$$Z_1 = 432\frac{C_9^2}{C_6^3}, \quad Z_2 = 3\frac{C_{12}}{C_6^2}. \tag{4.58}$$

Note that the relation

$$432C_9^2 = C_6^3 - 3C_6C_{12} + 2C_{18}$$

implies that

$$Z_1 + Z_2 - 1 = 2\frac{C_{18}}{C_6^3}. \tag{4.59}$$

The relation

$$1728\mathfrak{C}_{12}^3 = C_{18}^2 - C_{12}^3$$

gives that

$$12\frac{\mathfrak{C}_{12}}{C_6^2} = \sqrt[3]{\frac{1}{4}(Z_1 + Z_2 - 1)^2 - \frac{1}{27}Z_2^3}. \tag{4.60}$$

We find that

$$\begin{aligned} &Z_1^2 : Z_2^3 : (Z_1 + Z_2 - 1)^2 : [27(Z_1 + Z_2 - 1)^2 - 4Z_2^3] : 1 \\ =&432^2 C_9^4 : 27C_{12}^3 : 4C_{18}^2 : 432^2\mathfrak{C}_{12}^3 : C_6^6 \\ =&432^2(\xi^3 - \eta^3)^4(\eta^3 - 1)^4(1 - \xi^3)^4 \\ &:27(\xi^3 + \eta^3 + 1)^3[(\xi^3 + \eta^3 + 1)^3 + 216\xi^3\eta^3]^3 \\ &:4[(\xi^3 + \eta^3 + 1)^6 - 540\xi^3\eta^3(\xi^3 + \eta^3 + 1)^3 - 5832\xi^6\eta^6]^2 \\ &:432^2\xi^3\eta^3[27\xi^3\eta^3 - (\xi^3 + \eta^3 + 1)^3]^3 \\ &:[\xi^6 + \eta^6 + 1 - 10(\xi^3\eta^3 + \xi^3 + \eta^3)]^6. \end{aligned} \tag{4.61}$$

In fact,

$$Z_1 = \frac{1}{128}\frac{X_1 Y_3^2}{X_3^3}, \quad Z_2 = 3 + \frac{9}{2}\frac{X_1 Y_4}{X_3^2}. \tag{4.62}$$

Let

$$F_1 = C_{18}, \quad F_2 = \mathfrak{C}_{12}, \quad F_3 = C_{12}, \quad F_4 = C_9, \quad F_5 = C_6 \tag{4.63}$$

and

$$W = \frac{F_1 F_5}{F_3 F_4}. \tag{4.64}$$

Then

$$F_1 = \frac{Z_1^3 Z_2^6 W^6}{2^7 \cdot 3^{15}(Z_1 + Z_2 - 1)^5}, \tag{4.65}$$

$$F_3 = \frac{Z_1^2 Z_2^5 W^4}{2^4 \cdot 3^{11}(Z_1 + Z_2 - 1)^4}, \tag{4.66}$$

$$F_4=\frac{Z_1^2Z_2^3W^3}{2^5\cdot 3^9(Z_1+Z_2-1)^3},\tag{4.67}$$

$$F_5=\frac{Z_1Z_2^2W^2}{2^2\cdot 3^5(Z_1+Z_2-1)^2},\tag{4.68}$$

$$F_2=\frac{Z_1^2Z_2^4W^4}{2^6\cdot 3^{11}(Z_1+Z_2-1)^4}\sqrt[3]{\frac{1}{4}(Z_1+Z_2-1)^2-\frac{1}{27}Z_2^3}.\tag{4.69}$$

Put

$$\tau_1=\xi^3+\eta^3,\quad \tau_2=\xi\eta.\tag{4.70}$$

Then

$$\begin{cases} C_6=\tau_1^2-12\tau_2^3-10\tau_1+1,\\ C_{12}=(\tau_1+1)[(\tau_1+1)^3+216\tau_2^3],\\ \mathfrak{C}_{12}=\tau_2[27\tau_2^3-(\tau_1+1)^3],\\ C_{18}=(\tau_1+1)^6-540\tau_2^3(\tau_1+1)^3-5832\tau_2^6,\\ C_9^2=(\tau_1^2-4\tau_2^3)(\tau_2^3-\tau_1+1)^2.\end{cases}\tag{4.71}$$

Thus,

$$\begin{aligned} &Z_1^2:Z_2^3:(Z_1+Z_2-1)^2:[27(Z_1+Z_2-1)^2-4Z_2^3]:1\\ =&432^2(\tau_1^2-4\tau_2^3)^2(\tau_2^3-\tau_1+1)^4\\ &:27(\tau_1+1)^3[(\tau_1+1)^3+216\tau_2^3]^3\\ &:4[(\tau_1+1)^6-540\tau_2^3(\tau_1+1)^3-5832\tau_2^6]^2\\ &:432^2\tau_2^3[27\tau_2^3-(\tau_1+1)^3]^3\\ &:(\tau_1^2-12\tau_2^3-10\tau_1+1)^6.\end{aligned}\tag{4.72}$$

5. Invariant theory for the system of algebraic equations

In the affine coordinates, we have

$$\begin{cases} A(\xi,\eta)=\left(\frac{\eta}{\xi},\frac{1}{\xi}\right),\\ B(\xi,\eta)=\left(\frac{\xi}{\eta},\frac{1}{\eta}\right),\\ C(\xi,\eta)=(\omega\xi,\omega^2\eta),\\ D(\xi,\eta)=(\omega^2\xi,\eta),\\ E(\xi,\eta)=\left(\frac{\xi+\eta+1}{\xi+\omega^2\eta+\omega},\frac{\xi+\omega\eta+\omega^2}{\xi+\omega^2\eta+\omega}\right).\end{cases}\tag{5.1}$$

Note that $E^2 = -B$, $B^2 = I$, $A^3 = I$. We find that

$$\langle A, B\rangle = \{A, A^2, B, AB, BA, I\}. \tag{5.2}$$

Here,

$$A^2 = \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}, \quad AB = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{pmatrix}, \quad BA = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}. \tag{5.3}$$

There are three involutions:

$$B^2 = (AB)^2 = (BA)^2 = I. \tag{5.4}$$

(1) $B(\xi, \eta) = (\frac{\xi}{\eta}, \frac{1}{\eta})$.

Let $B(\xi, \eta) = (\xi, \eta)$, we find that

$$\xi = \xi\eta \quad \text{and} \quad \eta^2 = 1.$$

Hence,

$$z_1(z_2 - z_3) = 0 \quad \text{and} \quad (z_2 + z_3)(z_2 - z_3) = 0.$$

The common factor is $z_2 - z_3 = 0$.

(2) $AB(\xi, \eta) = (\frac{1}{\xi}, \frac{\eta}{\xi})$.

Let $AB(\xi, \eta) = (\xi, \eta)$, we find that

$$\xi^2 = 1 \quad \text{and} \quad \eta = \xi\eta.$$

Hence,

$$(z_3 + z_1)(z_3 - z_1) = 0 \quad \text{and} \quad z_2(z_3 - z_1) = 0.$$

The common factor is $z_3 - z_1 = 0$.

(3) $BA(\xi, \eta) = (\eta, \xi)$.

Let $BA(\xi, \eta) = (\xi, \eta)$, we find that

$$\eta = \xi \quad \text{and} \quad \xi = \eta.$$

Hence,

$$z_2 - z_1 = 0 \quad \text{and} \quad z_1 - z_2 = 0.$$

The common factor is $z_1 - z_2 = 0$.

Put

$$G(z_1, z_2, z_3) := (z_1 - z_2)(z_2 - z_3)(z_3 - z_1). \tag{5.5}$$

The Hessian of G is zero.

Note that

$$E(z_1, z_2, z_3) = \frac{1}{\sqrt{-3}}(z_1 + z_2 + z_3, z_1 + \omega z_2 + \omega^2 z_3, z_1 + \omega^2 z_2 + \omega z_3). \tag{5.6}$$

According to Maschke [Mas2], put

$$\varphi = z_1 z_2 z_3, \quad \psi = z_1^3 + z_2^3 + z_3^3, \quad \chi = z_1^3 z_2^3 + z_2^3 z_3^3 + z_3^3 z_1^3. \tag{5.7}$$

We have

$$\psi(E(z_1, z_2, z_3)) = \frac{-1}{\sqrt{-3}}(\psi + 6\varphi). \tag{5.8}$$

$$\varphi(E(z_1, z_2, z_3)) = \frac{-1}{3\sqrt{-3}}(\psi - 3\varphi). \tag{5.9}$$

Let

$$H(z_1, z_2, z_3) := \frac{1}{3}[(z_1 + z_2 + z_3)^3 + (z_1 + \omega z_2 + \omega^2 z_3)^3 + (z_1 + \omega^2 z_2 + \omega z_3)^3]. \tag{5.10}$$

$$K(z_1, z_2, z_3) := (z_1 + z_2 + z_3)(z_1 + \omega z_2 + \omega^2 z_3)(z_1 + \omega^2 z_2 + \omega z_3). \tag{5.11}$$

Then

$$H = \psi + 6\varphi, \quad K = \psi - 3\varphi. \tag{5.12}$$

The Hessian of H:

$$\text{Hessian}(H) = \begin{vmatrix} \frac{\partial^2 H}{\partial z_1^2} & \frac{\partial^2 H}{\partial z_1 \partial z_2} & \frac{\partial^2 H}{\partial z_1 \partial z_3} \\ \frac{\partial^2 H}{\partial z_1 \partial z_2} & \frac{\partial^2 H}{\partial z_2^2} & \frac{\partial^2 H}{\partial z_2 \partial z_3} \\ \frac{\partial^2 H}{\partial z_1 \partial z_3} & \frac{\partial^2 H}{\partial z_2 \partial z_3} & \frac{\partial^2 H}{\partial z_3^2} \end{vmatrix} = -216K. \tag{5.13}$$

The Hessian of K:

$$\text{Hessian}(K) = \begin{vmatrix} \frac{\partial^2 K}{\partial z_1^2} & \frac{\partial^2 K}{\partial z_1 \partial z_2} & \frac{\partial^2 K}{\partial z_1 \partial z_3} \\ \frac{\partial^2 K}{\partial z_1 \partial z_2} & \frac{\partial^2 K}{\partial z_2^2} & \frac{\partial^2 K}{\partial z_2 \partial z_3} \\ \frac{\partial^2 K}{\partial z_1 \partial z_3} & \frac{\partial^2 K}{\partial z_2 \partial z_3} & \frac{\partial^2 K}{\partial z_3^2} \end{vmatrix} = -54K. \tag{5.14}$$

In fact,

$$\begin{cases} C_6 = \psi^2 - 12\chi, \\ C_{12} = \psi(\psi^3 + 216\varphi^3), \\ \mathfrak{C}_{12} = \varphi(27\varphi^3 - \psi^3), \\ C_{18} = \psi^6 - 540\varphi^3\psi^3 - 5832\varphi^6. \end{cases} \tag{5.15}$$

Theorem 5.1 (Main Theorem 4). *The invariants G, H and K satisfy the following algebraic equations, which are the form-theoretic resolvents* (*algebraic resolvents*) *of G, H, K*:

$$\begin{cases} 4G^3 + H^2G - C_6G - 4C_9 = 0, \\ H(H^3 + 8K^3) - 9C_{12} = 0, \\ K(K^3 - H^3) - 27\mathfrak{C}_{12} = 0. \end{cases} \tag{5.16}$$

Proof. We have

$$\begin{aligned} \frac{C_9}{G} =&(z_1^2 + z_1z_2 + z_2^2)(z_2^2 + z_2z_3 + z_3^2)(z_3^2 + z_3z_1 + z_1^2) \\ =&3z_1^2z_2^2z_3^2 + z_1^4z_2^2 + z_2^4z_3^2 + z_3^4z_1^2 + z_1^2z_2^4 + z_2^2z_3^4 + z_3^2z_1^4 + \\ &+ z_1^3z_2^3 + z_2^3z_3^3 + z_3^3z_1^3 + z_1^4z_2z_3 + z_2^4z_3z_1 + z_3^4z_1z_2 + \\ &+ 2z_1z_2z_3(z_1^2z_2 + z_2^2z_3 + z_3^2z_1 + z_1z_2^2 + z_2z_3^2 + z_3z_1^2). \end{aligned}$$

Set

$$\sigma_1 = z_1 + z_2 + z_3, \quad \sigma_2 = z_1z_2 + z_2z_3 + z_3z_1.$$

We have

$$z_1^2z_2 + z_2^2z_3 + z_3^2z_1 + z_1z_2^2 + z_2z_3^2 + z_3z_1^2 = \sigma_1\sigma_2 - 3\varphi.$$

$$z_1^4z_2^2 + z_2^4z_3^2 + z_3^4z_1^2 + z_1^2z_2^4 + z_2^2z_3^4 + z_3^2z_1^4 = (\sigma_1^2 - 2\sigma_2)(\sigma_2^2 - 2\varphi\sigma_1) - 3\varphi^2.$$

Thus,

$$\frac{C_9}{G} = 3\varphi^2 + \chi + \varphi\psi + (\sigma_1^2 - 2\sigma_2)(\sigma_2^2 - 2\varphi\sigma_1) - 3\varphi^2 + 2\varphi(\sigma_1\sigma_2 - 3\varphi).$$

Note that

$$\psi = \sigma_1(\sigma_1^2 - 3\sigma_2) + 3\varphi, \quad \chi = \sigma_2(\sigma_2^2 - 3\varphi\sigma_1) + 3\varphi^2.$$

This gives that

$$(\sigma_1^2 - 2\sigma_2)(\sigma_2^2 - 2\varphi\sigma_1) = -2\varphi(\psi - 3\varphi) - 2\varphi\sigma_1\sigma_2 + \sigma_2^2(\sigma_1^2 - 2\sigma_2).$$

Hence,

$$\frac{C_9}{G} = \chi - \varphi\psi + \sigma_2^2(\sigma_1^2 - 2\sigma_2).$$

In fact,

$$\sigma_2^3 = \chi - 3\varphi^2 + 3\varphi\sigma_1\sigma_2.$$

This implies that

$$\begin{aligned}\frac{C_9}{G} =& \chi - \varphi\psi + \sigma_1^2\sigma_2^2 - 2\sigma_2^3 \\ =& -\chi - \varphi\psi + 6\varphi^2 + \sigma_1^2\sigma_2^2 - 6\varphi\sigma_1\sigma_2.\end{aligned}$$

We have

$$G = (z_1z_2^2 + z_2z_3^2 + z_3z_1^2) - (z_1^2z_2 + z_2^2z_3 + z_3^2z_1).$$

Hence,

$$\begin{aligned}G^2 =& (z_1z_2^2 + z_2z_3^2 + z_3z_1^2 + z_1^2z_2 + z_2^2z_3 + z_3^2z_1)^2 + \\ & - 4(z_1z_2^2 + z_2z_3^2 + z_3z_1^2)(z_1^2z_2 + z_2^2z_3 + z_3^2z_1) \\ =& (\sigma_1\sigma_2 - 3\varphi)^2 - 4(\chi + 3\varphi^2 + \varphi\psi) \\ =& \sigma_1^2\sigma_2^2 - 6\varphi\sigma_1\sigma_2 - 3\varphi^2 - 4\chi - 4\varphi\psi.\end{aligned}$$

Therefore,

$$\frac{C_9}{G} - G^2 = 3(\chi + \varphi\psi + 3\varphi^2).$$

We find that

$$4\left(\frac{C_9}{G} - G^2\right) + C_6 = (\psi + 6\varphi)^2 = H^2.$$

Hence,

$$4G^3 + H^2G - C_6G - 4C_9 = 0.$$

By

$$H = \psi + 6\varphi, \quad K = \psi - 3\varphi,$$

we have

$$\varphi = \frac{1}{9}(H - K), \quad \psi = \frac{1}{3}(H + 2K).$$

The relations

$$C_{12} = \psi(\psi^3 + 216\varphi^3), \quad \mathfrak{C}_{12} = \varphi(27\varphi^3 - \psi^3)$$

give that

$$H(H^3 + 8K^3) = 9C_{12}, \quad K(K^3 - H^3) = 27\mathfrak{C}_{12}.$$

□

Let

$$r_1 = \frac{G^2}{C_6}, \quad r_2 = \frac{H^2}{C_6}, \quad r_3 = \frac{K^2}{C_6}. \tag{5.17}$$

PROPOSITION 5.2. *The following identities hold*:

$$\begin{cases} Z_1 = 27r_1(4r_1 + r_2 - 1)^2, \\ Z_2 = \frac{1}{3}r_2^2 + \frac{8}{3}r_3 \cdot \sqrt{r_2 r_3}, \\ \sqrt[3]{\frac{1}{4}(Z_1 + Z_2 - 1)^2 - \frac{1}{27}Z_2^3} = \frac{4}{9}(r_3^2 - r_2 \cdot \sqrt{r_2 r_3}). \end{cases} \tag{5.18}$$

Proof. By

$$4G^3 + H^2 G - C_6 G - 4C_9 = 0,$$

we have

$$C_9 = G\left(G^2 + \frac{1}{4}H^2 - \frac{1}{4}C_6\right).$$

Thus

$$\frac{C_9^2}{C_6^3} = \frac{G^2}{C_6}\left(\frac{G^2}{C_6} + \frac{1}{4}\frac{H^2}{C_6} - \frac{1}{4}\right)^2.$$

This gives that

$$\frac{1}{432}Z_1 = r_1\left(r_1 + \frac{1}{4}r_2 - \frac{1}{4}\right)^2.$$

Hence,

$$Z_1 = 27r_1(4r_1 + r_2 - 1)^2.$$

By

$$C_{12} = \frac{1}{9}H(H^3 + 8K^3),$$

we have

$$\frac{C_{12}}{C_6^2} = \frac{1}{9}\left(\frac{H^2}{C_6}\right)^{\frac{1}{2}}\left[\left(\frac{H^2}{C_6}\right)^{\frac{3}{2}} + 8\left(\frac{K^2}{C_6}\right)^{\frac{3}{2}}\right].$$

This gives that

$$Z_2 = \frac{1}{3}r_2^2 + \frac{8}{3}r_3 \cdot \sqrt{r_2 r_3}.$$

By

$$\mathfrak{C}_{12} = \frac{1}{27}K(K^3 - H^3),$$

we have

$$\frac{\mathfrak{C}_{12}}{C_6^2}=\frac{1}{27}\left(\frac{K^2}{C_6}\right)^{\frac{1}{2}}\left[\left(\frac{K^2}{C_6}\right)^{\frac{3}{2}}-\left(\frac{H^2}{C_6}\right)^{\frac{3}{2}}\right].$$

This gives that

$$\sqrt[3]{\frac{1}{4}(Z_1+Z_2-1)^2-\frac{1}{27}Z_2^3}=\frac{4}{9}(r_3^2-r_2\cdot\sqrt{r_2r_3}).$$

□

This gives the following theorem:

THEOREM 5.3 (MAIN THEOREM 5). *The function-theoretic resolvents* (*analytic resolvents*) *of* Z_1 *and* Z_2 *are given by*:

$$\begin{aligned}&Z_1^2:Z_2^3:(Z_1+Z_2-1)^2:[27(Z_1+Z_2-1)^2-4Z_2^3]:1\\=&729^2r_1^2(4r_1+r_2-1)^4:27(r_2^2+8r_3\cdot\sqrt{r_2r_3})^3\\&:[256(r_3^2-r_2\cdot\sqrt{r_2r_3})^3+4(r_2^2+8r_3\cdot\sqrt{r_2r_3})^3]\\&:6912(r_3^2-r_2\cdot\sqrt{r_2r_3})^3:729.\end{aligned}\tag{5.19}$$

For the Hessian polyhedral equations, the corresponding J-invariants (J_1,J_2) satisfy:

$$\begin{aligned}&J_1^2:J_2^3:(J_1+J_2-1)^2:[27(J_1+J_2-1)^2-4J_2^3]:1\\=&729^2v_1^2(4v_1+v_2-1)^4:27(v_2^2+8v_3\cdot\sqrt{v_2v_3})^3\\&:[256(v_3^2-v_2\cdot\sqrt{v_2v_3})^3+4(v_2^2+8v_3\cdot\sqrt{v_2v_3})^3]\\&:6912(v_3^2-v_2\cdot\sqrt{v_2v_3})^3:729,\end{aligned}\tag{5.20}$$

where

$$v_1=v_1(\omega_1,\omega_2),\quad v_2=v_2(\omega_1,\omega_2),\quad v_3=v_3(\omega_1,\omega_2)$$

are three Picard modular functions with $(\omega_1,\omega_2)\in\mathfrak{S}_2=\{(z_1,z_2)\in\mathbb{C}^2:z_1+\overline{z_1}-z_2\overline{z_2}>0\}$.

Put

$$u=\frac{C_6G}{C_9},\quad v=\frac{C_9H}{C_{12}},\quad w=\frac{C_9K}{\mathfrak{C}_{12}}.\tag{5.21}$$

We will denote the next three equations as the resolvents of the u, v and w's.

THEOREM 5.4 (MAIN THEOREM 6). *The functions* u, v *and* w *satisfy the following algebraic equations, which are the analytic and algebraic resolvents of* u, v *and* w:

$$\begin{cases}Z_1^2u^3+5184Z_2^2uv^2-108Z_1u-432Z_1=0,\\6912Z_2^3v^4+8[27(Z_1+Z_2-1)^2-4Z_2^3]vw^3-9Z_1^2=0,\\{[27(Z_1+Z_2-1)^2-4Z_2^3]}w^4-6912Z_2^3v^3w-27Z_1^2=0.\end{cases}\tag{5.22}$$

Proof. We have

$$G = \frac{C_9 u}{C_6}, \quad H = \frac{C_{12} v}{C_9}, \quad K = \frac{\mathfrak{C}_{12} w}{C_9}.$$

By

$$4G^3 + H^2 G - C_6 G - 4C_9 = 0,$$

we have

$$4\frac{C_9^3 u^3}{C_6^3} + \frac{C_{12}^2 v^2}{C_9^2} \cdot \frac{C_9 u}{C_6} - C_6 \cdot \frac{C_9 u}{C_6} - 4C_9 = 0,$$

i.e.,

$$4C_9^4 u^3 + C_6^2 C_{12}^2 v^2 u - C_6^3 C_9^2 u - 4C_6^3 C_9^2 = 0.$$

Thus,

$$4\frac{C_9^4}{C_6^6} u^3 + \frac{C_{12}^2}{C_6^4} uv^2 - \frac{C_9^2}{C_6^3} u - 4\frac{C_9^2}{C_6^3} = 0.$$

This implies that

$$4\left(\frac{Z_1}{432}\right)^2 u^3 + \left(\frac{Z_2}{3}\right)^2 uv^2 - \frac{Z_1}{432} u - 4\frac{Z_1}{432} = 0.$$

Hence,

$$Z_1^2 u^3 + 5184 Z_2^2 uv^2 - 108 Z_1 u - 432 Z_1 = 0.$$

Note that

$$9C_{12} = H(H^3 + 8K^3) = \frac{C_{12} v}{C_9} \left(\frac{C_{12}^3 v^3}{C_9^3} + 8\frac{\mathfrak{C}_{12}^3 w^3}{C_9^3} \right).$$

i.e.,

$$9C_9^4 = C_{12}^3 v^4 + 8\mathfrak{C}_{12}^3 v w^3.$$

Thus,

$$9\left(\frac{C_9^2}{C_6^3}\right)^2 = \left(\frac{C_{12}}{C_6^2}\right)^3 v^4 + 8\left(\frac{\mathfrak{C}_{12}}{C_6^2}\right)^3 v w^3.$$

This implies that

$$9\left(\frac{Z_1}{432}\right)^2 = \left(\frac{Z_2}{3}\right)^3 v^4 + 8 \cdot \frac{1}{1728} \left[\frac{1}{4}(Z_1 + Z_2 - 1)^2 - \frac{1}{27} Z_2^3 \right] v w^3.$$

Hence,

$$9Z_1^2 = 6912 Z_2^3 v^4 + 8[27(Z_1 + Z_2 - 1)^2 - 4Z_2^3] v w^3.$$

Note that

$$27\mathfrak{C}_{12}=K(K^3-H^3)=\frac{\mathfrak{C}_{12}w}{C_9}\left(\frac{\mathfrak{C}_{12}^3w^3}{C_9^3}-\frac{C_{12}^3v^3}{C_9^3}\right),$$

i.e.,

$$27C_9^4=\mathfrak{C}_{12}^3w^4-C_{12}^3v^3w.$$

Thus,

$$27\left(\frac{C_9^2}{C_6^3}\right)^2=\left(\frac{\mathfrak{C}_{12}}{C_6^2}\right)^3w^4-\left(\frac{C_{12}}{C_6^2}\right)^3v^3w.$$

This implies that

$$27\cdot\left(\frac{Z_1}{432}\right)^2=\frac{1}{1728}\left[\frac{1}{4}(Z_1+Z_2-1)^2-\frac{1}{27}Z_2^3\right]w^4-\left(\frac{Z_2}{3}\right)^3v^3w.$$

Hence,

$$27Z_1^2=[27(Z_1+Z_2-1)^2-4Z_2^3]w^4-6912Z_2^3v^3w.$$

□

THEOREM 5.5 (MAIN THEOREM 7). *The functions* u, v *and* w *can be expressed as the algebraic functions of* r_1, r_2 *and* r_3:

$$\begin{cases} u=\dfrac{4}{4r_1+r_2-1}, \\ v=\dfrac{9}{4}\dfrac{(4r_1+r_2-1)\sqrt{r_1}}{r_2\sqrt{r_2}+8r_3\sqrt{r_3}}, \\ w=\dfrac{27}{4}\dfrac{(4r_1+r_2-1)\sqrt{r_1}}{r_3\sqrt{r_3}-r_2\sqrt{r_2}}. \end{cases} \tag{5.23}$$

Proof. We have

$$u^2=\frac{C_6^2G^2}{C_9^2}=\frac{\frac{G^2}{C_6}}{\frac{C_9^2}{C_6^3}}=\frac{r_1}{\frac{1}{432}Z_1}=\frac{16}{(4r_1+r_2-1)^2}.$$

So,

$$u=\frac{4}{4r_1+r_2-1}.$$

On the other hand,

$$v^2=\frac{C_9^2H^2}{C_{12}^2}=\frac{\frac{C_9^2}{C_6^3}}{(\frac{C_{12}}{C_6^2})^2}\cdot\frac{H^2}{C_6}=\frac{\frac{1}{432}Z_1}{(\frac{1}{3}Z_2)^2}r_2=\frac{\frac{1}{16}r_1(4r_1+r_2-1)^2}{(\frac{1}{9}r_2^2+\frac{8}{9}r_3\sqrt{r_2r_3})^2}r_2.$$

So,

$$v = \frac{9(4r_1 + r_2 - 1)\sqrt{r_1}}{4r_2\sqrt{r_2} + 32r_3\sqrt{r_3}}.$$

Similarly,

$$\begin{aligned} w^2 =& \frac{C_9^2 K^2}{\mathfrak{C}_{12}^2} = \frac{\frac{C_9^2}{C_6^3}}{(\frac{\mathfrak{C}_{12}}{C_6^2})^2} \cdot \frac{K^2}{C_6} = \frac{\frac{1}{432}Z_1}{\frac{1}{144}[\frac{1}{4}(Z_1 + Z_2 - 1)^2 - \frac{1}{27}Z_2^3]^{\frac{2}{3}}} r_3 \\ =& \frac{\frac{1}{16}r_1(4r_1 + r_2 - 1)^2 r_3}{\frac{1}{144}(\frac{4}{9})^2(r_3^2 - r_2 \cdot \sqrt{r_2 r_3})^2}. \end{aligned}$$

So,

$$w = \frac{27}{4}\frac{(4r_1 + r_2 - 1)\sqrt{r_1}}{r_3\sqrt{r_3} - r_2\sqrt{r_2}}.$$

$\square$

Set

$$M := M_\infty = 81\sqrt{-3}\varphi^3, \quad N := N_\infty = 3\sqrt{-3}\psi^3. \tag{5.24}$$

Then

$$M_0 := M(E(z_1, z_2, z_3)) = K^3, \quad N_0 := N(E(z_1, z_2, z_3)) = H^3. \tag{5.25}$$

The equations

$$H(H^3 + 8K^3) = 9C_{12}, \quad K(K^3 - H^3) = 27\mathfrak{C}_{12}$$

give that

$$N_0(N_0 + 8M_0)^3 = 9^3 C_{12}^3, \quad M_0(M_0 - N_0)^3 = 27^3\mathfrak{C}_{12}^3. \tag{5.26}$$

Let

(5.27)

$$M_i := M(ED^i(z_1, z_2, z_3)) = (\psi - 3\omega^{2i}\varphi)^3, \quad N_i := N(ED^i(z_1, z_2, z_3)) = (\psi + 6\omega^{2i}\varphi)^3.$$

We find that

$$M_\infty M_0 M_1 M_2 = -81\sqrt{-3}\mathfrak{C}_{12}^3, \quad N_\infty N_0 N_1 N_2 = 3\sqrt{-3}C_{12}^3. \tag{5.28}$$

Note that

$$M_\infty(D(z_1, z_2, z_3)) = M_\infty(z_1, z_2, z_3), \quad N_\infty(D(z_1, z_2, z_3)) = N_\infty(z_1, z_2, z_3). \tag{5.29}$$

$$M_1(E(z_1, z_2, z_3)) = M_2(z_1, z_2, z_3), \quad N_1(E(z_1, z_2, z_3)) = N_2(z_1, z_2, z_3). \tag{5.30}$$

$$M_2(E(z_1, z_2, z_3)) = -M_1(z_1, z_2, z_3), \quad N_2(E(z_1, z_2, z_3)) = -N_1(z_1, z_2, z_3). \tag{5.31}$$

$$M_\infty(E(z_1, z_2, z_3)) = M_0(z_1, z_2, z_3), \quad N_\infty(E(z_1, z_2, z_3)) = N_0(z_1, z_2, z_3). \tag{5.32}$$

$$M_0(E(z_1, z_2, z_3)) = -M_\infty(z_1, z_2, z_3), \quad N_0(E(z_1, z_2, z_3)) = -N_\infty(z_1, z_2, z_3). \tag{5.33}$$

Let

$$L(z_1, z_2, z_3) := (z_1 - z_2)^3 (z_2 - z_3)^3 (z_3 - z_1)^3. \tag{5.34}$$

Note that

$$AE = EC, \quad CE = EA^2, \quad AC = \omega CA. \tag{5.35}$$

Put

$$L_{i,j,k}(z_1, z_2, z_3) := L(E^i A^j C^k (z_1, z_2, z_3)). \tag{5.36}$$

We find that

$$\begin{cases} L_{0,0,0} = L_{0,1,0} = L_{0,2,0} = L_{2,0,0} = L_{2,1,0} = L_{2,2,0}, \\ L_{0,0,1} = L_{0,1,1} = L_{0,2,1} = L_{2,0,1} = L_{2,1,1} = L_{2,2,1}, \\ L_{0,0,2} = L_{0,1,2} = L_{0,2,2} = L_{2,0,2} = L_{2,1,2} = L_{2,2,2}, \end{cases} \tag{5.37}$$

and

$$\begin{cases} L_{1,0,0} = L_{1,0,1} = L_{1,0,2}, \\ L_{1,1,0} = L_{1,1,1} = L_{1,1,2}, \\ L_{1,2,0} = L_{1,2,1} = L_{1,2,2}. \end{cases} \tag{5.38}$$

Here,

$$\begin{cases} L_{1,0,0} = L(E(z_1, z_2, z_3)) = (z_2^3 - z_3^3)^3, \\ L_{1,1,0} = L(EA(z_1, z_2, z_3)) = (z_3^3 - z_1^3)^3, \\ L_{1,2,0} = L(EA^2(z_1, z_2, z_3)) = (z_1^3 - z_2^3)^3, \end{cases} \tag{5.39}$$

and

$$\begin{cases} L_{0,0,0} = L(z_1, z_2, z_3) = (z_1 - z_2)^3 (z_2 - z_3)^3 (z_3 - z_1)^3, \\ L_{0,0,1} = L(C(z_1, z_2, z_3)) = (z_1 - \omega z_2)^3 (\omega z_2 - \omega^2 z_3)^3 (\omega^2 z_3 - z_1)^3, \\ L_{0,0,2} = L(C^2(z_1, z_2, z_3)) = (z_1 - \omega^2 z_2)^3 (\omega^2 z_2 - \omega z_3)^3 (\omega z_3 - z_1)^3. \end{cases} \tag{5.40}$$

We find that

$$L_{1,0,0} L_{1,1,0} L_{1,2,0} = L_{0,0,0} L_{0,0,1} L_{0,0,2} = C_9^3. \tag{5.41}$$

In fact,

$$\begin{cases} L_{1,0,0}(E(z_1, z_2, z_3)) = L_{0,0,0}(z_1, z_2, z_3), \\ L_{1,0,0}(A(z_1, z_2, z_3)) = L_{1,1,0}(z_1, z_2, z_3), \\ L_{1,0,0}(C(z_1, z_2, z_3)) = L_{1,0,0}(z_1, z_2, z_3). \end{cases}$$

$$\begin{cases} L_{1,1,0}(E(z_1, z_2, z_3)) = L_{0,0,1}(z_1, z_2, z_3), \\ L_{1,1,0}(A(z_1, z_2, z_3)) = L_{1,2,0}(z_1, z_2, z_3), \\ L_{1,1,0}(C(z_1, z_2, z_3)) = L_{1,1,0}(z_1, z_2, z_3). \end{cases}$$

$$\begin{cases} L_{1,2,0}(E(z_1, z_2, z_3)) = L_{0,0,2}(z_1, z_2, z_3), \\ L_{1,2,0}(A(z_1, z_2, z_3)) = L_{1,0,0}(z_1, z_2, z_3), \\ L_{1,2,0}(C(z_1, z_2, z_3)) = L_{1,2,0}(z_1, z_2, z_3). \end{cases}$$

$$\begin{cases} L_{0,0,0}(E(z_1, z_2, z_3)) = L_{1,0,0}(z_1, z_2, z_3), \\ L_{0,0,0}(A(z_1, z_2, z_3)) = L_{0,0,0}(z_1, z_2, z_3), \\ L_{0,0,0}(C(z_1, z_2, z_3)) = L_{0,0,1}(z_1, z_2, z_3). \end{cases}$$

$$\begin{cases} L_{0,0,1}(E(z_1, z_2, z_3)) = L_{1,2,0}(z_1, z_2, z_3), \\ L_{0,0,1}(A(z_1, z_2, z_3)) = L_{0,0,1}(z_1, z_2, z_3), \\ L_{0,0,1}(C(z_1, z_2, z_3)) = L_{0,0,2}(z_1, z_2, z_3). \end{cases}$$

$$\begin{cases} L_{0,0,2}(E(z_1, z_2, z_3)) = L_{1,1,0}(z_1, z_2, z_3), \\ L_{0,0,2}(A(z_1, z_2, z_3)) = L_{0,0,2}(z_1, z_2, z_3), \\ L_{0,0,2}(C(z_1, z_2, z_3)) = L_{0,0,0}(z_1, z_2, z_3). \end{cases}$$

Therefore,

$$\begin{cases} L_{1,0,0}(E(z_1, z_2, z_3)) = L_{0,0,0}(z_1, z_2, z_3), \\ L_{0,0,0}(E(z_1, z_2, z_3)) = L_{1,0,0}(z_1, z_2, z_3). \end{cases} \tag{5.42}$$

$$\begin{cases} L_{1,1,0}(E(z_1, z_2, z_3)) = L_{0,0,1}(z_1, z_2, z_3), \\ L_{0,0,1}(E(z_1, z_2, z_3)) = L_{1,2,0}(z_1, z_2, z_3), \\ L_{1,2,0}(E(z_1, z_2, z_3)) = L_{0,0,2}(z_1, z_2, z_3), \\ L_{0,0,2}(E(z_1, z_2, z_3)) = L_{1,1,0}(z_1, z_2, z_3). \end{cases} \tag{5.43}$$

$$\begin{cases} L_{1,0,0}(A(z_1, z_2, z_3)) = L_{1,1,0}(z_1, z_2, z_3), \\ L_{1,1,0}(A(z_1, z_2, z_3)) = L_{1,2,0}(z_1, z_2, z_3), \\ L_{1,2,0}(A(z_1, z_2, z_3)) = L_{1,0,0}(z_1, z_2, z_3). \end{cases} \tag{5.44}$$

$$\begin{cases} L_{0,0,0}(A(z_1, z_2, z_3)) = L_{0,0,0}(z_1, z_2, z_3), \\ L_{0,0,1}(A(z_1, z_2, z_3)) = L_{0,0,1}(z_1, z_2, z_3), \\ L_{0,0,2}(A(z_1, z_2, z_3)) = L_{0,0,2}(z_1, z_2, z_3). \end{cases} \tag{5.45}$$

$$\begin{cases} L_{1,0,0}(C(z_1, z_2, z_3)) = L_{1,0,0}(z_1, z_2, z_3), \\ L_{1,1,0}(C(z_1, z_2, z_3)) = L_{1,1,0}(z_1, z_2, z_3), \\ L_{1,2,0}(C(z_1, z_2, z_3)) = L_{1,2,0}(z_1, z_2, z_3). \end{cases} \tag{5.46}$$

$$\begin{cases} L_{0,0,0}(C(z_1, z_2, z_3)) = L_{0,0,1}(z_1, z_2, z_3), \\ L_{0,0,1}(C(z_1, z_2, z_3)) = L_{0,0,2}(z_1, z_2, z_3), \\ L_{0,0,2}(C(z_1, z_2, z_3)) = L_{0,0,0}(z_1, z_2, z_3). \end{cases} \tag{5.47}$$

Put

$$\zeta_1 = \frac{M_0}{C_9}, \quad \zeta_2 = \frac{N_0}{C_9}. \tag{5.48}$$

THEOREM 5.6 (MAIN THEOREM 8). *The functions ζ_1 and ζ_2 satisfy the following equations*:

$$\begin{cases} Z_1^2 \zeta_2 (\zeta_2 + 8\zeta_1)^3 = 2^8 \cdot 3^9 Z_2^3, \\ Z_1^2 \zeta_1 (\zeta_1 - \zeta_2)^3 = 27^3 [27(Z_1 + Z_2 - 1)^2 - 4Z_2^3]. \end{cases} \tag{5.49}$$

Proof. We find that

$$\zeta_2(\zeta_2 + 8\zeta_1)^3 = 9^3 \frac{C_{12}^3}{C_9^4} = 9^3 \frac{(\frac{C_{12}}{C_6^2})^3}{(\frac{C_9^2}{C_6^3})^2} = 9^3 \frac{(\frac{Z_2}{3})^3}{(\frac{Z_1}{432})^2} = 2^8 \cdot 3^9 \frac{Z_2^3}{Z_1^2},$$

and

$$\zeta_1(\zeta_1-\zeta_2)^3 = 27^3\frac{\mathfrak{C}_{12}^3}{C_9^4} = 27^3\frac{(\frac{\mathfrak{C}_{12}}{C_6^2})^3}{(\frac{C_9^2}{C_6^3})^2} = 27^3\cdot\frac{1}{1728}\left[\frac{1}{4}(Z_1+Z_2-1)^2-\frac{1}{27}Z_2^3\right]\frac{1}{(\frac{Z_1}{432})^2}.$$

Thus,

$$Z_1^2\zeta_2(\zeta_2+8\zeta_1)^3 = 2^8\cdot 3^9 Z_2^3,$$

and

$$Z_1^2\zeta_1(\zeta_1-\zeta_2)^3 = 27^3[27(Z_1+Z_2-1)^2-4Z_2^3].$$

□

Put

$$\xi_1 := \frac{M_0^{\frac{1}{3}}}{C_6^{\frac{1}{2}}}, \quad \xi_2 := \frac{N_0^{\frac{1}{3}}}{C_6^{\frac{1}{2}}}. \tag{5.50}$$

THEOREM 5.7 (MAIN THEOREM 9). *The functions ξ_1 and ξ_2 satisfy the following equations*:

$$\begin{cases} \xi_2^3(\xi_2^3+8\xi_1^3)^3 = 27Z_2^3, \\ 256\xi_1^3(\xi_1^3-\xi_2^3)^3 = 27[27(Z_1+Z_2-1)^2-4Z_2^3]. \end{cases} \tag{5.51}$$

Proof. We find that

$$\xi_2^3(\xi_2^3+8\xi_1^3)^3 = 9^3\left(\frac{C_{12}}{C_6^2}\right)^3 = 27Z_2^3,$$

and

$$\xi_1^3(\xi_1^3-\xi_2^3)^3 = 27^3\left(\frac{\mathfrak{C}_{12}}{C_6^2}\right)^3 = \frac{27}{256}[27(Z_1+Z_2-1)^2-4Z_2^3].$$

□

Theorem 5.7 gives that

$$Z_1 = \frac{2}{27}[\xi_2^3(\xi_2^3+8\xi_1^3)^3+64\xi_1^3(\xi_1^3-\xi_2^3)^3]^{\frac{1}{2}}-\frac{1}{3}\xi_2(\xi_2^3+8\xi_1^3)+1. \tag{5.52}$$

Note that

$$\zeta_1^2 = \frac{\frac{M_0^2}{C_6^3}}{\frac{C_9^2}{C_6^3}} = \frac{432\xi_1^6}{Z_1}, \quad \zeta_2^2 = \frac{\frac{N_0^2}{C_6^3}}{\frac{C_9^2}{C_6^3}} = \frac{432\xi_2^6}{Z_1}. \tag{5.53}$$

Hence, ξ_1, ξ_2 and ζ_1, ζ_2 satisfy the following relations:

$$\begin{cases} \dfrac{4}{729}\zeta_1^4[\xi_2^3(\xi_2^3+8\xi_1^3)^3+64\xi_1^3(\xi_1^3-\xi_2^3)^3]=[432\xi_1^6+\dfrac{1}{3}\zeta_1^2\xi_2(\xi_2^3+8\xi_1^3)-\zeta_1^2]^2, \\ \dfrac{4}{729}\zeta_2^4[\xi_2^3(\xi_2^3+8\xi_1^3)^3+64\xi_1^3(\xi_1^3-\xi_2^3)^3]=[432\xi_2^6+\dfrac{1}{3}\zeta_2^2\xi_2(\xi_2^3+8\xi_1^3)-\zeta_2^2]^2. \end{cases} \tag{5.54}$$

THEOREM 5.8 (MAIN THEOREM 10). *The function-theoretic resolvents* (*analytic resolvents*) *of* Z_1 *and* Z_2 *are also given by*:

$$\begin{aligned} &Z_1^2 : Z_2^3 : (Z_1+Z_2-1)^2 : [27(Z_1+Z_2-1)^2-4Z_2^3] : 1 \\ =&[2(\xi_2^3(\xi_2^3+8\xi_1^3)^3+64\xi_1^3(\xi_1^3-\xi_2^3)^3)^{\frac{1}{2}}-9\xi_2(\xi_2^3+8\xi_1^3)+27]^2 \\ &:27\xi_2^3(\xi_2^3+8\xi_1^3)^3 : 4[\xi_2^3(\xi_2^3+8\xi_1^3)^3+64\xi_1^3(\xi_1^3-\xi_2^3)^3] \\ &:6912\xi_1^3(\xi_1^3-\xi_2^3)^3 : 729. \end{aligned} \tag{5.55}$$

Proof. It is obtained by Theorem 5.7 and (5.52). □

Note that $CD = DC$, we have

$$C^i D^j(z_1,z_2,z_3)=(z_1,\omega^{i+j}z_2,\omega^{2i+j}z_3). \tag{5.56}$$

We find that

$$\begin{cases} C_6(C^iD^j(z_1,z_2,z_3))=C_6(z_1,z_2,z_3), \\ C_9(C^iD^j(z_1,z_2,z_3))=C_9(z_1,z_2,z_3), \\ C_{12}(C^iD^j(z_1,z_2,z_3))=C_{12}(z_1,z_2,z_3), \\ \mathfrak{C}_{12}(C^iD^j(z_1,z_2,z_3))=\omega^{2j}\mathfrak{C}_{12}(z_1,z_2,z_3). \end{cases} \tag{5.57}$$

Let

$$\begin{cases} G_0 := G(z_1,z_2,z_3)=(z_1-z_2)(z_2-z_3)(z_3-z_1), \\ G_1 := G(C(z_1,z_2,z_3))=(z_1-\omega z_2)(\omega z_2-\omega^2 z_3)(\omega^2 z_3-z_1), \\ G_2 := G(C^2(z_1,z_2,z_3))=(z_1-\omega^2 z_2)(\omega^2 z_2-\omega z_3)(\omega z_3-z_1). \end{cases} \tag{5.58}$$

A straightforward calculation gives that

$$G_0G_1G_2=C_9. \tag{5.59}$$

Hence,

$$u_0u_1u_2 := \frac{C_6G_0}{C_9}\cdot\frac{C_6G_1}{C_9}\cdot\frac{C_6G_2}{C_9}=\frac{C_6^3}{C_9^2}=\frac{432}{Z_1}. \tag{5.60}$$

On the other hand,

$$H(D^j(z_1, z_2, z_3)) = \psi + 6\omega^{2j}\varphi. \tag{5.61}$$

Put

$$H_0 := \psi + 6\varphi, \quad H_1 := \psi + 6\omega^2\varphi, \quad H_2 := \psi + 6\omega\varphi. \tag{5.62}$$

Then

$$H_0 H_1 H_2 = \psi^3 + 216\varphi^3 = \frac{C_{12}}{\psi}. \tag{5.63}$$

Hence,

$$v_0 v_1 v_2 := \frac{C_9 H_0}{C_{12}} \cdot \frac{C_9 H_1}{C_{12}} \cdot \frac{C_9 H_2}{C_{12}} = \frac{C_9^3}{C_{12}^2 \psi}. \tag{5.64}$$

Similarly,

$$K(D^j(z_1, z_2, z_3)) = \psi - 3\omega^{2j}\varphi. \tag{5.65}$$

Put

$$K_0 := \psi - 3\varphi, \quad K_1 := \psi - 3\omega^2\varphi, \quad K_2 := \psi - 3\omega\varphi. \tag{5.66}$$

Then

$$K_0 K_1 K_2 = \psi^3 - 27\varphi^3 = -\frac{\mathfrak{C}_{12}}{\varphi}. \tag{5.67}$$

Hence,

$$w_0 w_1 w_2 := \frac{C_9 K_0}{\mathfrak{C}_{12}} \cdot \frac{C_9 K_1}{\omega^2 \mathfrak{C}_{12}} \cdot \frac{C_9 K_2}{\omega \mathfrak{C}_{12}} = -\frac{C_9^3}{\mathfrak{C}_{12}^2 \varphi}. \tag{5.68}$$

We have

$$\sqrt[3]{M_\infty} = -3\sqrt{-3}\varphi, \quad \sqrt[3]{M_0} = \psi - 3\varphi, \quad \sqrt[3]{M_1} = \psi - 3\omega^2\varphi, \quad \sqrt[3]{M_2} = \psi - 3\omega\varphi. \tag{5.69}$$

Hence,

$$\begin{cases} \sqrt[3]{M_0} + \sqrt[3]{M_1} + \sqrt[3]{M_2} = 3\psi, \\ \sqrt{-3}\,\sqrt[3]{M_\infty} + \sqrt[3]{M_0} + \omega\,\sqrt[3]{M_1} + \omega^2\,\sqrt[3]{M_2} = 0, \\ \sqrt[3]{M_0} + \omega^2\,\sqrt[3]{M_1} + \omega\,\sqrt[3]{M_2} = 0. \end{cases} \tag{5.70}$$

Similarly,

$$\sqrt[3]{N_\infty}=-\sqrt{-3}\psi, \quad \sqrt[3]{N_0}=\psi+6\varphi, \quad \sqrt[3]{N_1}=\psi+6\omega^2\varphi, \quad \sqrt[3]{N_2}=\psi+6\omega\varphi. \tag{5.71}$$

Hence,

$$\begin{cases} -\sqrt{-3}\sqrt[3]{N_\infty}+\sqrt[3]{N_0}+\sqrt[3]{N_1}+\sqrt[3]{N_2}=0, \\ \sqrt[3]{N_0}+\omega\sqrt[3]{N_1}+\omega^2\sqrt[3]{N_2}=18\varphi, \\ \sqrt[3]{N_0}+\omega^2\sqrt[3]{N_1}+\omega\sqrt[3]{N_2}=0. \end{cases} \tag{5.72}$$

6. Ternary cubic forms associated to Hessian polyhedra

Put

$$X(z_1,z_2,z_3)=z_1^3, \quad Y(z_1,z_2,z_3)=z_2^3, \quad Z(z_1,z_2,z_3)=z_3^3. \tag{6.1}$$

$$Q_1(z_1,z_2,z_3)=z_1z_2^2+z_2z_3^2+z_3z_1^2, \quad Q_2(z_1,z_2,z_3)=z_1^2z_2+z_2^2z_3+z_3^2z_1. \tag{6.2}$$

Then

$$\sqrt[3]{L_{1,0,0}}=Y-Z, \quad \sqrt[3]{L_{1,1,0}}=Z-X, \quad \sqrt[3]{L_{1,2,0}}=X-Y. \tag{6.3}$$

$$\sqrt[3]{L_{0,0,0}}=Q_1-Q_2, \quad \sqrt[3]{L_{0,0,1}}=\omega^2Q_1-\omega Q_2, \quad \sqrt[3]{L_{0,0,2}}=\omega Q_1-\omega^2Q_2. \tag{6.4}$$

The functions φ, ψ, χ, X, Y, Z and Q_1, Q_2 satisfy the relations:

$$\begin{cases} \varphi^3=XYZ, \\ \psi=X+Y+Z, \\ \chi=XY+YZ+ZX, \\ \chi+3\varphi^2+\varphi\psi=Q_1Q_2, \\ \psi\chi+6\varphi\chi+6\varphi^2\psi+9\varphi^3=Q_1^3+Q_2^3, \\ (X-Y)(Y-Z)(Z-X)=Q_1^3-Q_2^3. \end{cases} \tag{6.5}$$

The invariants C_6, C_9, C_{12}, $\mathfrak{C}_{12}$ and C_{18} can be expressed as the functions of X, Y and Z:

$$\begin{cases} C_6=(X+Y+Z)^2-12(XY+YZ+ZX), \\ C_9=(X-Y)(Y-Z)(Z-X), \\ C_{12}=(X+Y+Z)[(X+Y+Z)^3+216XYZ], \\ \mathfrak{C}_{12}=\sqrt[3]{XYZ}[27XYZ-(X+Y+Z)^3], \\ C_{18}=(X+Y+Z)^6-540XYZ(X+Y+Z)^3-5832X^2Y^2Z^2. \end{cases} \tag{6.6}$$

Let

$$\begin{cases} W_2 = (X+Y+Z)^2 - 12(XY+YZ+ZX), \\ W_3 = (X-Y)(Y-Z)(Z-X), \\ \mathfrak{W}_3 = XYZ - \varphi^3, \\ W_4 = (X+Y+Z)[(X+Y+Z)^3 + 216\varphi^3], \\ \mathfrak{W}_4 = \varphi[27\varphi^3 - (X+Y+Z)^3], \\ W_6 = (X+Y+Z)^6 - 540\varphi^3(X+Y+Z)^3 - 5832\varphi^6. \end{cases} \tag{6.7}$$

We will regard (φ, X, Y, Z) as a point in the complex projective space $\mathbb{CP}^3$.

We find that

$$1728\mathfrak{W}_4^3 = W_6^2 - W_4^3 \tag{6.8}$$

and

$$\frac{\partial(\mathfrak{W}_3, W_4, \mathfrak{W}_4, W_6)}{\partial(\varphi, X, Y, Z)} = 0. \tag{6.9}$$

For the ternary cubic form

$$F_1(x_1, x_2, x_3) = Xx_1^3 + Yx_2^3 + Zx_3^3 - 3\varphi x_1x_2x_3, \tag{6.10}$$

we have

$$\begin{cases} \sqrt[3]{M_\infty} = -3\sqrt{-3}\varphi, \\ \sqrt[3]{M_0} = -3\varphi + X + Y + Z, \\ \sqrt[3]{M_1} = -3\omega^2\varphi + X + Y + Z, \\ \sqrt[3]{M_2} = -3\omega\varphi + X + Y + Z. \end{cases} \tag{6.11}$$

Correspondingly,

$$\begin{cases} \sqrt[3]{M_\infty}: & (\sqrt{-3}, 0, 0, 0) \in \mathbb{CP}^3, \\ \sqrt[3]{M_0}: & (1, 1, 1, 1) \in \mathbb{CP}^3, \\ \sqrt[3]{M_1}: & (\omega^2, 1, 1, 1) \in \mathbb{CP}^3, \\ \sqrt[3]{M_2}: & (\omega, 1, 1, 1) \in \mathbb{CP}^3. \end{cases} \tag{6.12}$$

For the ternary cubic form

$$F_2(x_1, x_2, x_3) = Xx_1^3 + Yx_2^3 + Zx_3^3 + 6\varphi x_1x_2x_3, \tag{6.13}$$

we have

$$
(6.14) \qquad \begin{cases} \sqrt[3]{N_\infty} = -\sqrt{-3}(X+Y+Z), \\ \sqrt[3]{N_0} = 6\varphi + X+Y+Z, \\ \sqrt[3]{N_1} = 6\omega^2\varphi + X+Y+Z, \\ \sqrt[3]{N_2} = 6\omega\varphi + X+Y+Z. \end{cases}
$$

Correspondingly,

$$
(6.15) \qquad \begin{cases} \sqrt[3]{N_\infty}: & (0, -\sqrt{-3}, -\sqrt{-3}, -\sqrt{-3}) \in \mathbb{CP}^3, \\ \sqrt[3]{N_0}: & (1,1,1,1) \in \mathbb{CP}^3, \\ \sqrt[3]{N_1}: & (\omega^2,1,1,1) \in \mathbb{CP}^3, \\ \sqrt[3]{N_2}: & (\omega,1,1,1) \in \mathbb{CP}^3. \end{cases}
$$

In fact, $W_2 = 0$ is a quadratic ruled surface, $W_3 = 0$ is a cubic ruled surface, $\mathfrak{W}_3 = 0$ is a cubic surface, $W_4 = 0$ is a quartic surface, $\mathfrak{W}_4 = 0$ is a quartic surface, $W_6 = 0$ is a surface of degree six.

The action of Hessian groups on $\mathbb{CP}^3$ has two orbits:

$$\{(\sqrt{-3},0,0,0),(1,1,1,1),(\omega^2,1,1,1),(\omega,1,1,1)\}$$

and

$$\{(0,-\sqrt{-3},-\sqrt{-3},-\sqrt{-3}),(1,1,1,1),(\omega^2,1,1,1),(\omega,1,1,1)\}.$$

Each orbit has four points. The invariant surface $\mathfrak{W}_3 = 0$ passes through the points $(1,1,1,1)$, $(\omega^2,1,1,1)$ and $(\omega,1,1,1)$. The invariant surface $W_2 = 0$ passes through the point $(\sqrt{-3},0,0,0)$. The invariant surface $W_3 = 0$ passes through all the five points $(\sqrt{-3},0,0,0)$, $(1,1,1,1)$, $(\omega^2,1,1,1)$, $(\omega,1,1,1)$ and $(0,-\sqrt{-3},-\sqrt{-3},-\sqrt{-3})$. The invariant surface $W_4 = 0$ passes through the point $(\sqrt{-3},0,0,0)$. The invariant surface $\mathfrak{W}_4 = 0$ passes through the points $(0,-\sqrt{-3},-\sqrt{-3},-\sqrt{-3})$, $(1,1,1,1)$, $(\omega^2,1,1,1)$ and $(\omega,1,1,1)$. The invariant surface $W_6 = 0$ does not pass through the five points.

When we restrict the action of Hessian groups to the cubic surface $\mathfrak{W}_3 = 0$, we get the action of Hessian groups on $\mathbb{CP}^2$. We set

$$
(6.16) \qquad \begin{cases} W_2 = (X+Y+Z)^2 - 12(XY+YZ+ZX), \\ W_3 = (X-Y)(Y-Z)(Z-X), \\ W_4 = (X+Y+Z)[(X+Y+Z)^3 + 216XYZ], \\ W_6 = (X+Y+Z)^6 - 540XYZ(X+Y+Z)^3 - 5832X^2Y^2Z^2, \\ W_{12} = XYZ[27XYZ - (X+Y+Z)^3]^3. \end{cases}
$$

The invariants W_2, W_3, W_4, W_6 and W_{12} satisfy the relations:

$$\begin{cases} 432W_3^2 = W_2^3 - 3W_2W_4 + 2W_6, \\ 1728W_{12} = W_6^2 - W_4^3. \end{cases} \tag{6.17}$$

We will study the algebraic curves in $\mathbb{CP}^2$:

$$W_2 = 0, \quad W_3 = 0, \quad W_4 = 0, \quad W_6 = 0, \quad W_{12} = 0.$$

$W_2 = 0$ is a conic. $W_3 = 0$ is the product of three lines: $X - Y = 0$, $Y - Z = 0$ or $Z - X = 0$. $W_4 = 0$, i.e., $X + Y + Z = 0$ or $(X + Y + Z)^3 + 216XYZ = 0$. $W_6 = 0$, i.e., $(X + Y + Z)^3 - (270 + 162\sqrt{3})XYZ = 0$ or $(X + Y + Z)^3 - (270 - 162\sqrt{3})XYZ = 0$. $W_{12} = 0$, i.e., $X = 0$, $Y = 0$, $Z = 0$ or $(X + Y + Z)^3 - 27XYZ = 0$.

The curves $W_2 = 0$, $W_4 = 0$, $W_6 = 0$, $W_{12} = 0$ intersect $W_3 = 0$ in 6, 12, 18, 15 points respectively.

$$\{W_2 = 0\} \cap \{W_3 = 0\} = \{(\alpha_i, 1, 1), (1, \alpha_i, 1), (1, 1, \alpha_i), i = 1, 2\},$$

where α_1 and α_2 are two roots of the quadratic equation:

$$\alpha^2 - 20\alpha - 8 = 0.$$

$$\begin{aligned} &\{W_4 = 0\} \cap \{W_3 = 0\} \\ =&\{(-2, 1, 1), (1, -2, 1), (1, 1, -2), (\beta_i, 1, 1), (1, \beta_i, 1), (1, 1, \beta_i), i = 1, 2, 3\}, \end{aligned}$$

where β_1, β_2 and β_3 are the roots of the cubic equation:

$$\beta^3 + 6\beta^2 + 228\beta + 8 = 0.$$

$$\{W_6 = 0\} \cap \{W_3 = 0\} = \{(\gamma_i, 1, 1), (1, \gamma_i, 1), (1, 1, \gamma_i), 1 \leq i \leq 6\},$$

where γ_i $(1 \leq i \leq 6)$ are the roots of the equation of the sixth degree:

$$[\gamma^3 + 6\gamma^2 - (258 + 162\sqrt{3})\gamma + 8][\gamma^3 + 6\gamma^2 - (258 - 162\sqrt{3})\gamma + 8] = 0.$$

$$\begin{aligned} \{W_{12} = 0\} \cap \{W_3 = 0\} = \{&(1, 0, 0), (0, 1, 0), (0, 0, 1), (1, 1, 0), (1, 0, 1), (0, 1, 1), \\ &(\delta_i, 1, 1), (1, \delta_i, 1), (1, 1, \delta_i), i = 1, 2, 3\}, \end{aligned}$$

where δ_1, δ_2 and δ_3 are the roots of the cubic equation:

$$\delta^3 + 6\delta^2 - 15\delta + 8 = 0.$$

In fact, by $W_3 = 0$, we have

$$(X, Y, Z) = (X, X, Z), (X, Y, Y), (Z, Y, Z).$$

Without loss of generality, for $(X, Y, Z) = (X, Y, Y)$, we find that

$$\begin{cases} W_2 = X^2 - 20XY - 8Y^2, \\ W_4 = (X + 2Y)(X^3 + 6X^2Y + 228XY^2 + 8Y^3), \\ W_6 = [X^3 + 6X^2Y - (258 + 162\sqrt{3})XY^2 + 8Y^3] \times \\ \qquad \times [X^3 + 6X^2Y - (258 - 162\sqrt{3})XY^2 + 8Y^3], \\ W_{12} = -XY^2(X^3 + 6X^2Y - 15XY^2 + 8Y^3)^3. \end{cases}$$

Note that

$$\begin{aligned} &\mathbb{C}[W_2, W_3, W_4, W_6, W_{12}]/(W_2^3 - 3W_2W_4 - 432W_3^2 + 2W_6, W_6^2 - W_4^3 - 1728W_{12}) \\ \cong &\mathbb{C}[W_2, W_3, W_4]. \end{aligned} \tag{6.18}$$

We find that

$$\begin{cases} \sqrt[3]{L_{1,0,0}} + \sqrt[3]{L_{1,1,0}} + \sqrt[3]{L_{1,2,0}} = 0, \\ \sqrt[3]{L_{0,0,0}} + \sqrt[3]{L_{0,0,1}} + \sqrt[3]{L_{0,0,2}} = 0. \end{cases} \tag{6.19}$$

Moreover,

$$\begin{cases} L_{1,0,0} + L_{1,1,0} + L_{1,2,0} = 3C_9, \\ L_{0,0,0} + L_{0,0,1} + L_{0,0,2} = 3C_9. \end{cases} \tag{6.20}$$

In fact,

$$\sqrt{-3}(M_0 + M_1 + M_2) = N_\infty - M_\infty, \quad \sqrt{-3}(N_0 + N_1 + N_2) = N_\infty + 8M_\infty. \tag{6.21}$$

$$\begin{aligned} (L_{1,0,0} + L_{1,1,0} + L_{1,2,0})^3 &= 27L_{1,0,0}L_{1,1,0}L_{1,2,0}, \\ (L_{0,0,0} + L_{0,0,1} + L_{0,0,2})^3 &= 27L_{0,0,0}L_{0,0,1}L_{0,0,2}. \end{aligned} \tag{6.22}$$

We have

$$\begin{cases} \varphi(A(z_1, z_2, z_3)) = \varphi(z_1, z_2, z_3), \\ \psi(A(z_1, z_2, z_3)) = \psi(z_1, z_2, z_3), \\ X(A(z_1, z_2, z_3)) = Y(z_1, z_2, z_3), \\ Y(A(z_1, z_2, z_3)) = Z(z_1, z_2, z_3), \\ Z(A(z_1, z_2, z_3)) = X(z_1, z_2, z_3), \\ Q_1(A(z_1, z_2, z_3)) = Q_1(z_1, z_2, z_3), \\ Q_2(A(z_1, z_2, z_3)) = Q_2(z_1, z_2, z_3). \end{cases} \tag{6.23}$$

$$\begin{cases} \varphi(B(z_1,z_2,z_3)) = \varphi(z_1,z_2,z_3), \\ \psi(B(z_1,z_2,z_3)) = \psi(z_1,z_2,z_3), \\ X(B(z_1,z_2,z_3)) = X(z_1,z_2,z_3), \\ Y(B(z_1,z_2,z_3)) = Z(z_1,z_2,z_3), \\ Z(B(z_1,z_2,z_3)) = Y(z_1,z_2,z_3), \\ Q_1(B(z_1,z_2,z_3)) = Q_2(z_1,z_2,z_3), \\ Q_2(B(z_1,z_2,z_3)) = Q_1(z_1,z_2,z_3). \end{cases} \tag{6.24}$$

$$\begin{cases} \varphi(C(z_1,z_2,z_3)) = \varphi(z_1,z_2,z_3), \\ \psi(C(z_1,z_2,z_3)) = \psi(z_1,z_2,z_3), \\ X(C(z_1,z_2,z_3)) = X(z_1,z_2,z_3), \\ Y(C(z_1,z_2,z_3)) = Y(z_1,z_2,z_3), \\ Z(C(z_1,z_2,z_3)) = Z(z_1,z_2,z_3), \\ Q_1(C(z_1,z_2,z_3)) = \omega^2 Q_1(z_1,z_2,z_3), \\ Q_2(C(z_1,z_2,z_3)) = \omega Q_2(z_1,z_2,z_3). \end{cases} \tag{6.25}$$

$$\begin{cases} \varphi(D(z_1,z_2,z_3)) = \omega^2\varphi(z_1,z_2,z_3), \\ \psi(D(z_1,z_2,z_3)) = \psi(z_1,z_2,z_3), \\ X(D(z_1,z_2,z_3)) = X(z_1,z_2,z_3), \\ Y(D(z_1,z_2,z_3)) = Y(z_1,z_2,z_3), \\ Z(D(z_1,z_2,z_3)) = Z(z_1,z_2,z_3). \end{cases} \tag{6.26}$$

$$\begin{cases} (\sqrt{-3})^3\varphi(E(z_1,z_2,z_3)) = \psi - 3\varphi, \\ -\sqrt{-3}\psi(E(z_1,z_2,z_3)) = \psi + 6\varphi, \\ (\sqrt{-3})^3 X(E(z_1,z_2,z_3)) = \psi + 6\varphi + 3Q_1 + 3Q_2, \\ (\sqrt{-3})^3 Y(E(z_1,z_2,z_3)) = \psi + 6\varphi + 3\omega^2 Q_1 + 3\omega Q_2, \\ (\sqrt{-3})^3 Z(E(z_1,z_2,z_3)) = \psi + 6\varphi + 3\omega Q_1 + 3\omega^2 Q_2, \\ -\sqrt{-3}Q_1(E(z_1,z_2,z_3)) = X + \omega^2 Y + \omega Z, \\ -\sqrt{-3}Q_2(E(z_1,z_2,z_3)) = X + \omega Y + \omega^2 Z. \end{cases} \tag{6.27}$$

We have

$$\sum_{\nu=0}^{2} \omega^{\nu} M_{\nu} = -27\psi^2\varphi, \quad \sum_{\nu=0}^{2} \omega^{2\nu} M_{\nu} = 81\psi\varphi^2. \tag{6.28}$$

$$\sum_{\nu=0}^{2}\omega^{\nu}N_{\nu}=54\psi^2\varphi,\quad \sum_{\nu=0}^{2}\omega^{2\nu}N_{\nu}=324\psi\varphi^2. \tag{6.29}$$

$$\sum_{\nu=0}^{2}\omega^{\nu}L_{0,0,\nu}=-9Q_1^2Q_2,\quad \sum_{\nu=0}^{2}\omega^{2\nu}L_{0,0,\nu}=9Q_1Q_2^2. \tag{6.30}$$

Hence,

$$\left(\sum_{\nu=0}^{2}\omega^{\nu}M_{\nu}\right)\cdot\left(\sum_{\nu=0}^{2}\omega^{2\nu}M_{\nu}\right)=3M_{\infty}N_{\infty}. \tag{6.31}$$

$$\left(\sum_{\nu=0}^{2}\omega^{\nu}N_{\nu}\right)\cdot\left(\sum_{\nu=0}^{2}\omega^{2\nu}N_{\nu}\right)=-24M_{\infty}N_{\infty}. \tag{6.32}$$

$$\left(\sum_{\nu=0}^{2}\omega^{\nu}L_{0,0,\nu}\right)\cdot\left(\sum_{\nu=0}^{2}\omega^{2\nu}L_{0,0,\nu}\right)=-81(\chi+3\varphi^2+\varphi\psi)^3. \tag{6.33}$$

In fact,

$$\begin{cases}\sqrt[3]{M_{\infty}}\sqrt[3]{M_0}\sqrt[3]{M_1}\sqrt[3]{M_2}=3\sqrt{-3}\mathfrak{C}_{12},\\ \sqrt[3]{N_{\infty}}\sqrt[3]{N_0}\sqrt[3]{N_1}\sqrt[3]{N_2}=-\sqrt{-3}C_{12},\\ \sqrt[3]{L_{1,0,0}}\sqrt[3]{L_{1,1,0}}\sqrt[3]{L_{1,2,0}}=C_9,\\ \sqrt[3]{L_{0,0,0}}\sqrt[3]{L_{0,0,1}}\sqrt[3]{L_{0,0,2}}=C_9.\end{cases} \tag{6.34}$$

The Hessian of C_6 is given by

$$\text{Hessian}(C_6)=\begin{vmatrix}\frac{\partial^2C_6}{\partial z_1^2} & \frac{\partial^2C_6}{\partial z_1\partial z_2} & \frac{\partial^2C_6}{\partial z_1\partial z_3}\\ \frac{\partial^2C_6}{\partial z_1\partial z_2} & \frac{\partial^2C_6}{\partial z_2^2} & \frac{\partial^2C_6}{\partial z_2\partial z_3}\\ \frac{\partial^2C_6}{\partial z_1\partial z_3} & \frac{\partial^2C_6}{\partial z_2\partial z_3} & \frac{\partial^2C_6}{\partial z_3^2}\end{vmatrix}=-108000\mathfrak{C}_{12}. \tag{6.35}$$

The Hessian of $\mathfrak{C}_{12}$ with respect to φ and ψ:

$$\text{Hessian}(\mathfrak{C}_{12})=\begin{vmatrix}\frac{\partial^2\mathfrak{C}_{12}}{\partial\varphi^2} & \frac{\partial^2\mathfrak{C}_{12}}{\partial\varphi\partial\psi}\\ \frac{\partial^2\mathfrak{C}_{12}}{\partial\varphi\partial\psi} & \frac{\partial^2\mathfrak{C}_{12}}{\partial\psi^2}\end{vmatrix}=-9C_{12}. \tag{6.36}$$

The Hessian of C_{12} with respect to φ and ψ:

$$\text{Hessian}(C_{12}) = \begin{vmatrix} \frac{\partial^2 C_{12}}{\partial \varphi^2} & \frac{\partial^2 C_{12}}{\partial \varphi \partial \psi} \\ \frac{\partial^2 C_{12}}{\partial \varphi \partial \psi} & \frac{\partial^2 C_{12}}{\partial \psi^2} \end{vmatrix} = -15552\mathfrak{C}_{12}. \tag{6.37}$$

The functional determinant of $\mathfrak{C}_{12}$ and C_{12} with respect to φ and ψ:

$$\frac{\partial(\mathfrak{C}_{12}, C_{12})}{\partial(\varphi, \psi)} = \begin{vmatrix} \frac{\partial \mathfrak{C}_{12}}{\partial \varphi} & \frac{\partial \mathfrak{C}_{12}}{\partial \psi} \\ \frac{\partial C_{12}}{\partial \varphi} & \frac{\partial C_{12}}{\partial \psi} \end{vmatrix} = -4C_{18}. \tag{6.38}$$

We have
(6.39)

$$C_{12}^3 : C_{18}^2 : 1728\mathfrak{C}_{12}^3 = \psi^3(\psi^3+216\varphi^3)^3 : (\psi^6-540\varphi^3\psi^3-5832\varphi^6)^2 : 1728\varphi^3(27\varphi^3-\psi^3)^3.$$

Note that

$$H_1 := \text{Hessian}(Q_1) = \begin{vmatrix} \frac{\partial^2 Q_1}{\partial z_1^2} & \frac{\partial^2 Q_1}{\partial z_1 \partial z_2} & \frac{\partial^2 Q_1}{\partial z_1 \partial z_3} \\ \frac{\partial^2 Q_1}{\partial z_1 \partial z_2} & \frac{\partial^2 Q_1}{\partial z_2^2} & \frac{\partial^2 Q_1}{\partial z_2 \partial z_3} \\ \frac{\partial^2 Q_1}{\partial z_1 \partial z_3} & \frac{\partial^2 Q_1}{\partial z_2 \partial z_3} & \frac{\partial^2 Q_1}{\partial z_3^2} \end{vmatrix} = -8(\psi - 3\varphi). \tag{6.40}$$

$$H_2 := \text{Hessian}(Q_2) = \begin{vmatrix} \frac{\partial^2 Q_2}{\partial z_1^2} & \frac{\partial^2 Q_2}{\partial z_1 \partial z_2} & \frac{\partial^2 Q_2}{\partial z_1 \partial z_3} \\ \frac{\partial^2 Q_2}{\partial z_1 \partial z_2} & \frac{\partial^2 Q_2}{\partial z_2^2} & \frac{\partial^2 Q_2}{\partial z_2 \partial z_3} \\ \frac{\partial^2 Q_2}{\partial z_1 \partial z_3} & \frac{\partial^2 Q_2}{\partial z_2 \partial z_3} & \frac{\partial^2 Q_2}{\partial z_3^2} \end{vmatrix} = -8(\psi - 3\varphi). \tag{6.41}$$

$$H_3 := \text{Hessian}(\varphi) = 2\varphi. \tag{6.42}$$

$$H_4 := \text{Hessian}(\psi) = 216\varphi. \tag{6.43}$$

We have

$$J_1 := \begin{vmatrix} \frac{\partial^2 Q_1}{\partial z_1^2} & \frac{\partial^2 Q_1}{\partial z_1 \partial z_2} & \frac{\partial^2 Q_1}{\partial z_1 \partial z_3} & \frac{\partial H_1}{\partial z_1} \\ \frac{\partial^2 Q_1}{\partial z_1 \partial z_2} & \frac{\partial^2 Q_1}{\partial z_2^2} & \frac{\partial^2 Q_1}{\partial z_2 \partial z_3} & \frac{\partial H_1}{\partial z_2} \\ \frac{\partial^2 Q_1}{\partial z_1 \partial z_3} & \frac{\partial^2 Q_1}{\partial z_2 \partial z_3} & \frac{\partial^2 Q_1}{\partial z_3^2} & \frac{\partial H_1}{\partial z_3} \\ \frac{\partial H_1}{\partial z_1} & \frac{\partial H_1}{\partial z_2} & \frac{\partial H_1}{\partial z_3} & 0 \end{vmatrix} = 6912[3Q_1^2 - (\psi + 6\varphi)Q_2]. \tag{6.44}$$

$$J_2 := \begin{vmatrix} \frac{\partial^2 Q_2}{\partial z_1^2} & \frac{\partial^2 Q_2}{\partial z_1 \partial z_2} & \frac{\partial^2 Q_2}{\partial z_1 \partial z_3} & \frac{\partial H_2}{\partial z_1} \\ \frac{\partial^2 Q_2}{\partial z_1 \partial z_2} & \frac{\partial^2 Q_2}{\partial z_2^2} & \frac{\partial^2 Q_2}{\partial z_2 \partial z_3} & \frac{\partial H_2}{\partial z_2} \\ \frac{\partial^2 Q_2}{\partial z_1 \partial z_3} & \frac{\partial^2 Q_2}{\partial z_2 \partial z_3} & \frac{\partial^2 Q_2}{\partial z_3^2} & \frac{\partial H_2}{\partial z_3} \\ \frac{\partial H_2}{\partial z_1} & \frac{\partial H_2}{\partial z_2} & \frac{\partial H_2}{\partial z_3} & 0 \end{vmatrix} = 6912[3Q_2^2 - (\psi + 6\varphi)Q_1]. \tag{6.45}$$

$$J_3 := \begin{vmatrix} \frac{\partial^2 \varphi}{\partial z_1^2} & \frac{\partial^2 \varphi}{\partial z_1 \partial z_2} & \frac{\partial^2 \varphi}{\partial z_1 \partial z_3} & \frac{\partial H_3}{\partial z_1} \\ \frac{\partial^2 \varphi}{\partial z_1 \partial z_2} & \frac{\partial^2 \varphi}{\partial z_2^2} & \frac{\partial^2 \varphi}{\partial z_2 \partial z_3} & \frac{\partial H_3}{\partial z_2} \\ \frac{\partial^2 \varphi}{\partial z_1 \partial z_3} & \frac{\partial^2 \varphi}{\partial z_2 \partial z_3} & \frac{\partial^2 \varphi}{\partial z_3^2} & \frac{\partial H_3}{\partial z_3} \\ \frac{\partial H_3}{\partial z_1} & \frac{\partial H_3}{\partial z_2} & \frac{\partial H_3}{\partial z_3} & 0 \end{vmatrix} = -12\varphi^2. \tag{6.46}$$

$$J_4 := \begin{vmatrix} \frac{\partial^2 \psi}{\partial z_1^2} & \frac{\partial^2 \psi}{\partial z_1 \partial z_2} & \frac{\partial^2 \psi}{\partial z_1 \partial z_3} & \frac{\partial H_4}{\partial z_1} \\ \frac{\partial^2 \psi}{\partial z_1 \partial z_2} & \frac{\partial^2 \psi}{\partial z_2^2} & \frac{\partial^2 \psi}{\partial z_2 \partial z_3} & \frac{\partial H_4}{\partial z_2} \\ \frac{\partial^2 \psi}{\partial z_1 \partial z_3} & \frac{\partial^2 \psi}{\partial z_2 \partial z_3} & \frac{\partial^2 \psi}{\partial z_3^2} & \frac{\partial H_4}{\partial z_3} \\ \frac{\partial H_4}{\partial z_1} & \frac{\partial H_4}{\partial z_2} & \frac{\partial H_4}{\partial z_3} & 0 \end{vmatrix} = -36^4 \chi. \tag{6.47}$$

$$K_1 := \begin{vmatrix} \frac{\partial Q_1}{\partial z_1} & \frac{\partial Q_1}{\partial z_2} & \frac{\partial Q_1}{\partial z_3} \\ \frac{\partial H_1}{\partial z_1} & \frac{\partial H_1}{\partial z_2} & \frac{\partial H_1}{\partial z_3} \\ \frac{\partial J_1}{\partial z_1} & \frac{\partial J_1}{\partial z_2} & \frac{\partial J_1}{\partial z_3} \end{vmatrix} = 6912 \cdot 24[(\psi + 6\varphi)^3 - 27Q_2^3]. \tag{6.48}$$

$$K_2 := \begin{vmatrix} \frac{\partial Q_2}{\partial z_1} & \frac{\partial Q_2}{\partial z_2} & \frac{\partial Q_2}{\partial z_3} \\ \frac{\partial H_2}{\partial z_1} & \frac{\partial H_2}{\partial z_2} & \frac{\partial H_2}{\partial z_3} \\ \frac{\partial J_2}{\partial z_1} & \frac{\partial J_2}{\partial z_2} & \frac{\partial J_2}{\partial z_3} \end{vmatrix} = -6912 \cdot 24[(\psi + 6\varphi)^3 - 27Q_1^3]. \tag{6.49}$$

$$K_3 := \begin{vmatrix} \frac{\partial \varphi}{\partial z_1} & \frac{\partial \varphi}{\partial z_2} & \frac{\partial \varphi}{\partial z_3} \\ \frac{\partial H_3}{\partial z_1} & \frac{\partial H_3}{\partial z_2} & \frac{\partial H_3}{\partial z_3} \\ \frac{\partial J_3}{\partial z_1} & \frac{\partial J_3}{\partial z_2} & \frac{\partial J_3}{\partial z_3} \end{vmatrix} = 0. \tag{6.50}$$

$$K_4 := \begin{vmatrix} \frac{\partial \psi}{\partial z_1} & \frac{\partial \psi}{\partial z_2} & \frac{\partial \psi}{\partial z_3} \\ \frac{\partial H_4}{\partial z_1} & \frac{\partial H_4}{\partial z_2} & \frac{\partial H_4}{\partial z_3} \\ \frac{\partial J_4}{\partial z_1} & \frac{\partial J_4}{\partial z_2} & \frac{\partial J_4}{\partial z_3} \end{vmatrix} = -36^5 \cdot 54(Q_1^3 - Q_2^3). \tag{6.51}$$

$$(G,H,K):=\begin{vmatrix} \frac{\partial G}{\partial z_1} & \frac{\partial G}{\partial z_2} & \frac{\partial G}{\partial z_3} \\ \frac{\partial H}{\partial z_1} & \frac{\partial H}{\partial z_2} & \frac{\partial H}{\partial z_3} \\ \frac{\partial K}{\partial z_1} & \frac{\partial K}{\partial z_2} & \frac{\partial K}{\partial z_3} \end{vmatrix}=27[3(Q_1^2+Q_2^2)-(\psi+6\varphi)(Q_1+Q_2)]. \tag{6.52}$$

Put

$$U_1=z_1z_2^2, \quad U_2=z_2z_3^2, \quad U_3=z_3z_1^2, \tag{6.53}$$

$$V_1=z_1^2z_2, \quad V_2=z_2^2z_3, \quad V_3=z_3^2z_1. \tag{6.54}$$

Then

$$U_1V_1=XY, \quad U_2V_2=YZ, \quad U_3V_3=ZX. \tag{6.55}$$

$$U_1U_2U_3=V_1V_2V_3=XYZ=\varphi^3. \tag{6.56}$$

$$Q_1=U_1+U_2+U_3, \quad Q_2=V_1+V_2+V_3, \quad \psi=X+Y+Z. \tag{6.57}$$

The ternary cubic form

$$\begin{aligned} F(x_1,x_2,x_3)=&Xx_1^3+Yx_2^3+Zx_3^3+6\varphi x_1x_2x_3+\\ &+3(U_1x_1x_2^2+U_2x_2x_3^2+U_3x_3x_1^2)+3(V_1x_1^2x_2+V_2x_2^2x_3+V_3x_3^2x_1)\\ =&(z_1x_1+z_2x_2+z_3x_3)^3. \end{aligned} \tag{6.58}$$

We have

$$\begin{cases} U_1(A(z_1,z_2,z_3))=U_2(z_1,z_2,z_3),\\ U_2(A(z_1,z_2,z_3))=U_3(z_1,z_2,z_3),\\ U_3(A(z_1,z_2,z_3))=U_1(z_1,z_2,z_3). \end{cases} \tag{6.59}$$

$$\begin{cases} V_1(A(z_1,z_2,z_3))=V_2(z_1,z_2,z_3),\\ V_2(A(z_1,z_2,z_3))=V_3(z_1,z_2,z_3),\\ V_3(A(z_1,z_2,z_3))=V_1(z_1,z_2,z_3). \end{cases} \tag{6.60}$$

$$\begin{cases} U_1(B(z_1,z_2,z_3))=V_3(z_1,z_2,z_3),\\ U_2(B(z_1,z_2,z_3))=V_2(z_1,z_2,z_3),\\ U_3(B(z_1,z_2,z_3))=V_1(z_1,z_2,z_3). \end{cases} \tag{6.61}$$

$$\begin{cases} V_1(B(z_1,z_2,z_3)) = U_3(z_1,z_2,z_3), \\ V_2(B(z_1,z_2,z_3)) = U_2(z_1,z_2,z_3), \\ V_3(B(z_1,z_2,z_3)) = U_1(z_1,z_2,z_3). \end{cases} \tag{6.62}$$

$$\begin{cases} U_1(C(z_1,z_2,z_3)) = \omega^2 U_1(z_1,z_2,z_3), \\ U_2(C(z_1,z_2,z_3)) = \omega^2 U_2(z_1,z_2,z_3), \\ U_3(C(z_1,z_2,z_3)) = \omega^2 U_3(z_1,z_2,z_3). \end{cases} \tag{6.63}$$

$$\begin{cases} V_1(C(z_1,z_2,z_3)) = \omega V_1(z_1,z_2,z_3), \\ V_2(C(z_1,z_2,z_3)) = \omega V_2(z_1,z_2,z_3), \\ V_3(C(z_1,z_2,z_3)) = \omega V_3(z_1,z_2,z_3). \end{cases} \tag{6.64}$$

$$\begin{cases} U_1(D(z_1,z_2,z_3)) = \omega^2 U_1(z_1,z_2,z_3), \\ U_2(D(z_1,z_2,z_3)) = U_2(z_1,z_2,z_3), \\ U_3(D(z_1,z_2,z_3)) = \omega U_3(z_1,z_2,z_3). \end{cases} \tag{6.65}$$

$$\begin{cases} V_1(D(z_1,z_2,z_3)) = \omega V_1(z_1,z_2,z_3), \\ V_2(D(z_1,z_2,z_3)) = V_2(z_1,z_2,z_3), \\ V_3(D(z_1,z_2,z_3)) = \omega^2 V_3(z_1,z_2,z_3). \end{cases} \tag{6.66}$$

$$\begin{cases} (\sqrt{-3})^3 U_1(E(z_1,z_2,z_3)) = X + \omega^2 Y + \omega Z + (\omega - 1)U_1 + (1 - \omega^2)U_2 + \\ \qquad + (\omega^2 - \omega)U_3 + (\omega - \omega^2)V_1 + (1 - \omega)V_2 + (\omega^2 - 1)V_3, \\ (\sqrt{-3})^3 U_2(E(z_1,z_2,z_3)) = X + \omega^2 Y + \omega Z + (1 - \omega^2)U_1 + (\omega^2 - \omega)U_2 + \\ \qquad + (\omega - 1)U_3 + (\omega^2 - 1)V_1 + (\omega - \omega^2)V_2 + (1 - \omega)V_3, \\ (\sqrt{-3})^3 U_3(E(z_1,z_2,z_3)) = X + \omega^2 Y + \omega Z + (\omega^2 - \omega)U_1 + (\omega - 1)U_2 + \\ \qquad + (1 - \omega^2)U_3 + (1 - \omega)V_1 + (\omega^2 - 1)V_2 + (\omega - \omega^2)V_3. \end{cases} \tag{6.67}$$

$$\begin{cases} (\sqrt{-3})^3 V_1(E(z_1,z_2,z_3)) = X + \omega Y + \omega^2 Z + (\omega - \omega^2)U_1 + (\omega^2 - 1)U_2 + \\ \qquad + (1 - \omega)U_3 + (1 - \omega^2)V_1 + (\omega - 1)V_2 + (\omega^2 - \omega)V_3, \\ (\sqrt{-3})^3 V_2(E(z_1,z_2,z_3)) = X + \omega Y + \omega^2 Z + (1 - \omega)U_1 + (\omega - \omega^2)U_2 + \\ \qquad + (\omega^2 - 1)U_3 + (\omega - 1)V_1 + (\omega^2 - \omega)V_2 + (1 - \omega^2)V_3, \\ (\sqrt{-3})^3 V_3(E(z_1,z_2,z_3)) = X + \omega Y + \omega^2 Z + (\omega^2 - 1)U_1 + (1 - \omega)U_2 + \\ \qquad + (\omega - \omega^2)U_3 + (\omega^2 - \omega)V_1 + (1 - \omega^2)V_2 + (\omega - 1)V_3. \end{cases} \tag{6.68}$$

7. Some rational invariants on $\mathbb{CP}^2$

For the Hessian polyhedral equations:

$$\begin{cases} 432\dfrac{C_9^2}{C_6^3} = R_1(x, y), \\ 3\dfrac{C_{12}}{C_6^2} = R_2(x, y), \end{cases} \tag{7.1}$$

there are some particular cases:

$$\begin{cases} 432\dfrac{C_9^2}{C_6^3} = \dfrac{a_1 x + a_2 y + a_3}{c_1 x + c_2 y + c_3}, \\ 3\dfrac{C_{12}}{C_6^2} = \dfrac{b_1 x + b_2 y + b_3}{c_1 x + c_2 y + c_3}, \end{cases} \tag{7.2}$$

where $\begin{pmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{pmatrix} \in GL(3, \mathbb{C})$.

Set

$$x_1 = C_6, \quad x_2 = 12C_9, \quad x_3 = C_{12}, \quad x_4 = 12\mathfrak{C}_{12}, \quad x_5 = C_{18}.$$

Then

$$3x_2^2 = x_1^3 - 3x_1 x_3 + 2x_5, \quad x_4^3 = x_5^2 - x_3^3.$$

Put

$$A = \mathbb{C}[x_1, x_2, x_3, x_4, x_5]/I, \tag{7.3}$$

where the ideal

$$I = (x_1^3 - 3x_1 x_3 - 3x_2^2 + 2x_5, x_3^3 + x_4^3 - x_5^2). \tag{7.4}$$

We find that

$$A \cong \mathbb{C}[x_1, x_2, x_3, x_4]/(x_3^3 + x_4^3 - \frac{1}{4}(x_1^3 - 3x_1 x_3 - 3x_2^2)^2). \tag{7.5}$$

The work of Belyi ([Be]) represents one of the most recent advances in Galois theory, known as the theory of Dessins d'Enfants called by its founder Grothendieck (see [Gro]). The core notion of the theory is that of a Belyi function (see [Be]). In the planar case, a Belyi function is a rational function defined on the Riemann complex sphere $\mathbb{CP}^1$ which has at most three critical values, namely, 0, 1 and ∞. A pre-image of the segment $[1, \infty]$

under such a function is a planar map. The coefficients of such a function are algebraic numbers. This leads to an action of the absolute Galois group on planar maps (see [MZ]).

According to [CG], let us recall a classical result due to Klein concerning the classification of genus zero Galois coverings, and related to the classification of regular polyhedra ([Kl9]). We will introduce a certain number of Galois genus zero coverings of the sphere, corresponding to the well-known dessins.

The first family corresponds to the dessins consisting of a star with e rays where e is a positive integer. A corresponding Belyi function is

$$y = f(x) = x^e$$

where e is the degree of the covering, totally ramified over 0 and ∞ and unramified elsewhere. We call these dessins C_e. Their topological Galois group is the cyclic group with e elements, C_e.

The second family corresponds to the polygon with $2e$ edges and admits the following Belyi function

$$-4y = x^e + x^{-e} - 2.$$

We call these dessins D_{2e}. Their topological Galois group is the dihedral group with $2e$ elements, D_{2e}.

The third family corresponds to the tetrahedron T of degree 12 with Galois group the alternating permutation group on 4 letters A_4. A corresponding Belyi function is given by

$$yx^3(x^3+8)^3 = 2^6(x^3-1)^3.$$

The fourth family corresponds to the octahedron O of degree 24 with Galois group the full symmetric group on 4 letters S_4. A corresponding Belyi function is given by

$$y(x^8+14x^4+1)^3 = 2^2 \cdot 3^3 \cdot x^4(x^4-1)^4.$$

The fifth family corresponds to the icosahedron I of degree 60 with Galois group the alternating permutation group on 5 letters A_5. A corresponding Belyi function is given by

$$y(x^{20}+228x^{15}+494x^{10}-228x^5+1)^3 = x^5(x^{10}-11x^5-1)^5.$$

Let us define three rational functions (invariants) associated to the complex projective plane $\mathbb{CP}^2$:

$$\begin{cases} f_1^{\text{Hessian}}(\xi,\eta) := 432\dfrac{C_9(\xi,\eta)^2}{C_6(\xi,\eta)^3}, \\ f_2^{\text{Hessian}}(\xi,\eta) := 3\dfrac{C_{12}(\xi,\eta)}{C_6(\xi,\eta)^2}, \\ f_3^{\text{Hessian}}(\xi,\eta) := \dfrac{C_{18}(\xi,\eta)^2}{1728\mathfrak{C}_{12}(\xi,\eta)^3}. \end{cases} \tag{7.6}$$

In fact,

$$\left\{\begin{aligned} f_1^{\text{Hessian}}(\xi,\eta) &= 432\frac{(\xi^3-1)^2(\eta^3-1)^2(\xi^3-\eta^3)^2}{(\xi^6-10\xi^3\eta^3+\eta^6-10\xi^3-10\eta^3+1)^3}, \\ f_2^{\text{Hessian}}(\xi,\eta) &= 3\frac{(\xi^3+\eta^3+1)[(\eta^3+\eta^3+1)^3+216\xi^3\eta^3]}{(\xi^6-10\xi^3\eta^3+\eta^6-10\xi^3-10\eta^3+1)^2}, \\ f_3^{\text{Hessian}}(\xi,\eta) &= \frac{[(\xi^3+\eta^3+1)^6-540\xi^3\eta^3(\xi^3+\eta^3+1)^3-5832\xi^6\eta^6]^2}{1728\xi^3\eta^3[27\xi^3\eta^3-(\xi^3+\eta^3+1)^3]^3}. \end{aligned}\right. \tag{7.7}$$

Let

$$\left\{\begin{aligned} F_1^{\text{Hessian}}(\xi,\eta) &:= (f_1^{\text{Hessian}}(\xi,\eta), f_3^{\text{Hessian}}(\xi,\eta)) = \left(\frac{432C_9(\xi,\eta)^2}{C_6(\xi,\eta)^3}, \frac{C_{18}(\xi,\eta)^2}{1728\mathfrak{C}_{12}(\xi,\eta)^3}\right), \\ F_2^{\text{Hessian}}(\xi,\eta) &:= (f_2^{\text{Hessian}}(\xi,\eta), f_3^{\text{Hessian}}(\xi,\eta)) = \left(\frac{3C_{12}(\xi,\eta)}{C_6(\xi,\eta)^2}, \frac{C_{18}(\xi,\eta)^2}{1728\mathfrak{C}_{12}(\xi,\eta)^3}\right). \end{aligned}\right. \tag{7.8}$$

We have

$$(F_1^{\text{Hessian}})^{-1}(0,0) = \{(\xi,\eta)\in\mathbb{C}^2 : C_9(\xi,\eta)=0, C_{18}(\xi,\eta)=0\}.$$

$$(F_1^{\text{Hessian}})^{-1}(\infty,0) = \{(\xi,\eta)\in\mathbb{C}^2 : C_6(\xi,\eta)=0, C_{18}(\xi,\eta)=0\}.$$

$$(F_1^{\text{Hessian}})^{-1}(0,1) = \{(\xi,\eta)\in\mathbb{C}^2 : C_9(\xi,\eta)=0, C_{12}(\xi,\eta)=0\}.$$

$$(F_1^{\text{Hessian}})^{-1}(\infty,1) = \{(\xi,\eta)\in\mathbb{C}^2 : C_6(\xi,\eta)=0, C_{12}(\xi,\eta)=0\}.$$

$$(F_1^{\text{Hessian}})^{-1}(0,\infty) = \{(\xi,\eta)\in\mathbb{C}^2 : C_9(\xi,\eta)=0, \mathfrak{C}_{12}(\xi,\eta)=0\}.$$

$$(F_1^{\text{Hessian}})^{-1}(\infty,\infty) = \{(\xi,\eta)\in\mathbb{C}^2 : C_6(\xi,\eta)=0, \mathfrak{C}_{12}(\xi,\eta)=0\}.$$

$$(F_2^{\text{Hessian}})^{-1}(0,0) = \{(\xi,\eta)\in\mathbb{C}^2 : C_{12}(\xi,\eta)=0, C_{18}(\xi,\eta)=0\}.$$

$$(F_2^{\text{Hessian}})^{-1}(\infty,0) = \{(\xi,\eta)\in\mathbb{C}^2 : C_6(\xi,\eta)=0, C_{18}(\xi,\eta)=0\}.$$

$$(F_2^{\text{Hessian}})^{-1}(0,1) = \{(\xi,\eta)\in\mathbb{C}^2 : C_{12}(\xi,\eta)=0\}.$$

$$(F_2^{\text{Hessian}})^{-1}(\infty,1) = \{(\xi,\eta)\in\mathbb{C}^2 : C_6(\xi,\eta)=0, C_{12}(\xi,\eta)=0\}.$$

$$(F_2^{\text{Hessian}})^{-1}(0,\infty) = \{(\xi,\eta)\in\mathbb{C}^2 : C_{12}(\xi,\eta)=0, \mathfrak{C}_{12}(\xi,\eta)=0\}.$$

$$(F_2^{\text{Hessian}})^{-1}(\infty,\infty) = \{(\xi,\eta)\in\mathbb{C}^2 : C_6(\xi,\eta)=0, \mathfrak{C}_{12}(\xi,\eta)=0\}.$$

Hence, the cardinal numbers are given by:

$$\sharp(F_1^{\text{Hessian}})^{-1}(0,0) = 9\times 18 = 162.$$

$$\sharp(F_1^{\text{Hessian}})^{-1}(\infty, 0) = 6 \times 18 = 108.$$

$$\sharp(F_1^{\text{Hessian}})^{-1}(0, 1) = 9 \times 12 = 108.$$

$$\sharp(F_1^{\text{Hessian}})^{-1}(\infty, 1) = 6 \times 12 = 72.$$

$$\sharp(F_1^{\text{Hessian}})^{-1}(0, \infty) = 9 \times 12 = 108.$$

$$\sharp(F_1^{\text{Hessian}})^{-1}(\infty, \infty) = 6 \times 12 = 72.$$

$$\sharp(F_2^{\text{Hessian}})^{-1}(0, 0) = 12 \times 18 = 216.$$

$$\sharp(F_2^{\text{Hessian}})^{-1}(\infty, 0) = 6 \times 18 = 108.$$

$$\sharp(F_2^{\text{Hessian}})^{-1}(0, 1) = \infty.$$

$$\sharp(F_2^{\text{Hessian}})^{-1}(\infty, 1) = 6 \times 12 = 72.$$

$$\sharp(F_2^{\text{Hessian}})^{-1}(0, \infty) = 12 \times 12 = 144.$$

$$\sharp(F_2^{\text{Hessian}})^{-1}(\infty, \infty) = 6 \times 12 = 72.$$

The function $F_1^{\text{Hessian}}(\xi, \eta)$ is ramified at the points $(0, 0)$, $(\infty, 0)$, $(0, 1)$, $(\infty, 1)$, $(0, \infty)$, (∞, ∞), with degrees 162, 108, 108, 72, 108, 72, respectively. Their least common multiple is 648. The function $F_2^{\text{Hessian}}(\xi, \eta)$ is ramified at the points $(0, 0)$, $(\infty, 0)$, $(0, 1)$, $(\infty, 1)$, $(0, \infty)$, (∞, ∞), with degrees 216, 108, ∞, 72, 144, 72, respectively. Their least common multiple is 1296. Thus, the rational invariants F_1^{Hessian} and F_2^{Hessian} correspond to the Hessian groups G_{648} and G_{1296}, respectively.

Let us consider the four lines:

$$\begin{cases} l_1 : f_1^{\text{Hessian}} = 0, \\ l_2 : f_2^{\text{Hessian}} = 0, \\ l_3 : f_1^{\text{Hessian}} = \infty, f_2^{\text{Hessian}} = \infty, \\ l_4 : f_1^{\text{Hessian}} + f_2^{\text{Hessian}} = 1. \end{cases}$$

In fact,

$$l_1 = \{(\xi, \eta) \in \mathbb{C}^2 : C_9(\xi, \eta) = 0\},$$

$$l_2 = \{(\xi, \eta) \in \mathbb{C}^2 : C_{12}(\xi, \eta) = 0\},$$

$$l_3 = \{(\xi, \eta) \in \mathbb{C}^2 : C_6(\xi, \eta) = 0\},$$

$$l_4 = \{(\xi, \eta) \in \mathbb{C}^2 : C_{18}(\xi, \eta) = 0\}.$$

This leads to the geometric object:

$$\mathbb{CP}^2 - \{l_1, l_2, l_3, l_4\}.$$

Let

$$T = \begin{pmatrix} -1 & 0 & 1 \\ & -1 & 1 \\ & & 1 \end{pmatrix}, \quad S_1 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}, \quad S_2 = \begin{pmatrix} & & 1 \\ & 1 & \\ 1 & & \end{pmatrix}. \tag{7.9}$$

Then

$$T(x,y) = (1-x, 1-y). \tag{7.10}$$

$$\begin{cases} S_1(x,y) = \left(\dfrac{x}{y}, \dfrac{1}{y}\right), \\ S_1T(x,y) = \left(\dfrac{1-x}{1-y}, \dfrac{1}{1-y}\right), \\ TS_1(x,y) = \left(\dfrac{y-x}{y}, \dfrac{y-1}{y}\right), \\ S_1TS_1(x,y) = \left(\dfrac{y-x}{y-1}, \dfrac{y}{y-1}\right). \end{cases} \tag{7.11}$$

$$\begin{cases} S_2(x,y) = \left(\dfrac{1}{x}, \dfrac{y}{x}\right), \\ S_2T(x,y) = \left(\dfrac{1}{1-x}, \dfrac{1-y}{1-x}\right), \\ TS_2(x,y) = \left(\dfrac{x-1}{x}, \dfrac{x-y}{x}\right), \\ S_2TS_2(x,y) = \left(\dfrac{x}{x-1}, \dfrac{x-y}{x-1}\right). \end{cases} \tag{7.12}$$

For $(x,y) = (\infty, 0)$, we have

$$T(x,y) = (\infty, 1).$$

$$\begin{cases} S_1(x,y) = (\infty, \infty), \\ S_1T(x,y) = (\infty, 1), \\ TS_1(x,y) = (\infty, \infty), \\ S_1TS_1(x,y) = (\infty, 0). \end{cases}$$

$$\begin{cases} S_2(x,y) = (0,0), \\ S_2T(x,y) = (0,0), \\ TS_2(x,y) = (1,1), \\ S_2TS_2(x,y) = (1,1). \end{cases}$$

The orbit consists of five points:

$$\{(\infty,0),(\infty,1),(\infty,\infty),(0,0),(1,1)\}.$$

For $(x,y)=(\infty,1)$, we have

$$T(x,y) = (\infty,0).$$

$$\begin{cases} S_1(x,y) = (\infty,1), \\ S_1T(x,y) = (\infty,\infty), \\ TS_1(x,y) = (\infty,0), \\ S_1TS_1(x,y) = (\infty,\infty). \end{cases}$$

$$\begin{cases} S_2(x,y) = (0,0), \\ S_2T(x,y) = (0,0), \\ TS_2(x,y) = (1,1), \\ S_2TS_2(x,y) = (1,1). \end{cases}$$

The orbit consists of five points:

$$\{(\infty,1),(\infty,0),(\infty,\infty),(0,0),(1,1)\}.$$

For $(x,y)=(0,1)$, we have

$$T(x,y) = (1,0).$$

$$\begin{cases} S_1(x,y) = (0,1), \\ S_1T(x,y) = (\infty,\infty), \\ TS_1(x,y) = (1,0), \\ S_1TS_1(x,y) = (\infty,\infty). \end{cases}$$

$$\begin{cases} S_2(x,y) = (\infty,\infty), \\ S_2T(x,y) = (1,0), \\ TS_2(x,y) = (\infty,\infty), \\ S_2TS_2(x,y) = (0,1). \end{cases}$$

The orbit consists of three points:

$$\{(0,1),(1,0),(\infty,\infty)\}.$$

According to [Y], let

$$G=\langle T_1,T_2,S,U_1,U_2\rangle \tag{7.13}$$

where

$$T_1=\begin{pmatrix}1&1&-\omega\\&1&1\\&&1\end{pmatrix},\quad T_2=\begin{pmatrix}1&\omega&-\omega\\&1&\overline{\omega}\\&&1\end{pmatrix}, \tag{7.14}$$

$$S=-\overline{\omega}J,\quad J=\begin{pmatrix}&&1\\&-1&\\1&&\end{pmatrix}, \tag{7.15}$$

$$U_1=\begin{pmatrix}1&&\\&-\omega&\\&&1\end{pmatrix},\quad U_2=\begin{pmatrix}-1&&\\&-\omega&\\&&-1\end{pmatrix}. \tag{7.16}$$

We find that
(7.17)

$$\begin{aligned}G&\cong\langle S,ST_1,ST_2,SU_1,SU_2:S^2=(ST_1)^4=(ST_2)^4=\omega I,(SU_1)^6=(SU_2)^6=I\rangle\\&\cong\langle a_1,a_2,a_3,a_4,a_5:a_1^2=a_2^4=a_3^4=1,a_4^6=a_5^6=1\rangle\\&\cong\mathbb{Z}_2*\mathbb{Z}_4*\mathbb{Z}_4*\mathbb{Z}_6*\mathbb{Z}_6.\end{aligned}$$

Note that

$$\begin{cases}\det(\lambda I-S)=(\lambda+\overline{\omega})(\lambda-\overline{\omega})^2,\\ \det(\lambda I-ST_1)=(\lambda+\omega)(\lambda^2+\overline{\omega}),\\ \det(\lambda I-ST_2)=(\lambda+\omega)(\lambda^2+\overline{\omega}),\\ \det(\lambda I-SU_1)=(\lambda+1)(\lambda+\overline{\omega})(\lambda-\overline{\omega}),\\ \det(\lambda I-SU_2)=(\lambda+1)(\lambda+\overline{\omega})(\lambda-\overline{\omega}).\end{cases}$$

One of the most interesting problems in algebraic geometry is to obtain a generalization of the theory of elliptic modular functions to the case of higher genus. We know that in the elliptic case the invariant j plays a fundamental role. Igusa (see [I]) studied the j invariants for the case of genus two, i.e. the hyperelliptic curves. He gave the invariants J_2, J_4, J_6, J_8 and J_{10}.

For the Picard curve

$$y^3 = x(x-1)(x-k_1)(x-k_2), \tag{7.18}$$

put

$$\begin{cases} J_1 = J_1(k_1, k_2) = \dfrac{k_2^2(k_2-1)^2}{k_1^2(k_1-1)^2(k_1-k_2)^2}, \\ J_2 = J_2(k_1, k_2) = \dfrac{k_1^2(k_1-1)^2}{k_2^2(k_2-1)^2(k_2-k_1)^2}. \end{cases} \tag{7.19}$$

Let us study the following six Picard curves:

$$\begin{cases} y^3 = x(x-1)(x-k_1)(x-k_2), \\ y^3 = x(x-1)(x-k_2)(x-k_3), \\ y^3 = x(x-1)(x-k_3)(x-k_1), \\ y^3 = x(x-1)(x-k_0)(x-k_1), \\ y^3 = x(x-1)(x-k_0)(x-k_2), \\ y^3 = x(x-1)(x-k_0)(x-k_3), \end{cases} \tag{7.20}$$

with $k_0 = k_0(z_1, z_2)$, $k_1 = k_1(z_1, z_2)$, $k_2 = k_2(z_1, z_2)$ and $k_3 = k_3(z_1, z_2)$ (see [Y]) where $(z_1, z_2) \in \mathfrak{S}_2 = \{(z_1, z_2) \in \mathbb{C}^2 : z_1 + \overline{z_1} - z_2\overline{z_2} > 0\}$. We are interested in the next six maps:

$$\begin{cases} (J_1(k_1(z_1,z_2), k_2(z_1,z_2)), J_2(k_1(z_1,z_2), k_2(z_1,z_2))) : \mathfrak{S}_2 \to \mathbb{CP}^2, \\ (J_1(k_2(z_1,z_2), k_3(z_1,z_2)), J_2(k_2(z_1,z_2), k_3(z_1,z_2))) : \mathfrak{S}_2 \to \mathbb{CP}^2, \\ (J_1(k_3(z_1,z_2), k_1(z_1,z_2)), J_2(k_3(z_1,z_2), k_1(z_1,z_2))) : \mathfrak{S}_2 \to \mathbb{CP}^2, \\ (J_1(k_0(z_1,z_2), k_1(z_1,z_2)), J_2(k_0(z_1,z_2), k_1(z_1,z_2))) : \mathfrak{S}_2 \to \mathbb{CP}^2, \\ (J_1(k_0(z_1,z_2), k_2(z_1,z_2)), J_2(k_0(z_1,z_2), k_2(z_1,z_2))) : \mathfrak{S}_2 \to \mathbb{CP}^2, \\ (J_1(k_0(z_1,z_2), k_3(z_1,z_2)), J_2(k_0(z_1,z_2), k_3(z_1,z_2))) : \mathfrak{S}_2 \to \mathbb{CP}^2. \end{cases} \tag{7.21}$$

In [Y], we find that

(1) Picard curves and Picard integrals

$$s_1\left(\frac{2}{3}, \frac{2}{3}, -\frac{1}{3}; \lambda_1, \lambda_2\right), s_2\left(\frac{2}{3}, \frac{2}{3}, -\frac{1}{3}; \lambda_1, \lambda_2\right).$$

(2) Picard modular functions

$$s_1\left(\frac{3}{4}, \frac{1}{2}, -\frac{1}{4}; J_1, J_2\right), s_2\left(\frac{3}{4}, \frac{1}{2}, -\frac{1}{4}; J_1, J_2\right).$$

Let us recall the following facts:

(1) $GL(2)$: $\mathbb{H}/SL(2,\mathbb{Z})(2) \cong \mathbb{P}^1 - \{0,1,\infty\}$.

(2) $GL(3)$: $\mathfrak{S}_2/U(2,1;\mathcal{O}_K)(1-\omega) \cong \mathbb{P}^2 - \{\text{four points}\}$.

This leads us to study the special functions on the configuration

$$\begin{aligned}&\mathbb{P}^2 - \{\text{four points}\}\\ \cong&\mathbb{P}^2 - \{(1,0),(0,1),(\infty,\infty),(0,0)\} \cong \mathbb{P}^2 - \{(1,0),(0,1),(\infty,\infty),(1,1)\}.\end{aligned}$$

In fact, the fundamental modular function J (Hauptmodul) was first constructed by Dedekind in 1877 and independently by Klein one year later. In our case, we constructed two fundamental modular functions J_1 and J_2.

(1) $GL(2)$ (Dedekind, Klein):

$$J(1-J)\frac{d^2z}{dJ^2} + \left(\frac{2}{3} - \frac{7}{6}J\right)\frac{dz}{dJ} - \frac{1}{144}z = 0.$$

A solution is given by

$$z = F\left(\frac{1}{12}, \frac{1}{12}; \frac{2}{3}; J\right).$$

(2) $GL(3)$ (see [Y]):

(7.22)

$$\begin{cases} J_1(1-J_1)\dfrac{\partial^2 z}{\partial J_1^2} + J_2(1-J_1)\dfrac{\partial^2 z}{\partial J_1 \partial J_2} + \left(1 - \dfrac{3}{2}J_1\right)\dfrac{\partial z}{\partial J_1} - \dfrac{1}{4}J_2\dfrac{\partial z}{\partial J_2} - \dfrac{1}{16}z = 0, \\ J_2(1-J_2)\dfrac{\partial^2 z}{\partial J_2^2} + J_1(1-J_2)\dfrac{\partial^2 z}{\partial J_1 \partial J_2} + \left(1 - \dfrac{3}{2}J_2\right)\dfrac{\partial z}{\partial J_2} - \dfrac{1}{4}J_1\dfrac{\partial z}{\partial J_1} - \dfrac{1}{16}z = 0. \end{cases}$$

A solution is given by

$$z = F_1\left(\frac{1}{4}; \frac{1}{4}, \frac{1}{4}; 1; J_1, J_2\right). \tag{7.23}$$

The following five values are interesting:

$$(J_1, J_2) = (1,0), \quad (0,1), \quad (\infty,\infty), \quad (0,0) \quad \text{or} \quad (1,1).$$

Department of Mathematics, Peking University
Beijing 100871, P. R. China
E-mail address: yanglei@math.pku.edu.cn

References

[AK] P. Appell and J. Kampé de Fériet, *Fonctions Hypergéométriques et Hypersphériques, Polynomes d'Hermite*, Gauthier-Villars, 1926.

[At] M. F. Atiyah, The icosahedron, Math. Medley **18** (1990), 1-12.

[AtS] M. F. Atiyah and P. Sutcliffe, Polyhedra in physics, chemistry and geometry, math-ph/0303071.

[Be] G. V. Belyĭ, On Galois extensions of a maximal cyclotomic field, Math. USSR Izv. **14** (1980), 247-256.

[Bu1] H. Burkhardt, Untersuchungen aus dem Gebiete der hyperelliptischen Modulfunctionen, II, Math. Ann. **38** (1891), 161-224.

[Bu2] H. Burkhardt, Untersuchungen aus dem Gebiete der hyperelliptischen Modulfunctionen, III, Math. Ann. **41** (1893), 313-343.

[Cob1] A. B. Coble, An application of the form-problems associated with certain Cremona groups to the solution of equations of higher degree, Trans. Amer. Math. Soc. **9** (1908), 396-424.

[Cob2] A. B. Coble, An application of Moore's cross-ratio group to the solution of the sextic equation, Trans. Amer. Math. Soc. **12** (1911), 311-325.

[Cob3] A. B. Coble, Point sets and allied Cremona groups, I, Trans. Amer. Math. Soc. **16** (1915), 155-198.

[Cob4] A. B. Coble, Point sets and allied Cremona groups, II, Trans. Amer. Math. Soc. **17** (1916), 345-385.

[Cob5] A. B. Coble, Point sets and allied Cremona groups, III, Trans. Amer. Math. Soc. **18** (1917), 331-372.

[C] A. M. Cohen, Finite complex reflection groups, Ann. Sci. Éc. Norm. Sup. 4^{e} série **9** (1976), 379-436.

[CW1] P. B. Cohen and J. Wolfart, Algebraic Appell-Lauricella functions, Analysis **12** (1992), 359-376.

[CW2] P. B. Cohen and J. Wolfart, Fonctions hypergéométriques en plusieurs variables et espaces des modules de variétés abéliennes, Ann. Sci. Éc. Norm. Sup. **26** (1993), 665-690.

[CC] J. H. Conway, R. T. Curtis, S. P. Norton, R. A. Parker and R. A. Wilson, *Atlas of Finite Groups, Maximal Subgroups and Ordinary Characters for Simple Groups*, Clarendon Press, Oxford, 1985.

[CG] J.-M. Couveignes and L. Granboulan, Dessins from a geometric point of view, In: *The Grothendieck Theory of Dessins d'Enfants*, Edited by L. Schneps, 79-113, London Math. Soc. Lecture Note Series **200**, Cambridge University Press, 1994.

[Co1] H. S. M. Coxeter, *Regular Polytopes*, Second Edition, The MacMillan Company, New York, 1963.

[Co2] H. S. M. Coxeter, *Regular Complex Polytopes*, Cambridge University Press, 1974.

[Co3] H. S. M. Coxeter, Finite groups generated by unitary reflections, Abh. Math. Sem. Univ. Hamburg **31** (1967), 125-135.

[DM1] P. Deligne and G. D. Mostow, Monodromy of hypergeometric functions and non-lattice integral monodromy, Publ. Math. I.H.E.S. **63** (1986), 5-89.

[DM2] P. Deligne and G. D. Mostow, *Commensurabilities among Lattices in* $PU(1,n)$, Annals of Math. Studies **132**, Princeton University Press, 1993.

[F] R. Fricke, Lehrbuch der Algebra, Vol. II, Braunschweig, 1926.

[G] J. J. Gray, *Linear Differential Equations and Group Theory from Riemann to Poincaré*, Second Edition, Birkhäuser, 2000.

[Gro] A. Grothendieck, Esquisse d'un programme, In: *Geometric Galois Actions, 1. Around Grothendieck's Esquisse d'un Programme*, Edited by L. Schneps and P. Lochak, 5-48, London Math. Soc. Lecture Note Series **242**, Cambridge University Press, 1997.

[He] C. Hermite, *Œuvres de Charles Hermite*, Vol. II, E. Picard Ed., Gauthier-Villars, 1908.

[H] D. Hilbert, Mathematical problems, Bull. Amer. Math. Soc. **8** (1902), 437-479.

[Hi] F. Hirzebruch, The ring of Hilbert modular forms for real quadratic fields of small discriminant, In: *Modular Functions of One Variable VI, Proceedings, Bonn 1976*, Edited by J. P. Serre and D. B. Zagier, 287-323, Lecture Notes in Math. **627**, Springer-Verlag, 1977.

- [Ho] R. P. Holzapfel, *Geometry and arithmetic around Euler partial differential equations*, Dt. Verl. d. Wiss., Berlin/Reidel, Dordrecht 1986.
- [Hu] B. Hunt, *The Geometry of some special Arithmetic Quotients*, Lecture Notes in Math. **1637**, Springer-Verlag, 1996.
- [I] J. Igusa, Arithmetic variety of moduli for genus two, Ann. of Math. **72** (1960), 612-649.
- [KlE] F. Klein, Vergleichende Betrachtungen über neuere geometrische Forschungen (Das Erlanger Programm), Programm zum Eintritt in die philosophische Fakultät und den Senat der k. Friedrich-Alexanders-Universität zu Erlangen, Erlangen, A. Deichert, 1872, In: *Gesammelte Mathematische Abhandlungen*, Bd. I, Springer-Verlag, Berlin, 1921, 460-497.
- [Kl] F. Klein, *Gesammelte Mathematische Abhandlungen*, Bd. II, Springer-Verlag, Berlin, 1922.
- [Kl1] F. Klein, Über binäre Formen mit linearen Transformationen in sich selbst, Math. Ann. **9** (1875), In: *Gesammelte Mathematische Abhandlungen*, Bd. II, Springer-Verlag, Berlin, 1922, 275-301.
- [Kl2] F. Klein, Über (algebraisch integrierbare) lineare Differentialgleichungen (Erster Aufsatz), Math. Ann. **11** (1876), In: *Gesammelte Mathematische Abhandlungen*, Bd. II, Springer-Verlag, Berlin, 1922, 302-306.
- [Kl3] F. Klein, Über (algebraisch integrierbare) lineare Differentialgleichungen (Zweiter Aufsatz), Math. Ann. **12** (1877), In: *Gesammelte Mathematische Abhandlungen*, Bd. II, Springer-Verlag, Berlin, 1922, 307-320.
- [Kl4] F. Klein, Weitere Untersuchungen über das Ikosaeder, Math. Ann. **12** (1877), In: *Gesammelte Mathematische Abhandlungen*, Bd. II, Springer-Verlag, Berlin, 1922, 321-384.
- [Kl5] F. Klein, Über die Auflösung gewisser Gleichungen vom siebenten und achten Grade, Math. Ann. **15** (1879), In: *Gesammelte Mathematische Abhandlungen*, Bd. II, Springer-Verlag, Berlin, 1922, 390-428.
- [Kl6] F. Klein, Zur Theorie der allgemeinen Gleichungen sechsten und siebenten Grades, Math. Ann. **28** (1886/87), In: *Gesammelte Mathematische Abhandlungen*, Bd. II, Springer-Verlag, Berlin, 1922, 439-472.
- [Kl7] F. Klein, Sur la résolution, par les fonctions hyperelliptiques, de l'équation du vingt-septième degré, de laquelle dépend la détermination des vingt-sept droites d'une surface cubique, J. de Math. pure et appl. **4** (1888), In: *Gesammelte Mathematische Abhandlungen*, Bd. II, Springer-Verlag, Berlin, 1922, 473-479.
- [Kl8] F. Klein, Über die Auflösung der allgemeinen Gleichungen fünften und sechsten Grades, Math. Ann. **61** (1905), In: *Gesammelte Mathematische Abhandlungen*, Bd. II, Springer-Verlag, Berlin, 1922, 481-504.
- [Kl9] F. Klein, *Lectures on the Icosahedron and the Solution of Equations of the Fifth Degree*, Translated by G. G. Morrice, second and revised edition, Dover Publications, Inc., 1956.
- [Kl10] F. Klein, *Vorlesungen über die Entwicklung der Mathematik im 19. Jahrhundert*, Berlin, 1926.
- [KlS] F. Klein, *Vorlesungen über das Ikosaeder und die Auflösung der Gleichungen vom fünften Grade*, Reprint of the 1884 original. Edited, with an introduction and commentary by Peter Slodowy, Birkhäuser Verlag, Basel; B. G. Teubner, Stuttgart, 1993.
- [Kl11] F. Klein, Ueber die Transformation der elliptischen Functionen und die Auflösung der Gleichungen fünften Grades, Math. Ann. **14** (1879), 111-172.
- [Kl12] F. Klein, Ueber die Transformation siebenter Ordnung der elliptischen Functionen, Math. Ann. **14** (1879), 428-471.
- [L1] R. P. Langlands, Problems in the theory of automorphic forms, In: *Lectures in modern analysis and applications*, III, 18-61, Lecture Notes in Math. **170**, Springer, Berlin, 1970.
- [L2] R. P. Langlands, Representation theory: its rise and its role in number theory, In: *Proceedings of the Gibbs Symposium* (New Haven, CT, 1989), 181-210, Amer. Math. Soc., Providence, RI, 1990.
- [MZ] N. Magot and A. Zvonkin, Belyi functions for Archimedean solids, Discrete Math. **217** (2000), 249-271.

[Ma] Yu. I. Manin, *Cubic Forms, Algebra, Geometry, Arithmetic*, Translated from Russian by M. Hazewinkel, Second Edition, North-Holland, 1986.

[Mas1] H. Maschke, Ueber die quaternäre, endliche, lineare Substitutionsgruppe der Borchardt'schen Moduln, Math. Ann. **30** (1887), 496-515.

[Mas2] H. Maschke, Aufstellung des vollen Formensystems einer quaternären Gruppe von 51840 linearen Substitutionen, Math. Ann. **33** (1889), 317-344.

[MBD] G. A. Miller, H. F. Blichfeldt and L. E. Dickson, *Theory and Applications of Finite Groups*, Wiley, New York, 1916.

[M] G. D. Mostow, Generalized Picard lattices arising from half-integral conditions, Publ. Math. I.H.E.S. **63** (1986), 91-106.

[Mu] D. Mumford, *Tata Lectures on Theta II, Jacobian theta functions and differential equations*, Progress in Math. Vol. **43**, Birkhäuser, 1984.

[Se] J.-P. Serre, Extensions icosaédriques, Séminaire de Théorie des Nombres de Bordeaux 1979/80, No. 19, In: *Œuvres Collected Papers*, Vol. III, 1972-1984, 550-554, Springer-Verlag, 1986.

[ST] C. C. Shephard and J. A. Todd, Finite unitary reflection groups, Canad. J. Math. **6** (1954), 274-304.

[Sl] P. Slodowy, Das Ikosaeder und die Gleichungen fünften Grades, In: *Arithmetik und Geometrie*, 71-113, Math. Miniaturen, **3**, Birkhäuser, 1986.

[U] H. Umemura, Resolution of algebraic equations by theta constants, In: D. Mumford, *Tata Lectures on Theta II*, Progress in Math. Vol. **43**, Birkhäuser, 1984, 261-272.

[We] H. Weber, *Lehrbuch der Algebra*, Vol. II, Chelsea Publishing Company, New York, 1961.

[W] H. Weyl, *Symmetry*, Princeton University Press, 1952.

[Wi] A. Witting, Ueber Jacobi'sche Functionen k^{ter} Ordnung zweier Variabler, Math. Ann. **29** (1887), 157-170.

[Y] L. Yang, Geometry and arithmetic associated to Appell hypergeometric partial differential equations, math.NT/0309415.